

Assessment of trace elements distribution in French agricultural soils

Agathe Herbout, Romain Armand, Raul E. Martinez, Davide Rizzo, Olivier

Pourret

► To cite this version:

Agathe Herbout, Romain Armand, Raul E. Martinez, Davide Rizzo, Olivier Pourret. Assessment of trace elements distribution in French agricultural soils. Goldschmidt, Aug 2019, Barcelona, Spain. 10.5281/zenodo.3249218 . hal-03154002

HAL Id: hal-03154002 https://inria.hal.science/hal-03154002v1

Submitted on 13 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Assessment of trace elements distribution in French agricultural soils

AGATHE HERBOUT¹, ROMAIN ARMAND¹, RAUL E. MARTINEZ², DAVIDE RIZZO³, OLIVIER POURRET¹

 ¹AGHYLE, UniLaSalle, Beauvais, France; olivier.pourret@unilasalle.fr
²Max Planck Institute for Biogeochemistry, Jena, Germany
³INTERACT, UniLaSalle, Beauvais, France

Although trace element distribution and mobility have been investigated in agricultural soils, precise information about the extent to which anthropogenic activities influence soil quality remains limited. This study quantified trace element distribution (As, Bi, Cd, Cu, Ni, Pb, Pd, Sb, Zn) in agricultural and grazing land soils in metropolitan France based on the results from the GEMAS project [1]. The dataset is composed of214 samples from agricultural soils (Ap) and 218 samples from grazing land soils (Gr) for total and mobile fractions. The geological substrates have been divided into five main classes to correlate with geology: (i) shales, (ii) sandstones and sands, (iii) carbonate rocks, (iv) metamorphic rocks, and (v) intrusive and igneous rocks. These nine metal elements were selected using the enrichment factor higher than 40 in most of the five rock classes, specifically As and Cd. Arsenic had a median of 9.04 mg/kg and 9.5 mg/kg for Ap and Gr, respectively, on all kind of bedrocks. These values are similar, and this is also the case for Cd : 0.215 mg/kg and 0.249 mg/kg for Ap and Gr, soils respectively. Kriging interpolation was used to analyze the spatial distribution of the both elements in French soils. A correlation between geology and mining was observed for the distribution of As in France, for example southern part of Massif Central [2]. However, Cd shows a different pattern. The highest concentrations are situated in the sedimentary basin (Aquitaine and Parisien), as well as in the South Massif Central and South Vosges. This could be related to anthropogenic activities [3]. If the high concentration level of As is not due to anthropisation, its distribution could be used as a reference for pollution in France. The concentrations differ depending on the bedrock. On the contrary, Cd concentrations are not correlated to geology, but high Cd concentrations are linked with cultural practices. The lowest level Cd concentration could only be used for a reference.

[1] Reinman et al. (2014) Chemistry of Europe's agricultural soils [2] Birke et al. (2017) J Geochem Explor 173, 13-30 [3] Tarvainen et al. (2013) Appl Geochem 28, 2-10.