



HAL
open science

A reduced parallel transport equation on Lie Groups with a left-invariant metric

Nicolas Guigui, Xavier Pennec

► **To cite this version:**

Nicolas Guigui, Xavier Pennec. A reduced parallel transport equation on Lie Groups with a left-invariant metric. 2021. hal-03154318v1

HAL Id: hal-03154318

<https://inria.hal.science/hal-03154318v1>

Preprint submitted on 1 Mar 2021 (v1), last revised 2 Mar 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A reduced parallel transport equation on Lie Groups with a left-invariant metric

Nicolas Guigui^[0000–0002–7901–0732] and Xavier Pennec

Universit Cte d’Azur, Inria Epione project team, France

Abstract. This paper presents a derivation of the parallel transport equation expressed in the Lie algebra of a Lie group endowed with a left-invariant metric. The use of this equation is exemplified on the group of rigid body motions $SE(3)$, using basic numerical integration schemes, and compared to the pole ladder algorithm. This results in a stable and efficient implementation of parallel transport. The implementation leverages the python package [geomstats](#) and is available online.

Keywords: Parallel transport · Lie Groups

1 Introduction

Lie groups are ubiquitous in geometry, physics and many application domains such as robotics [?], medical imaging [?] or computer vision [?], giving rise to a prolific research avenue. Structure preserving numerical methods have demonstrated significant qualitative and quantitative improvements over extrinsic methods [?]. Moreover, machine learning [?] and optimisation methods [?] are being developed to deal with Lie group data.

In this context, parallel transport is a natural tool to define statistical models and optimisation procedures, such as the geodesic or spline regression [?,?], or to normalise data represented by tangent vectors [?,?].

Different geometric structures are compatible with the group structure, such as its canonical Cartan connection, whose geodesics are one-parameter subgroups, or left-invariant Riemannian metrics. In this work we focus on the latter case, that is fundamental in geometric mechanics [?] and has been studied in depth since the foundational papers of Arnold [?] and Milnor [?]. The fundamental idea of Euler-Poincaré reduction is that the geodesic equation can be expressed entirely in the Lie algebra thanks to the symmetry of left-invariance [?], alleviating the burden of coordinate charts.

However, to the best of our knowledge, there is no literature on a similar treatment of the parallel transport equation. We present here a derivation of the parallel transport equation expressed in the Lie algebra of a Lie group endowed with a left-invariant metric. We exemplify the use of this equation on the group of rigid body motions $SE(3)$, using common numerical integration schemes, and compare it to the pole ladder approximation algorithm. This results in a stable and efficient implementation of parallel transport. The implementation leverages the python package [geomstats](#) and is available online at <http://geomstats.ai>.

In section 2, we give the general notations and recall some basic facts from Lie group theory. Then we derive algebraic expressions of the Levi-Civita connection associated to the left-invariant metric in section 3. The equation of parallel transport is deduced from this expression and its integration is exemplified in section 4.

2 Notations

Let G be a lie group of (finite) dimension n . Let e be its identity element, $\mathfrak{g} = T_e G$ be its tangent space at e , and for any $g \in G$, let $L_g : h \in G \mapsto gh$ denote the left-translation map, and dL_g its differential map. Let \mathfrak{g}^L be the Lie algebra of left-invariant vector fields of G : $X \in \mathfrak{g}^L \iff \forall g \in G, X|_g = dL_g X_e$.

\mathfrak{g} and \mathfrak{g}^L are in one-to-one correspondence, and we will write \tilde{x} the left-invariant field generated by $x \in \mathfrak{g}$: $\forall g \in G, \tilde{x}_g = dL_g x$. The bracket defined on \mathfrak{g} by $[x, y] = [\tilde{x}, \tilde{y}]_e$ turns \mathfrak{g} into a Lie algebra that is isomorphic to \mathfrak{g}^L . One can also check that this bracket coincides with the adjoint map defined by $\text{ad}_x(y) = d_e(g \mapsto \text{Ad}_g y)$, where $\text{Ad}_g = d_e(h \mapsto ghg^{-1})$. For a matrix group, it is the commutator.

Let (e_1, \dots, e_n) be an orthonormal basis of \mathfrak{g} , and the associated left-invariant vector fields $X_i^L = \tilde{e}_i = g \mapsto dL_g e_i$. As dL_g is an isomorphism, $(X_1^L|_g, \dots, X_n^L|_g)$ form a basis of $T_g G$ for any $g \in G$, so one can write $X|_g = f^i(g) X_i^L|_g$ where for $i = 1, \dots, n$, $g \mapsto f^i(g)$ is a smooth real-valued function on G . Any vector field on G can thus be expressed as a linear combination of the X_i^L with function coefficients.

Finally, let θ be the Maurer-Cartan form defined on G by:

$$\forall g \in G, \forall v \in T_g G, \theta|_g(v) = (dL_g)^{-1} v \in \mathfrak{g} \quad (1)$$

It is a \mathfrak{g} -valued 1-form and for a vector field X on G we write $\theta(X)|_g = \theta|_g(X|_g)$ to simplify the notations.

3 Left-invariant metric and connection

A Riemannian metric $\langle \cdot, \cdot \rangle$ on G is called left-invariant if the differential map of the left translation is an isometry between tangent spaces, that is

$$\forall g, h \in G, \forall u, v \in T_g G, \langle u, v \rangle_g = \langle dL_h u, dL_h v \rangle_{hg}.$$

It is thus uniquely determined by an inner product on the tangent space at the identity $T_e G = \mathfrak{g}$ of G . Furthermore, the metric dual to the adjoint map is defined such that

$$\forall a, b, c \in \mathfrak{g}, \langle \text{ad}_a^*(b), c \rangle = \langle b, \text{ad}_a(c) \rangle = \langle [a, c], b \rangle. \quad (2)$$

As the bracket can be computed explicitly in the Lie algebra, so can ad^* thanks to the orthonormal basis of \mathfrak{g} . Now let ∇ be the Levi-Civita connection associated

to the metric. It is also left-invariant and can be characterised by a bi-linear form on \mathfrak{g} that verifies [?,?]:

$$\forall x, y \in \mathfrak{g}, \quad \alpha(x, y) := (\nabla_{\tilde{x}} \tilde{y})_e = \frac{1}{2}([x, y] - \text{ad}_x^*(y) - \text{ad}_y^*(x)) \quad (3)$$

Indeed by the left-invariance, for two left-invariant vector fields $X = \tilde{x}, Y = \tilde{y} \in \mathfrak{g}^L$, the map $g \mapsto \langle X, Y \rangle_g$ is constant, so for any vector field $Z = \tilde{z}$ we have $Z(\langle X, Y \rangle) = 0$. Kozsul formula thus becomes

$$\begin{aligned} 2\langle \nabla_X Y, Z \rangle &= \langle [X, Y], Z \rangle - \langle [Y, Z], X \rangle - \langle [X, Z], Y \rangle \\ 2\langle \nabla_X Y, Z \rangle_e &= \langle [x, y], z \rangle_e - \langle \text{ad}_y(z), x \rangle_e - \langle \text{ad}_x(z), y \rangle_e \\ 2\langle \alpha(x, y), z \rangle_e &= \langle [x, y], z \rangle_e - \langle \text{ad}_y^*(x), z \rangle_e - \langle \text{ad}_x^*(y), z \rangle_e. \end{aligned} \quad (4)$$

Note however that this formula is only valid for left-invariant vector fields. We will now generalise to any vector fields defined along a smooth curve on G , using the left-invariant basis (X_1^L, \dots, X_n^L) .

Let $\gamma : [0, 1] \rightarrow G$ be a smooth curve, and Y a vector field defined along γ . Write $Y = g^i X_i^L$, $\dot{\gamma} = f^i X_i^L$. Let's also define the *left-angular velocities* $\omega(t) = \theta|_{\gamma(t)} \dot{\gamma}(t) = (f^i \circ \gamma)(t) e_i \in \mathfrak{g}$ and $\zeta(t) = \theta(Y)|_{\gamma(t)} = (g^j \circ \gamma)(t) e_j \in \mathfrak{g}$. Then the covariant derivative of Y along γ is

$$\begin{aligned} \nabla_{\dot{\gamma}(t)} Y &= (f^i \circ \gamma)(t) \nabla_{X_i^L} (g^j X_j^L) \\ &= (f^i \circ \gamma)(t) X_i^L (g^j) X_j^L + (f^i \circ \gamma)(t) (g^j \circ \gamma)(t) (\nabla_{X_i^L} X_j^L)|_{\gamma(t)} \\ dL_{\gamma(t)}^{-1} \nabla_{\dot{\gamma}(t)} Y &= (f^i \circ \gamma)(t) X_i^L (g^j) e_j + (f^i \circ \gamma)(t) (g^j \circ \gamma)(t) dL_{\gamma(t)}^{-1} (\nabla_{X_i^L} X_j^L)|_{\gamma(t)} \\ &= (f^i \circ \gamma)(t) X_i^L (g^j) e_j + (f^i \circ \gamma)(t) (g^j \circ \gamma)(t) \nabla_{e_i} e_j \end{aligned}$$

where Leibniz formula and the invariance of the connection is used in $(\nabla_{X_i^L} X_j^L) = dL_{\gamma(t)} \nabla_{e_i} e_j$. Therefore for $k = 1..n$

$$\begin{aligned} \langle dL_{\gamma(t)}^{-1} \nabla_{\dot{\gamma}(t)} Y, e_k \rangle &= (f^i \circ \gamma)(t) X_i^L (g^j) \langle e_j, e_k \rangle \\ &\quad + (f^i \circ \gamma)(t) (g^j \circ \gamma)(t) \langle \nabla_{e_i} e_j, e_k \rangle \end{aligned} \quad (5)$$

but on one hand

$$\begin{aligned} \zeta(t) &= \theta(Y)|_{\gamma(t)} = \theta|_{\gamma(t)} ((g^j \circ \gamma)(t) X_j^L|_{\gamma(t)}) \\ &= (g^j \circ \gamma)(t) e_j \end{aligned} \quad (6)$$

$$\begin{aligned} \dot{\zeta}(t) &= (g^j \circ \gamma)'(t) e_j = d_{\gamma(t)} g^j \dot{\gamma}(t) e_j \\ &= d_{\gamma(t)} g^j \left((f^i \circ \gamma)(t) X_i^L|_{\gamma(t)} \right) e_j \\ &= (f^i \circ \gamma)(t) d_{\gamma(t)} g^j X_i^L|_{\gamma(t)} e_j \\ &= (f^i \circ \gamma)(t) X_i^L (g^j) e_j \end{aligned} \quad (7)$$

and on the other hand, using (4):

$$\begin{aligned}
(f^i \circ \gamma)(g^j \circ \gamma) \langle \nabla_{e_i} e_j, e_k \rangle &= \frac{1}{2} (f^i \circ \gamma)(g^j \circ \gamma) (\langle [e_i, e_j], e_k \rangle \\
&\quad - \langle [e_j, e_k], e_i \rangle - \langle [e_i, e_k], e_j \rangle) \\
&= \frac{1}{2} (\langle [(f^i \circ \gamma)e_i, (g^j \circ \gamma)e_j], e_k \rangle \\
&\quad - \langle [(g^j \circ \gamma)e_j, e_k], (f^i \circ \gamma)e_i \rangle \\
&\quad - \langle [(f^i \circ \gamma)e_i, e_k], (g^j \circ \gamma)e_j \rangle) \\
&= \frac{1}{2} ([\omega, \zeta] - \text{ad}_\omega^* \zeta - \text{ad}_\zeta^* \omega) = \alpha(\omega, \zeta) \quad (8)
\end{aligned}$$

Thus, we obtain an algebraic expression for the covariant derivative of any vector field Y along a smooth curve γ . It will be the main ingredient of this paper.

$$dL_{\gamma(t)}^{-1} \nabla_{\dot{\gamma}(t)} Y(t) = \dot{\zeta}(t) + \alpha(\omega(t), \zeta(t)) \quad (9)$$

A similar expression can be found in [?,?]. As all the variables of the right-hand side are defined in \mathfrak{g} , they can be computed with matrix operations and an orthonormal basis.

4 Parallel Transport

We now focus on two particular cases of (9) to derive the equations of geodesics and of parallel transport along a curve.

4.1 Geodesic equation

The first particular case is for $Y(t) = \dot{\gamma}(t)$. It is then straightforward to deduce from (9) the Euler-Poincaré equation for a geodesic curve [?,?]. Indeed in this case, recall that $\omega = \theta|_{\gamma(t)} \dot{\gamma}(t)$ is the left-angular velocity, $\zeta = \omega$ and $\alpha(\omega, \omega) = -\text{ad}_\omega^* \omega$. Hence γ is a geodesic if and only if $dL_{\gamma(t)}^{-1} \nabla_{\dot{\gamma}(t)} \dot{\gamma}(t) = 0$ i.e. setting the left-hand side of (9) to 0. We obtain

$$\begin{cases} \dot{\gamma}(t) &= dL_{\gamma(t)} \omega(t) \\ \dot{\omega}(t) &= \text{ad}_{\omega(t)}^* \omega(t). \end{cases} \quad (10)$$

Remark 1. One can show that the metric is bi-invariant if and only if the adjoint map is skew-symmetric (see [?] or [?, Prop. 20.7]). In this case $\text{ad}_\omega^* \omega = 0$ and (10) coincides with the equation of one-parameter subgroups on G .

4.2 Reduced Parallel Transport Equation

The second case is for a vector Y that is parallel along the curve γ , that is, $\forall t, \nabla_{\dot{\gamma}(t)} Y(t) = 0$. Similarly to the geodesic equation, we deduce from (9) the parallel transport equation expressed in the Lie algebra.

Theorem 1. *Let γ be a smooth curve on G . The vector Y is parallel along γ if and only if it is solution to*

$$\begin{cases} \omega(t) &= dL_{\gamma(t)}^{-1} \dot{\gamma}(t) \\ Y(t) &= dL_{\gamma(t)} \zeta(t) \\ \dot{\zeta}(t) &= -\alpha(\omega(t), \zeta(t)) \end{cases} \quad (11)$$

Note that in order to parallel transport along a geodesic curve, (10) and (11) are solved jointly.

4.3 Application

We now exemplify Theorem 1 on the group of isometries of \mathbb{R}^3 , $SE(3)$, endowed with a left-invariant metric g . $SE(3)$, is the semi-direct product of the group of three-dimensional rotations $SO(3)$ with \mathbb{R}^3 , i.e. the group multiplicative law for $R, R' \in SO(3), t, t' \in \mathbb{R}^3$ is given by

$$(R, t) \cdot (R', t') = (RR', t + Rt').$$

It can be seen as a subgroup of $GL(4)$ and represented by homogeneous coordinates:

$$(R, t) = \begin{pmatrix} R & t \\ 0 & 1 \end{pmatrix},$$

and all group operations then correspond to the matrix operations. Let the metric matrix at the identity be diagonal: $G = \text{diag}(1, 1, 1, \beta, 1, 1)$ for some $\beta > 0$, the anisotropy parameter. An orthonormal basis of the Lie algebra $\mathfrak{se}(3)$ is

$$\begin{aligned} e_1 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & e_2 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & e_3 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \\ e_4 &= \frac{1}{\sqrt{\beta}} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & e_5 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & e_6 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \end{aligned}$$

Define the corresponding structure constants $C_{ij}^k = \langle [e_i, e_j], e_k \rangle$, where the Lie bracket $[\cdot, \cdot]$ is the usual matrix commutator. It is straightforward to compute

$$C_{ij}^k = \frac{1}{\sqrt{2}} \text{ if } ijk \text{ is a direct cycle of } \{1, 2, 3\}; \quad (12)$$


$$C_{15}^6 = -C_{16}^5 = -\sqrt{\beta} C_{24}^6 = \frac{1}{\sqrt{\beta}} C_{26}^4 = \sqrt{\beta} C_{34}^5 = -\frac{1}{\sqrt{\beta}} C_{35}^4 = \frac{1}{\sqrt{2}}. \quad (13)$$

and all others that cannot be deduced by skew-symmetry of the bracket are equal to 0. The connection can then easily be computed using

$$\alpha(e_i, e_j) = \nabla_{e_i} e_j = \frac{1}{2} \sum_k (C_{ij}^k - C_{jk}^i + C_{ki}^j) e_k,$$

For $\beta = 1$, $(SE(3), G)$ is a symmetric space and the metric corresponds to the direct product metric of $SO(3) \times \mathbb{R}^3$. However, for $\beta \neq 1$, the geodesics cannot be computed in closed-form and we resort to a numerical scheme to integrate (10). According to [?], the pole ladder can be used with only one step of a fourth-order scheme to compute the exponential and logarithm maps at each rung of the ladder. We use a Runge-Kutta (RK) scheme of order 4. The Riemannian logarithm is computed with a gradient descent on the initial velocity, where the gradient of the exponential is computed by automatic differentiation. All of these are available in the `InvariantMetric` class of the package `geomstats` [?].

We now compare the integration of (11) to the pole ladder [?] for $\beta = 1.5, 2$ to parallel transport a tangent vector along a geodesic. The results are displayed on Figure 1 in a log-log plot. As expected, we reach convergence speeds of order



two for the pole ladder and the RK2 scheme, while the RK4 schemes is of order four. Both integration methods are very stable, while the pole ladder is less stable for $\sim n \geq 200$.

5 Acknowledgments

This work was partially funded by the ERC grant Nr. 786854 G-Statistics from the European Research Council under the European Unions Horizon 2020 research and innovation program. It was also supported by the French government through the 3IA Cte dAzur Investments ANR-19-P3IA-0002 managed by the National Research Agency.

References

1. Arnold, V.: Sur la gomtrie differentielle des groupes de Lie de dimension infinie et ses applications l'hydrodynamique des fluides parfaits. *Annales de l'institut Fourier* **16**(1), 319–361 (1966). <https://doi.org/10.5802/aif.233>
2. Barbaresco, F., Gay-Balmaz, F.: Lie Group Cohomology and (Multi)Symplectic Integrators: New Geometric Tools for Lie Group Machine Learning Based on Souriau Geometric Statistical Mechanics. *Entropy* **22**(5), 498 (May 2020). <https://doi.org/10.3390/e22050498>, number: 5 Publisher: Multidisciplinary Digital Publishing Institute
3. Barrau, A., Bonnabel, S.: The Invariant Extended Kalman Filter as a Stable Observer. *IEEE Transactions on Automatic Control* **62**(4), 1797–1812 (Apr 2017). <https://doi.org/10.1109/TAC.2016.2594085>, conference Name: IEEE Transactions on Automatic Control
4. Brooks, D., Schwander, O., Barbaresco, F., Schneider, J.Y., Cord, M.: Riemannian batch normalization for SPD neural networks. In: Wallach, H., Larochelle, H., Beygelzimer, A., d'Alch Buc, F., Fox, E., Garnett, R. (eds.) *Advances in Neural Information Processing Systems* 32. pp. 15489–15500. Curran Associates, Inc. (2019), <http://papers.nips.cc/paper/9682-riemannian-batch-normalization-for-spd-neural-networks.pdf>
5. Cendra, H., Holm, D.D., Marsden, J.E., Ratiu, T.S.: Lagrangian Reduction, the Euler-Poincar Equations, and Semidirect Products. *American Mathematical Society Translations* **186**(1), 1–25 (1998), number: 1 Publisher: American Mathematical Society
6. Gallier, J., Quaintance, J.: *Differential Geometry and Lie Groups: A Computational Perspective*. *Geometry and Computing*, Springer International Publishing (2020). <https://doi.org/10.1007/978-3-030-46040-2>
7. Gay-Balmaz, F., Holm, D.D., Meier, D.M., Ratiu, T.S., Vialard, F.X.: Invariant higher-order variational problems II. *J Nonlinear Sci* **22**(4), 553–597 (Aug 2012). <https://doi.org/10.1007/s00332-012-9137-2>, arXiv: 1112.6380
8. Guigui, N., Pennec, X.: Numerical Accuracy of Ladder Schemes for Parallel Transport on Manifolds (Jul 2020), <https://hal.inria.fr/hal-02894783>
9. Hauberg, S., Lauze, F., Pedersen, K.S.: Unscented Kalman Filtering on Riemannian Manifolds. *J Math Imaging Vis* **46**(1), 103–120 (May 2013). <https://doi.org/10.1007/s10851-012-0372-9>

10. Iserles, A., Munthe-Kaas, H., Nørsett, S., Zanna, A.: Lie-group methods. *Acta Numerica* (2005). <https://doi.org/10.1017/S0962492900002154>, Cambridge University Press (CUP)
11. Journee, M., Absil, P.A., Sepulchre, R.: Gradient-optimization on the orthogonal group for Independent Component Analysis. In: *Independent Component Analysis and Signal Separation*. LNCS, vol. 4666, pp. 57–64. Springer (2007)
12. Kim, K.R., Dryden, I.L., Le, H., Severn, K.E.: Smoothing splines on Riemannian manifolds, with applications to 3D shape space. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* (Dec 2020)
13. Kolev, B.: Lie Groups and mechanics: an introduction. *Journal of Nonlinear Mathematical Physics* **11**(4), 480–498 (Jan 2004). <https://doi.org/10.2991/jnmp.2004.11.4.5>, arXiv: math-ph/0402052
14. Lorenzi, M., Pennec, X.: Efficient Parallel Transport of Deformations in Time Series of Images: From Schild to Pole Ladder. *J Math Imaging Vis* **50**(1), 5–17 (Sep 2014). <https://doi.org/10.1007/s10851-013-0470-3>
15. Marsden, J.E., Ratiu, T.S.: Mechanical Systems: Mechanical systems: Symmetries and Reduction. In: Meyers, R.A. (ed.) *Encyclopedia of Complexity and Systems Science*, pp. 5482–5510. Springer, New York, NY (2009). https://doi.org/10.1007/978-0-387-30440-3_26
16. Milnor, J.: Curvatures of left invariant metrics on lie groups. *Advances in Mathematics* **21**(3), 293–329 (Sep 1976). [https://doi.org/10.1016/S0001-8708\(76\)80002-3](https://doi.org/10.1016/S0001-8708(76)80002-3)
17. Miolane, N., Guigui, N., Brigant, A.L., Mathe, J., Hou, B., Thanwerdas, Y., Heyder, S., Peltre, O., Koep, N., Zaatiti, H., Hajri, H., Cabanes, Y., Gerald, T., Chauchat, P., Shewmake, C., Brooks, D., Kainz, B., Donnat, C., Holmes, S., Pennec, X.: Geomstats: A Python Package for Riemannian Geometry in Machine Learning. *Journal of Machine Learning Research* **21**(223), 1–9 (2020), <http://jmlr.org/papers/v21/19-027.html>
18. Nava-Yazdani, E., Hege, H.C., Sullivan, T.J., von Tycowicz, C.: Geodesic Analysis in Kendall's Shape Space with Epidemiological Applications. *J Math Imaging Vis* **62**(4), 549–559 (May 2020). <https://doi.org/10.1007/s10851-020-00945-w>
19. Pennec, X., Arsigny, V.: Exponential Barycenters of the Canonical Cartan Connection and Invariant Means on Lie Groups. Springer (May 2012). https://doi.org/10.1007/978-3-642-30232-9_7, pages: 123-168
20. Yair, O., Ben-Chen, M., Talmon, R.: Parallel Transport on the Cone Manifold of SPD Matrices for Domain Adaptation. In: *IEEE Transactions on Signal Processing*. vol. 67, pp. 1797–1811 (Apr 2019). <https://doi.org/10.1109/TSP.2019.2894801>