



HAL
open science

Estimation of the tail-index and extreme quantiles from a mixture of heavy-tailed distributions

Stéphane Girard, Emmanuel Gobet

► **To cite this version:**

Stéphane Girard, Emmanuel Gobet. Estimation of the tail-index and extreme quantiles from a mixture of heavy-tailed distributions. RESIM 2021 - 13th International Workshop on Rare-Event Simulation, May 2021, Paris / Virtual, France. pp.1. hal-03235031

HAL Id: hal-03235031

<https://inria.hal.science/hal-03235031v1>

Submitted on 25 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Estimation of the tail-index and extreme quantiles from a mixture of heavy-tailed distributions

Stéphane Girard*^{1,2} and Emmanuel Gobet³

¹Inria Grenoble - Rhône-Alpes – Institut National de Recherche en Informatique et en Automatique –
France

²Laboratoire Jean Kuntzmann – Institut National de Recherche en Informatique et en Automatique,
Centre National de la Recherche Scientifique, Université Grenoble Alpes, Institut polytechnique de
Grenoble - Grenoble Institute of Technology – France

³Ecole Polytechnique [Palaiseau] – Ecole Polytechnique – École Polytechnique, 91128 Palaiseau Cedex,
France

Abstract

The estimation of extreme quantiles requires adapted methods to extrapolate beyond the largest observation of the sample. Extreme-value theory provides a mathematical framework to tackle this problem together with statistical procedures based on the estimation of the so-called tail-index describing the distribution tail. We focus on heavy-tailed distributions and consider the case where the shape of the distribution tail depends on unknown auxiliary variables. As a consequence, one has to deal with observations from a mixture of heavy-tailed distributions, and it is shown that, in such a situation, usual extreme-value estimators suffer from a strong bias. We propose several methods to mitigate this bias. Their asymptotic properties are established and their finite sample performance is illustrated both on simulated and real financial data. This is joint work with Emmanuel Gobet.

Keywords: Extreme, values, heavy, tailed distribution, mixture.

*Speaker