



HAL
open science

An equilibrated a posteriori error estimator for the curl-curl problem

Théophile Chaumont-Frelet, Martin Vohralík

► **To cite this version:**

Théophile Chaumont-Frelet, Martin Vohralík. An equilibrated a posteriori error estimator for the curl-curl problem. EFEF 2021 - 18th European Finite Element Fair, Sep 2021, Paris, France. hal-03403997

HAL Id: hal-03403997

<https://inria.hal.science/hal-03403997v1>

Submitted on 26 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An equilibrated a posteriori error estimator for the curl–curl problem

T. Chaumont-Frelet^{*,†}, M. Vohralík^{§,‡}

18th European finite element fair, September 2021

* Inria Sophia-Antipolis, † Laboratoire J.A. Dieudonné, § Inria Paris, ‡ CERMICS

Outline

- 1 Equilibration for the Poisson problem
- 2 Equilibration for the curl–curl problem

The Poisson problem

Poisson problem

Consider a Lipschitz polyhedral domain Ω and $f \in L^2(\Omega)$.

Our first model problem is to find $u \in H_0^1(\Omega)$ such that

$$(\nabla u, \nabla v)_\Omega = (f, v)_\Omega$$

for all $v \in H_0^1(\Omega)$.

u_h is the Lagrange FEM approximation of u with degree $p + 1$.

For the sake of simplicity, I will assume that $f = f_h \in \mathcal{P}_p(\mathcal{T}_h)$.

The idea of flux equilibration

Assume that we have a field $\sigma_h \in \mathbf{H}(\text{div}, \Omega)$ such that

$$\nabla \cdot \sigma_h = f_h \text{ in } \Omega$$

at our disposal.

Then, we have

$$\begin{aligned}(\nabla(u - u_h), \nabla v)_\Omega &= (f_h, v)_\Omega - (\nabla u_h, \nabla v) \\ &= (\nabla \cdot \sigma_h, v)_\Omega - (\nabla u_h, \nabla v) \\ &= -(\sigma_h + \nabla u_h, \nabla v),\end{aligned}$$

for all $v \in H_0^1(\Omega)$ and in particular

$$\|\nabla(u - u_h)\|_\Omega \leq \|\sigma_h + \nabla u_h\|_\Omega.$$

Prager-Synge theorem

Equilibrated flux

$$\boldsymbol{\sigma}_h \in \mathbf{H}(\operatorname{div}, \Omega); \quad \nabla \cdot \boldsymbol{\sigma}_h = \mathbf{f}_h \text{ in } \Omega$$

Error estimate

$$\|\nabla(u - u_h)\|_{\Omega} \leq \|\boldsymbol{\sigma}_h + \nabla u_h\|_{\Omega}.$$



W. Prager and J.L. Synge, 1947

The particular choice $\boldsymbol{\sigma} := -\nabla u$ saturates the bound.


Local flux constructions

The idea flux would be $\sigma := -\nabla u$.

Let's characterize it locally and cook up a discrete computable version.

Let us set

$$\sigma^a := \psi_a \sigma$$



where ψ_a is the “hat function” associated with the vertex a , so that

$$\sigma = \sum_{a \in \mathcal{V}_h} \sigma^a.$$

Characterization of the local contributions

We know that $\boldsymbol{\sigma}^a := -\psi_a \nabla u$. So that in particular

$$\boldsymbol{\sigma}^a \in \mathbf{H}_0(\text{div}, \omega_a)$$

where ω_a is the set of tetrahedra $K \in \mathcal{T}_h$ sharing the vertex \mathbf{a} , and

$$\nabla \cdot \boldsymbol{\sigma}^a = \psi_a f_h - \nabla \psi_a \cdot \nabla u.$$

As a result, we have the characterization

$$\boldsymbol{\sigma}^a = \arg \min_{\substack{\mathbf{v} \in \mathbf{H}_0(\text{div}, \omega_a) \\ \nabla \cdot \mathbf{v} = \psi_a f_h - \nabla \psi_a \cdot \nabla u}} \|\mathbf{v} + \psi_a \nabla u\|_{\omega_a},$$

since the minimum is zero and achieved when $\mathbf{v} = -\psi_a \nabla u$.

Discrete construction

Recall that

$$\sigma^a = \arg \min_{\substack{\mathbf{v} \in \mathbf{H}_0(\operatorname{div}, \omega_a) \\ \nabla \cdot \mathbf{v} = \psi_a f_h - \nabla \psi_a \cdot \nabla u}} \|\mathbf{v} + \psi_a \nabla u\|_{\omega_a}.$$

As a discrete counterpart, we set

$$\sigma_h^a := \arg \min_{\substack{\mathbf{v}_h \in \mathbf{RT}_{p+1}(\mathcal{T}_h^a) \cap \mathbf{H}_0(\operatorname{div}, \omega_a) \\ \nabla \cdot \mathbf{v}_h = \psi_a f_h - \nabla \psi_a \cdot \nabla u_h}} \|\mathbf{v}_h + \psi_a \nabla u_h\|_{\omega_a}.$$

This is indeed well-defined since

$$(\psi_a f - \nabla \psi_a \cdot \nabla u_h, 1)_{\omega_a} = (f, \psi_a)_{\Omega} - (\nabla u_h, \nabla \psi_a)_{\omega_a} = 0$$

whenever $a \notin \partial\Omega$.

Summation provides an equilibrated flux:

$$\boldsymbol{\sigma}_h := \sum_{\mathbf{a} \in \mathcal{V}_h} \boldsymbol{\sigma}_h^{\mathbf{a}},$$

and reliability follows from the Prager–Synge theorem.

Efficiency

$$\|\boldsymbol{\sigma}_h + \nabla u_h\|_{\mathcal{K}} \lesssim \|\nabla(u - u_h)\|_{\tilde{\mathcal{K}}}$$

The hidden constant does not depend on p .

The curl–curl problem

The curl–curl problem

For the sake of simplicity, assume that Ω is simply connected, and consider a divergence-free right-hand side $\mathbf{J}_h \in \mathbf{RT}_p(\mathcal{T}_h)$.

Our model problem is then to find $\mathbf{A} \in \mathbf{H}_0(\mathbf{curl}, \Omega)$ such that

$$(\nabla \times \mathbf{A}, \nabla \times \mathbf{v})_\Omega = (\mathbf{J}_h, \mathbf{v})_\Omega, \quad (\mathbf{A}, \nabla q)_\Omega = 0,$$

for all $\mathbf{v} \in \mathbf{H}_0(\mathbf{curl}, \Omega)$ and $q \in H_0^1(\Omega)$.

$\mathbf{A}_h \in \mathbf{N}_p(\mathcal{T}_h) \cap \mathbf{H}_0(\mathbf{curl}, \Omega)$ is the Nédélec approximation of \mathbf{A} .

Prager-Synge theorem

Equilibrated flux

$$\mathbf{B}_h \in \mathbf{H}(\text{curl}, \Omega); \quad \nabla \times \mathbf{B}_h = \mathbf{J}_h$$

Error estimate

$$\|\nabla \times (\mathbf{A} - \mathbf{A}_h)\|_{\Omega} \leq \|\mathbf{B}_h - \nabla \times \mathbf{A}_h\|_{\Omega}$$

The “ideal” flux $\mathbf{B} := \nabla \times \mathbf{A}$ saturates the bound.

The issue with localization (1/2)

We follow the same steps than for the Poisson problem.

The ideal flux is $\mathbf{B} := \nabla \times \mathbf{A}$. If we set $\mathbf{B}^a := \psi_a \nabla \times \mathbf{A}$, then

$$\nabla \times \mathbf{B}^a = \psi_a \mathbf{J}_h + \nabla \psi_a \times \nabla \times \mathbf{A},$$

so that

$$\mathbf{B}^a = \arg \min_{\substack{\mathbf{v} \in H_0(\text{curl}, \omega_a) \\ \nabla \times \mathbf{v} = \psi_a \mathbf{J} + \nabla \psi_a \times \nabla \times \mathbf{A}}} \|\mathbf{v} - \psi_a \nabla \times \mathbf{A}\|_{\omega_a}.$$

The issue with localization (2/2)

Recall that

$$B^a = \arg \min_{\substack{\mathbf{v} \in H_0(\text{curl}, \omega_a) \\ \nabla \times \mathbf{v} = \psi_a \mathbf{J}_h + \nabla \psi_a \times \nabla \times \mathbf{A}}} \|\mathbf{v} - \psi_a \nabla \times \mathbf{A}\|_{\omega_a}.$$

Unfortunately, we can not set

$$B_h^a := \arg \min_{\substack{\mathbf{v}_h \in N_{p+1}(\mathcal{T}_h^a) \cap H_0(\text{curl}, \omega_a) \\ \nabla \times \mathbf{v}_h = \psi_a \mathbf{J}_h + \nabla \psi_a \times \nabla \times \mathbf{A}_h}} \|\mathbf{v}_h - \psi_a \nabla \times \mathbf{A}_h\|_{\omega_a}$$

as the minimization set is empty: the field

$$\psi_a \mathbf{J}_h + \nabla \psi_a \times \nabla \times \mathbf{A}_h$$

is not divergence-free!

Possible solutions

An initial idea for lowest-order elements:



D. Braess and J. Schöberl, 2008

Recently developed extensions:



J. Gedicke, S. Geeveres and I. Perugia, 2019



J. Gedicke, S. Geeveres, I. Perugia and J. Schöberl, 2020



T. Chaumont-Frelet and M. Vohralík, 2021

Here, I will detail the last construction.

The idea

Let $\boldsymbol{\theta}^a := \nabla\psi_a \times \nabla \times \mathbf{A}$. There are two important properties:

$$\sum_{a \in \mathcal{V}_h} \boldsymbol{\theta}^a = \mathbf{0}, \quad \nabla \cdot \boldsymbol{\theta}^a = -\nabla\psi_a \cdot \mathbf{J}_h.$$

At the discrete level, we have

$$\sum_{a \in \mathcal{V}_h} (\nabla\psi_a \times \nabla \times \mathbf{A}_h) = \mathbf{0}, \quad \nabla \cdot (\nabla\psi_a \times \nabla \times \mathbf{A}_h) \neq -\nabla\psi_a \cdot \mathbf{J}_h.$$

It is tempting to set

$$\boldsymbol{\theta}_h^a := \arg \min_{\substack{\mathbf{v} \in \mathbf{RT}_p(\mathcal{T}_h^a) \cap \mathbf{H}_0(\text{div}, \omega_a) \\ \nabla \cdot \mathbf{v} = -\nabla\psi_a \cdot \mathbf{J}_h}} \|\mathbf{v} - \nabla\psi_a \times \nabla \times \mathbf{A}_h\|_{\omega_a},$$

but it does not sum up to zero.

Trick #1: Over-constrained minimization

We instead set

$$\hat{\boldsymbol{\theta}}_h^a := \arg \min_{\substack{\mathbf{v} \in \mathbf{RT}_p(\mathcal{T}_h^a) \cap \mathbf{H}_0(\operatorname{div}, \omega_a) \\ (\mathbf{v} - \nabla \psi_a \times \nabla \times \mathbf{A}_h, \mathbf{r})_{\omega_a} = 0 \quad \forall \mathbf{r} \in \mathcal{P}_0(\mathcal{T}_h^a) \\ \nabla \cdot \mathbf{v} = -\nabla \psi_a \cdot \mathbf{J}_h}} \|\mathbf{v} - \nabla \psi_a \times \nabla \times \mathbf{A}_h\|_{\omega_a},$$

where the mean-value constrain works because \mathbf{A}_h solves the discrete problem. We then set

$$\hat{\boldsymbol{\theta}}_h := \sum_{\mathbf{a} \in \mathcal{V}_h} \hat{\boldsymbol{\theta}}_h^a.$$

Importantly, we have

$$(\hat{\boldsymbol{\theta}}_h, \mathbf{r})_{\Omega} = 0 \quad \forall \mathbf{r} \in \mathcal{P}_0(\mathcal{T}_h).$$

Trick #2: Divergence-free decomposition of $\widehat{\boldsymbol{\theta}}$ (1/2)

The $\widehat{\boldsymbol{\theta}}_h^a$ have the correct divergence, but does not sum up to zero.
We need a divergence-free decomposition of $\widehat{\boldsymbol{\theta}}_h$ into contributions $\widetilde{\boldsymbol{\theta}}_h^a$.

We set for each $\mathbf{a} \in \mathcal{V}_h$ and $K \in \mathcal{T}_h^a$

$$\widetilde{\boldsymbol{\theta}}_h^a|_K := \arg \min_{\substack{\mathbf{v}_h \in \mathbf{RT}_{p+1}(K) \\ \nabla \cdot \mathbf{v}_h = 0 \\ \mathbf{v}_h \cdot \mathbf{n}_K = \psi_a \widehat{\boldsymbol{\theta}}_h \cdot \mathbf{n}_K \text{ on } \partial K}} \|\mathbf{v}_h - \psi_a \widehat{\boldsymbol{\theta}}_h\|_K$$

This is indeed well-posed since

$$\begin{aligned} (\psi_a \widehat{\boldsymbol{\theta}}_h \cdot \mathbf{n}_K, 1)_{\partial K} &= (\psi_a, \widehat{\boldsymbol{\theta}}_h \cdot \mathbf{n}_K)_{\partial K} \\ &= (\nabla \psi_a, \widehat{\boldsymbol{\theta}}_h)_K + (\nabla \cdot \widehat{\boldsymbol{\theta}}_h, \psi_a)_K \\ &= 0. \end{aligned}$$

Trick #2: Divergence-free decomposition of $\widehat{\theta}$ (2/2)

We can then actually show that

$$\sum_{\mathbf{a} \in \mathcal{V}_h} \widetilde{\theta}_h^{\mathbf{a}} = \widehat{\theta}_h,$$

so that setting

$$\theta_h^{\mathbf{a}} := \widehat{\theta}_h^{\mathbf{a}} - \widetilde{\theta}_h^{\mathbf{a}} \in \mathbf{H}_0(\text{div}, \omega_{\mathbf{a}}),$$

we have

$$\nabla \cdot \theta_h^{\mathbf{a}} = -\nabla \psi_{\mathbf{a}} \cdot \mathbf{J}_h, \quad \sum_{\mathbf{a} \in \mathcal{V}_h} \theta_h^{\mathbf{a}} = \mathbf{0}.$$

Local flux construction and efficiency

Since the θ_h^a have all the required properties, we can now set

$$B_h^a := \arg \min_{\substack{\mathbf{v}_h \in \mathbf{N}_{p+1}(\mathcal{T}_h^a) \cap \mathbf{H}_0(\text{curl}, \omega_a) \\ \nabla \times \mathbf{v}_h = \psi_a \mathbf{J}_h + \theta_h^a}} \|\mathbf{v}_h - \psi_a \nabla \times \mathbf{A}_h\|_{\omega_a},$$

and we construct an equilibrated flux as

$$B_h := \sum_{a \in \mathcal{V}_h} B_h^a.$$

Efficiency

$$\|B_h - \nabla \times A_h\|_{\mathcal{K}} \lesssim \|\nabla \times (A - A_h)\|_{\tilde{\mathcal{K}}}$$

The hidden constant does not depend on p .



T. Chaumont-Frelet, A. Ern and M. Vohralík, 2021