



Eukaryotic stress induced mutagenesis is limited by a local control of Translesion Synthesis

Katarzyna Maslowska, Florencia Villafanez, Luisa Laureti, Shigenori Iwai,
Vincent Pagès

► To cite this version:

Katarzyna Maslowska, Florencia Villafanez, Luisa Laureti, Shigenori Iwai, Vincent Pagès. Eukaryotic stress induced mutagenesis is limited by a local control of Translesion Synthesis. *Nucleic Acids Research*, 2022, 50 (4), pp.2074-2080. 10.1093/nar/gkac044 . hal-03413120

HAL Id: hal-03413120

<https://hal.science/hal-03413120>

Submitted on 8 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Eukaryotic stress-induced mutagenesis is limited by a local control of translesion synthesis

Katarzyna H. Masłowska¹, Florencia Villafañez¹, Luisa Laureti¹, Shigenori Iwai² and Vincent Pagès^{1,*}

¹Cancer Research Center of Marseille: Team DNA Damage and Genome Instability | CNRS, Aix Marseille Univ, Inserm, Institut Paoli-Calmettes, Marseille 13009, France and ²Graduate School of Engineering Science, Osaka University, Osaka 560-8531, Japan

Received August 26, 2021; Revised January 11, 2022; Editorial Decision January 13, 2022; Accepted January 14, 2022

ABSTRACT

The DNA damage response (DDR) preserves the genetic integrity of the cell by sensing and repairing damages after a genotoxic stress. Translesion Synthesis (TLS), an error-prone DNA damage tolerance pathway, is controlled by PCNA ubiquitination. In this work, we raise the question whether TLS is controlled locally or globally. Using a recently developed method that allows to follow the bypass of a single lesion inserted into the yeast genome, we show that (i) TLS is controlled locally at each individual lesion by PCNA ubiquitination, (ii) a single lesion is enough to induce PCNA ubiquitination and (iii) PCNA ubiquitination is imperative for TLS to occur. More importantly, we show that the activation of the DDR that follows a genotoxic stress does not increase TLS at individual lesions. We conclude that unlike the SOS response in bacteria, the eukaryotic DDR does not promote TLS and mutagenesis.

INTRODUCTION

Accurate DNA replication is essential for genome stability. Since DNA is constantly insulted by endogenous and exogenous DNA-damaging agents, organisms have evolved several mechanisms to deal with DNA damage. The DNA damage response (DDR) includes cell cycle arrest, lesion repair and lesion tolerance (1). Numerous repair systems remove various modifications from DNA in an error-free manner (2). However, despite their efficient action, it is inevitable that some lesions might be present during replication. Most DNA damage impedes DNA synthesis by high-fidelity replicative DNA polymerases. Therefore, to complete replication and maintain cell survival in the presence of residual DNA damage, cells have evolved two lesion tolerance mechanisms: Damage Avoidance (DA) and Translesion Synthesis (TLS). Damage avoidance (also named strand switch, copy choice or homology directed gap

repair) is a pathway relying on the information of the newly replicated sister chromatid to circumvent the lesion in an error-free manner ((3,4), reviewed in (5)). Translesion synthesis is a potentially mutagenic pathway that employs specialized DNA polymerases able to insert a nucleotide directly opposite the lesion (reviewed in (6) and (7)).

How the DDR controls mutagenesis has been widely studied in prokaryotic cells: the SOS response, by increasing the expression level of TLS polymerases in response to a genotoxic stress, greatly contributes to mutagenesis and therefore, to the adaptive response to environmental stress (8). A good example of this phenomenon is the importance of the SOS response in resistance to antibiotics (9). Moreover, experiments involving the study of a single lesion have shown that the pre-induction of the SOS system by a genotoxic stress greatly increases the level of TLS and mutagenesis at the studied lesion (10,11). Hence, in bacteria, two factors contribute to the increase in mutagenesis in response to a genotoxic stress: (i) the number of lesions (the higher the number of lesions, the higher the probability to generate a mutation) and (ii) the increased level of TLS polymerases in response to the induction of SOS (the more TLS polymerases, the higher probability to bypass the lesion by TLS). Thus, the SOS response is a global response that favors TLS and mutagenesis.

The eukaryotic DDR includes mostly posttranslational modifications such as phosphorylation and ubiquitination (12). Damage-induced transcriptional regulation is less common but has also been reported (13). Proliferating Cell Nuclear Antigen (PCNA) ubiquitination regulates lesion tolerance in response to DNA damage. After the replication fork encounters a DNA lesion, PCNA stalls at the lesion site, and single-stranded DNA (ssDNA) is generated downstream of the lesion. The formation of RPA protein-coated ssDNA leads to the recruitment of the Rad6-Rad18 complex and the subsequent mono-ubiquitination of PCNA at lysine K164 (14–16). This mono-ubiquitination can be further extended by Rad5 and Ubc13-Mms2, through the formation of K63-linked ubiquitin chains (17,18). It is well established that in eukaryotic cells, PCNA

*To whom correspondence should be addressed. Tel: +33 4 86 977 384; Email: vincent.pages@cnrs.fr



A Outline of the integration system**B PCNA ubiquitination is imperative for TLS**

Figure 1. PCNA ubiquitination is imperative for TLS. (A) Outline of the integration system: A non-replicative plasmid containing a single lesion is integrated into a yeast chromosome using a Cre/lox site-specific recombination. The integrative vector carrying a selection marker (LEU2) and the 5'-end of the *lacZ* reporter gene containing a single lesion is introduced into a specific locus of the chromosome with the 3'-end of *lacZ*. The precise integration of the plasmid DNA into the chromosome restores a functional *lacZ* gene, enabling the phenotypical detection of TLS and DA events (as blue and white colonies on X-gal indicator media). (B) Bypass of the N2dG-AAF lesion in strains deficient in TLS polymerases (Rev1, Pol η /rad30, Pol ζ /rev3) and strains deficient in PCNA ubiquitination (*rad18* and *pol30K164R* mutant). T-tests were performed to compare the TLS values from the different mutants to the parental strain: * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$.

mono-ubiquitination stimulates TLS, while PCNA polyubiquitination triggers DA (reviewed in (19,20)).

Forward mutagenesis assays have shown that the mutation frequency rapidly increases with the amount of genotoxic stress inflicted to the cell. While it is expected that the level of mutagenesis increases with the number of lesions (more lesions lead to more mutations), it is not known if an additional regulatory mechanism also contributes to the increase in mutagenesis in eukaryotic cells. Traditional bulk approaches do not allow to determine if the mutagenesis level is solely correlated to the number of lesions, or if a more global DNA damage response also favors mutagenesis.

As PCNA ubiquitination controls TLS, and PCNA ubiquitination increases in response to genotoxic stress, it appears intuitive that such global response exists: the more PCNA is mono-ubiquitinated, the more TLS will be used by the cell. However, the existence of such global response has never been demonstrated.

In this study, we set out to determine whether the level of TLS is regulated solely at the local level, or if the amount of

damage present in the cell could favor TLS in a more global manner. We have recently devised a method to introduce a single lesion in the genome of a yeast cell (Figure 1A) (21). Such approach allows to dissect the regulation of the tolerance mechanisms in different genetic backgrounds as well as in different conditions of external stress for the cell. We have used our assay to determine whether the ratio between TLS and DA at the level of a single lesion is modulated by increasing level of genotoxic stress resulting in global DNA damage response and increased PCNA ubiquitination.

MATERIALS AND METHODS

Strains and media

All strains used in the present study are derivative of strain EMY74.7 (22) (MATa his3- Δ 1 leu2-3,112 trp1- Δ ura3- Δ met25- Δ phr1- Δ rad14- Δ msh2 Δ ::hisG). Gene disruptions were achieved using PCR-mediated seamless gene deletion (23) or URABlast (24) techniques. All strains used in the study are listed in Table 1.

Table 1. Strains used in the study

Strain	Relevant Genotype
EVP161	(All yeast strains are: <i>MATa his3-ΔI leu2-3,112 trp1-Δ ura3-Δ met25-Δ rad14-Δ phr1-Δ msh2Δ::hisG</i>)
SC22	<i>E. coli</i> MG1655 <i>phrB::kan</i>
SC53	<i>rad14 phr1</i>
SC55	VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC82	VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC83	<i>rev1-Δ</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC86	<i>rev1-Δ</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC87	<i>rad30Δ::hisG</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC181	<i>rad30Δ::hisG</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC182	<i>rev3Δ::hisG</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC203	<i>rev3Δ::hisG</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC206	<i>rad18Δ::hisG</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC236	<i>rad18Δ::hisG</i> III(75494–75499)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC237	<i>pol30-K164R</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC533	<i>pol30-K164R</i> VI(167260–167265)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC534	<i>pol30::3FLAG-kan</i> VI (167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
SC535	<i>pol30::3FLAG-kan</i> VI (167260–167265)::(<i>lox66-3' lacZ-MET25/lead</i>)
SC537	<i>ubc13Δ Pol30::3FLAG-kan</i> VI (167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)
	<i>pol30-K164R::3FLAG-kan</i> VI (167260–167265)::(<i>lox66-3' lacZ-MET25/lag</i>)

Integration system

Integration of plasmids carrying cis-syn TT dimer / 6–4 (TT) / N2dG-AAF lesions (or control plasmids without lesion) and result analysis was performed as previously described (21). In experiments where cells were treated with 4-NQO, the overnight culture was inoculated into 100 ml of yeast extract/peptone/dextrose medium (YPD) per integrated lesion to reach OD₆₀₀ = 0.3 and incubated at 30°C with shaking until OD₆₀₀ = 0.8. After the addition of 150 ng/ml of 4-NQO, cultures were incubated for 30 more minutes. 4-NQO was inactivated by adding an equal volume of 10% sodium thiosulfate, and then cells were further washed and processed the same way as untreated cultures. For UV treatment, the overnight culture was inoculated into 50 ml of YPD per integrated lesion to reach OD₆₀₀ = 0.3 and incubated at 30°C with shaking until OD₆₀₀ = 1.6. Cells were then harvested, resuspended in twice the initial volume of water, and treated with UV (4 J.m⁻²) in Petri dishes (15 cm Ø, 25 ml/dish). Cells were further washed and processed the same way as untreated cultures.

All experiments were performed in triplicate or more. Only the N2dG-AAF lesion in the UV-induced condition was done in duplicate. Graphs and statistical analysis were performed using GraphPad Prism applying unpaired *t*-test. Bars represent the mean value ± s.d.

Preparation of an oligonucleotide containing the (6-4) photoproduct

A 13-mer oligonucleotide, d(GCAAGTTAACACG), purchased from Tsukuba Oligo Service (150 nmol) was dissolved in water (7.5 ml), and after a nitrogen purge for 5 min, the solution was heated at 75°C for 5 min and cooled in ice. This solution was poured into a petri dish with a diameter of 8.5 cm and irradiated for 20 min on a Spectrolinker XL-1500 UV Crosslinker (Spectronics Corp.) equipped with six 15 W germicidal lamps (254 nm, 2.2 mW.cm⁻²). The reaction mixture was analyzed by reversed-phase HPLC using a Waters μBondasphere C18 5μm 300Å column (3.9 × 150 mm) at 50°C with an acetonitrile gra-

dient of 6 to 10% during 20 min in 0.1 M triethylammonium acetate (pH 7.0). The elution was monitored by using a Waters 2998 photodiode array detector, and the product with absorption at 325 nm (which is characteristic of the (6-4) photoproduct) was isolated by repeating injection of 500 μl. After concentration of the combined eluates, impurities detected by the second HPLC analysis using a GL Sciences Inertsil ODS-3 5μm column (4.6 × 250 mm) with an acetonitrile gradient of 8 to 11% were removed. The final eluate (chromatogram shown in Supplementary Figure S2) was concentrated and co-evaporated with water. The yield determined from the absorbance at 260 nm was 4.5 nmol. This process is similar to the one previously described by LeClerc *et al.* (25)

Immunoprecipitation and western blot

Cells expressing FLAG-tagged PCNA were grown and processed as for the integration of the plasmid with a lesion. After electroporation cells were left to recover during 30 min, harvested, washed with buffer containing glycerol (1× PBS, 10% glycerol and 1 mM EDTA) and frozen at –80°C. For protein extraction cells were resuspended in protein extraction buffer (1× PBS, 10% glycerol, 1 mM EDTA, 1 mM PMSF, 1× Roche cOmplete™ Protease Inhibitor Cocktail) and lysed in a bead beater. Lysate was clarified by centrifugation, and total protein concentration was determined using Qubit Fluorometer according to manufacturer’s protocol (ThermoFisher). Crude protein extract was incubated with Anti-FLAG® M2 Magnetic Beads (Merck) overnight, in an IP buffer (1× PBS, 5% glycerol, 0.5 mM EDTA, 1 mM PMSF, 1× Roche cOmplete™ Protease Inhibitor Cocktail, 2 mM DTT). Proteins were resolved by SDS-PAGE (Bio-Rad 12% Mini-PROTEAN® TGX™ Precast Protein Gels) and transferred onto PVDF membranes using Bio-Rad Trans-Blot Turbo Transfer System for Western blotting. Modified and non-modified PCNA was detected using ANTI-FLAG® M2-Peroxidase (HRP) antibody (Merck).

For Rad53 detection, protein extracts for western blot analysis were prepared by trichloroacetic acid (TCA) precipitation, as described previously (26). Proteins were

resolved by SDS-PAGE (Bio-Rad 10% Mini-PROTEAN® TGX™ Precast Protein Gels) and transferred onto PVDF membranes. Rad53, phosphorylated and not, was detected using anti-Rad53 polyclonal antibodies (Abcam ab104232).

Rifampicin mutagenesis assay

Escherichia coli strain EVP161 (MG1655 *phrB::kan*) was grown to exponential phase in LB media, centrifuged and resuspended in 10 mM MgSO₄ before UV irradiation with different UV doses (0–150 J.m⁻²). An aliquot of cells was plated on LB to assess survival. Cells were then diluted 1/20 in LB and grown for 6h before plating on LB and LB + Rif 100 µg/ml. Colonies were counted after 24 h on LB and 48 h on LB + Rif.

Canavanine mutagenesis assay

Mutagenesis experiments were performed as described in Unk and Daraba (33). Briefly, yeast cultures were grown to saturation in YPD, harvested, resuspended in water (10⁸ cells/ml) and sonicated to separate clumps of cells. Serial dilutions were plated on SD-agar plates containing canavanine for mutagenesis, and on SC-agar plates for survival. Plates were left to absorb the moisture, and irradiated with different doses of UV (0–100 J.m⁻²). Colonies were counted after 4 days of incubation.

RESULTS

PCNA ubiquitination occurs locally at a single lesion

The assay we have previously described (21) is based on a non-replicative plasmid containing a single lesion which is stably integrated into a specific site of the yeast genome. The precise integration of the plasmid DNA into the chromosome restores a functional *lacZ* gene, enabling the phenotypical detection of TLS and DA events (as blue and white colonies on X-gal indicator media). In order to study tolerance events, we inactivated the following repair mechanisms: nucleotide excision repair (*rad14*), photolyase (*phr1*) and mismatch repair system (*msh2*). Tolerance events are calculated as the ratio of colonies resulting from the integration of the damaged vector versus the non-damaged vector.

Using this method, we have introduced a N2dG-AAF (N2-dG-Acetylaminofluorene) adduct in the genome of cells deficient for repair mechanisms but proficient for lesion tolerance. 2-Acetylaminofluorene is a potent carcinogen known to produce liver tumors in rats (27). We observe for this lesion a TLS level of 18% that relies almost exclusively on the TLS polymerases Rev1 and Pol ζ (Figure 1B). We also observed a reduction of TLS in the *rad30* mutant showing that Pol η is also involved in the bypass of this lesion as previously shown (28). Indeed, it has been previously suggested that Pol η can participate to the insertion step at the N2dG-AAF lesion, while Pol ζ is required for the extension step (28).

We can note that the level of TLS is highly dependent on the type of lesion as we previously measured 55% of TLS for the cis-syn TT dimer lesion (cyclobutane pyrimidine dimer) and 5% of TLS for the (6-4)TT photoproduct

lesion (thymine–thymine pyrimidine(6-4)pyrimidone photoproduct) (21). Replication through all these lesions relies exclusively on TLS polymerases as the inactivation of *rad30*, *rev3* and/or *rev1* abolishes TLS (21).

We then introduced the N2dG-AAF lesion in cells that were unable to ubiquitinate PCNA, either in *rad18* strains or strains where the lysine 164 of PCNA was mutated into an arginine (pol30K164R) (Figure 1B). In both conditions, TLS was drastically reduced which is consistent with the current view of PCNA ubiquitination promoting TLS. The same results were previously observed for the cis-syn TT dimer and (6-4)TT photoproduct lesions (21). Therefore, PCNA ubiquitination appears as an imperative for TLS to occur since the level of TLS is close to 0% in its absence. Given that the lesions were introduced in the cells in the absence of any other DNA damaging treatment, the presence of the single lesion is enough to generate the signal required to trigger Rad6–Rad18-mediated PCNA ubiquitination. Hence, these data indicate that PCNA ubiquitination occurs locally, and that the single patch of ssDNA generated at one lesion is enough to recruit the Rad6–Rad18 complex that will ubiquitinate PCNA allowing TLS to occur at the lesion. We can note here that the DA level remains high despite the absence of PCNA ubiquitination. These events may be attributed to the ‘salvage recombination pathway’ that bypasses lesions independently of PCNA ubiquitination and may utilize Rad52 (5,29,30). The roles and regulation of the ‘salvage’ homologous recombination pathway in lesion bypass will be further explored in future studies.

We then wondered if in addition to this local regulation, a more global control of TLS exists. In other words, can the activation of the DDR and/or the increased level of ubiquitylated PCNA generated by a genotoxic stress increase the use of TLS at each lesion?

Is TLS modulated by a global stress response?

To answer this question, we treated the cells with DNA damaging agents (either 4-NQO: 4-Nitroquinoline-1-oxide (31), or UV irradiation) prior to the integration of three types of lesions (cis-syn TT dimer, (6-4)TT photoproduct or N2dG-AAF). We used a dose that causes about 80% lethality. Such treatment is known to induce a strong DDR, which includes PCNA ubiquitination and Rad53 phosphorylation. Indeed, we showed by western blot that 4-NQO treatment leads to PCNA ubiquitination (Figure 2B). Similarly, we showed that UV-irradiation leads to PCNA ubiquitination (Supplementary Figure S1). We also verified that the treatment to make the cells competent for electroporation and the electroporation itself were not inducing stress triggering PCNA ubiquitination. This confirms that the TLS data obtained previously (Figure 1) is independent of any genotoxic stress and only related to the single inserted lesion. We also showed that 4NQO treatment leads to complete Rad53 phosphorylation (Figure 2C), indicating the activation of the DDR.

Having confirmed that PCNA is ubiquitinated and the DDR is activated in response to 4-NQO treatment, we introduced a single lesion cis-syn TT dimer, (6-4)TT photoproduct or N2dG-AAF after treating the cells with 4-NQO for 30 min, or after UV-irradiation. No increase in TLS was




Figure 2. TLS is not modulated by a global stress response. (A) Partition of DNA Damage Tolerance events for different lesions (cis-syn TT dimer, (6-4)TT photoproduct and N2dG-AAF) non-treated (left) and treated with 4-NQO or UV irradiation (right) prior to the integration. No significant difference was observed between the non-treated and treated conditions. (B) Western blot analysis revealing FLAG-tagged PCNA shows that the electroporation conditions do not induce significant level of PCNA ubiquitination. Upon treatment, two bands appear that correspond to mono- and bi-ubiquitination of PCNA. In the pol30-K164R mutant, these two bands are no longer present since the site of ubiquitination of PCNA (lysine 164) is absent. In the ubc13 mutant, the polyubiquitination band is absent. (C) Western blot analysis revealing Rad53 phosphorylation in response to 4-NQO treatment.

observed in cells pre-treated by 4-NQO as compared to the nontreated cells. The same results were obtained with UV-irradiated cells (Figure 2A).

It appears therefore that the activation of a global DDR (as monitored by Rad53 phosphorylation) does not modulate the bypass of individual lesion via TLS. This suggests that TLS is controlled at a local level, rather than globally.

Global versus local response to DNA damage

To confirm this hypothesis, and to avoid any bias that could be introduced by our integration system, we compared UV-induced mutagenesis in bacterial and yeast strains (Figure 3). In short, cells were UV-irradiated at different UV doses and mutants were counted as Rifampicin resistant colonies for *E. coli* (32), or as Canavanine resistant colonies for *Saccharomyces cerevisiae* (33).

In *E. coli*, we observe a quadratic correlation between the UV dose and mutagenesis. Indeed, the number of mutations increases with both the number of lesions, and with the probability of a lesion to turn into a mutation (that also increases with the amount of lesions that induce the SOS response). Consequently, the number of mutations increases with the square of the number of lesions (quadratic correlation). This reflects the global effect of the SOS response.

On the other hand, in yeast we observe a linear correlation that even tends to reach a plateau for high doses. This shows that the genotoxic stress does not favor TLS, and mu-

tagenesis only increases with the number of lesions. This reflects the absence of global regulation of TLS in this organism.

DISCUSSION

Our data show that a single lesion is enough to induce PCNA ubiquitination locally, and that PCNA ubiquitination is imperative for TLS to occur. More importantly, we show that a genotoxic stress leading to an increased level of PCNA ubiquitination and to the activation of the DNA damage response (leading to complete Rad53 phosphorylation) has no effect on the level of TLS at individual lesions. We conclude from these data that the eukaryotic DNA damage response does not favor the increase of TLS and mutagenesis. This observation was confirmed by UV-induced mutagenesis experiments showing that the mutagenic response increases only linearly with the level of damage in yeast. This response is quite different from the one in *E. coli* where it has previously been shown that the level of TLS at individual lesion increases upon UV-irradiation (10), and where the mutagenic response shows a quadratic correlation with the UV dose.

While the bacterial SOS response implies transcriptional activation, the eukaryotic DDR includes mostly post-translational modifications such as phosphorylation and ubiquitination (12). PCNA is ubiquitinated in response to DNA damage, but the level of ubiquitination generally observed (our study and (16)) is rather low, with free-PCNA




Figure 3. UV-induced mutagenesis in *E. coli* and *S. cerevisiae*. *E. coli* mutants were scored as rifampicin resistant colonies, *S. cerevisiae* as canavanine resistant colonies. Each point represents the average and standard deviation of at least three independent experiments. The curve in red represents the fit following a quadratic equation for the Rif mutagenesis ($R^2 = 0.93$) or a linear fit for the Can mutagenesis ($R^2 = 0.82$).

remaining predominant. Even in the absence of deubiquitylases, free-PCNA remains highly majoritarian (34). Given the low level of PCNA ubiquitination that is generally observed for *S. cerevisiae*, we can hypothesize that *de novo* ubiquitination occurs at each encounter with a lesion. This model is compatible with previous *in vitro* data that showed that PCNA remains at the lesion terminus where RPA coated ssDNA prevents its diffusion (35). In this model, a new PCNA is loaded downstream of the lesion allowing the replication to go on, while the PCNA that is maintained at the lesion site can be ubiquitinated and allows TLS to perform the gap filling reaction. Once the gap filling reaction is achieved, ubiquitinated PCNA is unloaded and will not contribute to TLS at another site.

Taken together, our data show that the regulation of TLS occurs locally, and that in eukaryotic cells, there is no global response capable of increasing the level of TLS and mutagenesis in response to a genotoxic stress. The level of mutagenesis depends solely on the number of lesions present in the genome (following a linear correlation). Similar mutagenesis assays in human cells have shown the same linear response (36,37). Unlike bacteria that show a quadratic increase in mutagenesis, allowing them to rapidly mutate and adapt to a stressful environment, eukaryotic cells seem to have evolved a more controlled mutagenic response to the environmental stress.

DATA AVAILABILITY

All data is held by the authors and is available on request.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

FUNDING

Fondation pour la Recherche Médicale [FRM - EQU2 01903007797]; Fondation de France (to K.M.). Funding for

open access charge: Fondation pour la Recherche Médicale [FRM - EQU2 01903007797].

Conflict of interest statement. None declared.

REFERENCES

- Hanawalt, P.C. (2015) Historical perspective on the DNA damage response. *DNA Repair (Amst)*, **36**, 2–7.
- Friedberg, E.C., Walker, G.C., Siede, W., Wood, R.D., Schultz, R.A. and Ellenberger, T. (2006) In: *DNA Repair And Mutagenesis*. Amer Society for Microbiology Press, Washington, DC.
- Hoege, C., Pfander, B., Moldovan, G.-L., Pyrowolakis, G. and Jentsch, S. (2002) RAD6-dependent DNA repair is linked to modification of PCNA by ubiquitin and SUMO. *Nature*, **419**, 135–141.
- Branzei, D., Seki, M. and Enomoto, T. (2004) Rad18/Rad5/Mms2-mediated polyubiquitination of PCNA is implicated in replication completion during replication stress. *Gene. Cells*, **9**, 1031–1042.
- Branzei, D. and Szakal, B. (2016) DNA damage tolerance by recombination: molecular pathways and DNA structures. *DNA Repair (Amst)*, **44**, 68–75.
- Prakash, S., Johnson, R.E. and Prakash, L. (2005) Eukaryotic translesion synthesis DNA polymerases: specificity of structure and function. *Ann. Rev. Biochem.*, **74**, 317–353.
- Waters, L.S., Minesinger, B.K., Wiltout, M.E., D'Souza, S., Woodruff, R.V. and Walker, G.C. (2009) Eukaryotic translesion polymerases and their roles and regulation in DNA damage tolerance. *Microbiol. Mole. Biol. Rev. MMBR*, **73**, 134–154.
- Masłowska, K.H., Makiela-Dzibenska, K. and Fijalkowska, I.J. (2018) The SOS system: a complex and tightly regulated response to DNA damage. *Environment. Mol. Mutagen.*, **7**, e1002244.
- Barrett, T.C., Mok, W.W.K., Murawski, A.M. and Brynildsen, M.P. (2019) Enhanced antibiotic resistance development from fluoroquinolone persisters after a single exposure to antibiotic. *Nat. Commun.*, **10**, 1177.
- Naiman, K., Philippin, G., Fuchs, R.P. and Pagès, V. (2014) Chronology in lesion tolerance gives priority to genetic variability. *Proc. Natl Acad. Sci. USA*, **111**, 5526–5531.
- Napolitano, R., Janel-Bintz, R., Wagner, J. and Fuchs, R.P. (2000) All three SOS-inducible DNA polymerases (Pol II, pol IV and pol V) are involved in induced mutagenesis. *Embo J.*, **19**, 6259–6265.
- McIntyre, J. and Woodgate, R. (2015) Regulation of translesion DNA synthesis: posttranslational modification of lysine residues in key proteins. *DNA Repair*, **29**, 166–179.

13. Fu, Y., Pastushok, L. and Xiao, W. (2008) DNA damage-induced gene expression in *saccharomyces cerevisiae*. *FEMS Microbiol. Rev.*, **32**, 908–926.
14. Jentsch, S., McGrath, J.P. and Varshavsky, A. (1987) The yeast DNA repair gene RAD6 encodes a ubiquitin-conjugating enzyme. *Nature*, **329**, 131–134.
15. Bailly, V., Lauder, S., Prakash, S. and Prakash, L. (1997) Yeast DNA repair proteins rad6 and rad18 form a heterodimer that has ubiquitin conjugating, DNA binding, and ATP hydrolytic activities. *J. Biol. Chemist.*, **272**, 23360–23365.
16. Davies, A.A., Huttner, D., Daigaku, Y., Chen, S. and Ulrich, H.D. (2008) Activation of ubiquitin-dependent DNA damage bypass is mediated by replication protein a. *Mol. Cell*, **29**, 625–636.
17. Ulrich, H.D. and Jentsch, S. (2000) Two RING finger proteins mediate cooperation between ubiquitin-conjugating enzymes in DNA repair. *Embo J.*, **19**, 3388–3397.
18. Parker, J.L. and Ulrich, H.D. (2009) Mechanistic analysis of PCNA poly-ubiquitylation by the ubiquitin protein ligases rad18 and rad5. *EMBO J.*, **28**, 3657–3666.
19. Andersen, P.L., Xu, F. and Xiao, W. (2008) Eukaryotic DNA damage tolerance and translesion synthesis through covalent modifications of PCNA. *Cell Res.*, **18**, 162–173.
20. Che, J., Hong, X. and Rao, H. (2021) PCNA ubiquitylation: instructive or permissive to DNA damage tolerance pathways. *Biomolecules*, **11**, 1543.
21. Maslowska, K.H., Laureti, L. and Pagès, V. (2019) iDamage: a method to integrate modified DNA into the yeast genome. *Nucleic Acids Res.*, **47**, e124.
22. Johnson, R.E., Torres-Ramos, C.A., Izumi, T., Mitra, S., Prakash, S. and Prakash, L. (1998) Identification of APN2, the *saccharomyces cerevisiae* homolog of the major human AP endonuclease HAP1, and its role in the repair of abasic sites. *Genes Dev.*, **12**, 3137–3143.
23. Akada, R., Kitagawa, T., Kaneko, S., Toyonaga, D., Ito, S., Kakiyama, Y., Hoshida, H., Morimura, S., Kondo, A. and Kida, K. (2006) PCR-mediated seamless gene deletion and marker recycling in *saccharomyces cerevisiae*. *Yeast (Chichester, England)*, **23**, 399–405.
24. Alani, E., Cao, L. and Kleckner, N. (1987) A method for gene disruption that allows repeated use of URA3 selection in the construction of multiply disrupted yeast strains. *Genetics*, **116**, 541–545.
25. LeClerc, J.E., Borden, A. and Lawrence, C.W. (1991) The thymine-thymine pyrimidine-pyrimidone(6-4) ultraviolet light photoproduct is highly mutagenic and specifically induces 3' thymine-to-cytosine transitions in *escherichia coli*. *Proc. Natl Acad. Sci. USA*, **88**, 9685–9689.
26. Marsella, A., Gobbini, E., Cassani, C., Tisi, R., Cannavo, E., Reginato, G., Cejka, P. and Longhese, M.P. (2021) Sae2 and rif2 regulate MRX endonuclease activity at DNA double-strand breaks in opposite manners. *Cell Rep.*, **34**, 108906.
27. Miller, E.C., Miller, J.A. and Hartmann, H.A. (1961) N-Hydroxy-2-acetylaminofluorene: a metabolite of 2-acetylaminofluorene with increased carcinogenic activity in the rat. *Cancer Res.*, **21**, 815–824.
28. Pagès, V., Bresson, A., Acharya, N., Prakash, S., Fuchs, R.P. and Prakash, L. (2008) Requirement of rad5 for DNA polymerase zeta-dependent translesion synthesis in *saccharomyces cerevisiae*. *Genetics*, **180**, 73–82.
29. Mortensen, U.H., Bendixen, C., Sunjevaric, I. and Rothstein, R. (1996) DNA strand annealing is promoted by the yeast rad52 protein. *Proc. Natl Acad. Sci. USA*, **93**, 10729–10734.
30. Gallagher, D.N., Pham, N., Tsai, A.M., Janto, A.N., Choi, J., Ira, G. and Haber, J.E. (2020) A Rad51-independent pathway promotes single-strand template repair in gene editing. *PLoS Genet.*, **16**, e1008689.
31. Nagao, M. and Sugimura, T. (1976) Molecular biology of the carcinogen, 4-nitroquinoline 1-oxide. *Adv. Cancer Res.*, **23**, 131–169.
32. Foster, P.L. (1991) In vivo mutagenesis. *Methods Enzymol.*, **204**, 114–125.
33. Unk, I. and Daraba, A. (2014) Measuring UV-induced mutagenesis at the CAN1 locus in *saccharomyces cerevisiae*. *Bio. Protoc.*, **4**, e1272.
34. Álvarez, V., Frattini, C., Sacristán, M.P., Gallego-Sánchez, A., Bermejo, R. and Bueno, A. (2019) PCNA deubiquitylases control DNA damage bypass at replication forks. *Cell Rep.*, **29**, 1323–1335.
35. Hedglin, M. and Benkovic, S.J. (2017) Replication protein a prohibits diffusion of the PCNA sliding clamp along single-stranded DNA. *Biochemistry*, **56**, 1824–1835.
36. Buglewicz, D.J., Mussallem, J.T., Haskins, A.H., Su, C., Maeda, J. and Kato, T.A. (2020) Cytotoxicity and mutagenicity of narrowband UVB to mammalian cells. *Genes (Basel)*, **11**, 646.
37. Tang, J.Y., Hwang, B.J., Ford, J.M., Hanawalt, P.C. and Chu, G. (2000) Xeroderma pigmentosum p48 gene enhances global genomic repair and suppresses UV-induced mutagenesis. *Mol. Cell*, **5**, 737–744.