



A Kappa model for hepatic stellate cells activation by TGFB1

Matthieu Bouguéon, Pierre Boutillier, Jerome Feret, Octave Hazard, Nathalie Theret

► To cite this version:

Matthieu Bouguéon, Pierre Boutillier, Jerome Feret, Octave Hazard, Nathalie Theret. A Kappa model for hepatic stellate cells activation by TGFB1. CMSB 2021 - 19th International Conference on Computational Methods in Systems Biology, Sep 2021, Bordeaux / Virtual, France. pp.1-6. hal-03545256

HAL Id: hal-03545256

<https://inria.hal.science/hal-03545256>

Submitted on 27 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Kappa model for hepatic stellate cells activation by TGFB1

^{1,2,3}Matthieu Bouguéon, Pierre Boutillier, Jérôme Feret, Octave Hazard and Nathalie Theret.



Hepatic Stellate Cells ; the drivers of tissue repair and fibrosis




Kappa

Rule-based language for modeling systems of interacting agents (<https://kappalanguage.org/>)

Entities are graphical structures, rules are graph-rewrite directives.

Rules locally modify the state of a system




'HSC(state{quiescent}) TGFB1_2{membrane}[.], TGFB1(cell[.], state{active}) -> HSC(state{quiescent}) TGFB1_2{intern}[1], TGFB1(cell[1] state{active}) @ 'Vr' * 'fixing_cell'

Acute simulation




Number of occurrences



1 input of **TGFB** at 100 hours

Number of occurrences




60 inputs of **TGFB** each 24 hours

Conclusion

- ❖ reverted HSC (iHSC) are key regulators
- ❖ **TGFB1** is necessary to induce HSC activation but doesn't allow to reach a fibrotic state characterised by persistent high level of MFB

Hypothesis ; extracellular matrix contribute to the sustained activation of HSCs ?

A New model to develop !





Thank you for your attention