

Optimal bacterial resource allocation strategies in batch processing

Agustín Gabriel Yabo, Jean-Baptiste Caillau, Jean-Luc Gouzé

► To cite this version:

Agustín Gabriel Yabo, Jean-Baptiste Caillau, Jean-Luc Gouzé. Optimal bacterial resource allocation strategies in batch processing. 2022. hal-03710681v1

HAL Id: hal-03710681 https://inria.hal.science/hal-03710681v1

Preprint submitted on 30 Jun 2022 (v1), last revised 25 Sep 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Manuscript submitted to AIMS' Journals Volume X, Number 0X, XX 200X doi:10.3934/xx.xxxxxx

pp. **X–XX**

OPTIMAL BACTERIAL RESOURCE ALLOCATION STRATEGIES IN BATCH PROCESSING

Agustín Gabriel Yabo^{1,2}

¹Université Côte d'Azur, Inria, INRAE, CNRS, Sorbonne Université, Biocore Team Sophia Antipolis, France ²MISTEA, Université Montpellier, INRAE, Institut Agro Montpellier, France

JEAN-BAPTISTE CAILLAU³

³Université Côte d'Azur, CNRS, Inria, LJAD France

JEAN-LUC GOUZÉ⁴

⁴Université Côte d'Azur, Inria, INRAE, CNRS, Sorbonne Université, Biocore Team Sophia Antipolis, France

(Communicated by the associate editor name)

ABSTRACT. The study of living microorganisms using resource allocation models has been key in elucidating natural behaviors of bacteria, by allowing allocation of microbial resources to be represented through optimal control strategies. The approach can also be applied to research in microbial cell factories, to investigate the optimal production of value-added compounds regulated by an external control. The latter is the subject of this paper, in which we study batch bioprocessing from a resource allocation perspective. Based on previous works, we propose a simple multiscale model accounting for the dynamics of the bioreactor and intracellular composition, and we analyze its asymptotic behavior and stability. Using optimization and optimal control theory, we study the production of biomass and metabolites of interest for infinite- and finite-time horizons. The resulting optimal control problems are studied using Pontryagin's Maximum Principle and numerical methods, and the solutions found are characterized by the presence of Fuller phenomenon (producing an infinite set of switching points occurring in a finite-time window) at the junctions with a second-order singular arc. The approach, inspired in biotechnological engineering, aims to shed light upon the role of cellular composition and resource allocation during batch processing and, at the same time, poses very interesting and challenging mathematical problems.

²⁰²⁰ Mathematics Subject Classification. Primary: 37N25, 49K15; Secondary: 92C42. Key words and phrases. mathematical systems theory, nonlinear systems; optimization problems; dynamic stability; systems biology; mathematical cell model dynamics and control; modeling and identification; industrial biotechnology; optimal control; bacterial resource allocation.

This work was partially supported by ANR project Maximic (ANR-17-CE40-0024-01), Inria IPL Cosy and Labex SIGNALIFE (ANR-11-LABX-0028-01).

^{*}Corresponding author: Agustín Gabriel Yabo.

1. Introduction. The study of living microorganisms through resource allocation models has become increasingly relevant for its capacity to elucidate natural behaviors of microbia through very simple dynamical models [6, 7, 10, 11, 20, 25]. The core idea is to represent the distribution of cellular resources through optimal control strategies, based on the assumption that evolutionary processes have tuned these endogenous allocation strategies to attain nearly-optimal levels [13]. Numerous problems arise in this context, one of them being the optimal production of metabolites regulated by an external control capable of arresting bacterial growth [9]. Growth control has proven a key engineering method for several industrial applications, such as in food preservation, biofuel production, and in combating antibiotics resistance [8]. To this end, a resource allocation approach can help understand how to modify the naturally-evolved allocation strategies so as to efficiently produce such chemical compounds [5].

These biosynthetic strategies have been studied in different frameworks. The simplest case describes the interactions between intracellular proteins with minimal interplay with the environment [26, 4, 21]. The latter can be modelled by omitting the dynamics of the substrate in the medium, representing the case where bacterial exponential growth can be attained. Another relevant, more complex case is continuous bioreactors [22, 24], used extensively in industries and in cell biology research for its capacity to reach and maintain steady-state growth conditions. The latter is accomplished through an inflow of fresh medium rich in substrate and an outflow of the culture at the same volumetric flow rate, which produce a constant volume of the culture in the device. In that case, optimization studies are mostly oriented to reach such steady state in a cost-effective way. In fed-batch fermentation, the process starts with an initial volume of bacterial culture inside a bioreactor, which is progressively filled up through an inflow of rich medium, increasing the volume of the culture until it reaches a maximum level [23]. Once the maximum volume is attained, the culture evolves as a closed process, known in the field as batch processing. As no mass comes in or out of the device, the remainder of the nutrients in the medium are progressively consumed until the mass is entirely transformed into final products.

The latter is the subject of this paper, which tackles batch processing from a resource allocation perspective. The novelty of the approach lies in the multi-scale nature of the model, which considers bioreactor dynamics, related to the classical physical and chemical laws in bioprocessing; and cellular composition, taking into account the intracellular components responsible for the main biological functions of bacteria. The problem has been first posed in [26], where a simpler mathematical model of resource allocation is studied through numerical optimal control. We extend these results from an analytical perspective—both for the dynamical analysis and the optimal control study—and including the case with no metabolite synthesis as a starting point. Based on previous works [6], a simple coarse-grained self-replicator model is introduced, including a heterologous pathway for the production of a value-added chemical compound [26, 21]. Additionally, the main biological assumptions of the mechanistic bacterial model are revised, based on empirical studies of exponentially growing E. coli cultures [16]. Specifically, we consider a class of growth rate-independent proteins in the cellular composition that accounts for housekeeping proteins and non-active ribosomes, known to take up more than 50%of the cell [15]. The inclusion of this class of proteins in previous models has shown considerable improvement in the agreement between simulations and experimental

data [25]. Using mass conservation laws related to the closedness of the bioprocess, it is possible to analyze the asymptotic behavior and stability of the dynamical system, showing that, for every possible allocation strategy, all component of the system are transformed either into proteins or into metabolites, a condition later defined as *Full depletion*. Then, two main studies are performed: the biomass maximization case, representing the natural objective of wild-type (i.e. not modified) microbial cultures; and the metabolite maximization case, using the full bacterial model that includes the pathway for metabolite synthesis for industrial purposes. Both problems are analyzed in infinite time and in finite time, the latter stated as OCPs (Optimal Control Problems), which are investigated through the application of PMP (Pontryagin's Maximum Principle) [14]. While the finite-time case is suitable for representing bioprocesses with predetermined duration, the analysis of the infinite-time case becomes crucial in understanding the nature and asymptotic trend of the process. The solutions of the OCPs are characterized by the presence of Fuller's phenomenon [3], producing arcs composed of an infinite set of switching points (i.e. bangs) over a finite-time window. These optimal solutions follow a Fuller-singular-Fuller structure, similar to the one found in [25], described by a single second-order singular arc which is delimited by two Fuller's arcs at the beginning and at the end of the process. In particular, the solution of the biomass maximization case is thoroughly studied from an analytical point of view, resulting in an explicit expression of the singular control in feedback form. The results here presented are also confirmed by numerical simulations obtained with Bocop [17], an optimal control solver based on direct methods.

The paper is organized as follows: in Section 2, the dynamical model is presented, and its dynamical behavior is studied in Section 3. The biomass and product maximization cases are introduced and investigated in Sections 4 and 5, respectively. Finally, the results are discussed in Section 6.

2. Model definition.

2.1. Self-replicator model. We define a self-replicator model describing the dynamics of a microbial population growing inside a closed bioreactor. The bacterial culture has constant volume \mathcal{V}_e , measured in liters. At the beginning of the experience, there is an initial mass of substrate S inside the bioreactor, that is gradually consumed by the bacterial population, and transformed into precursor metabolites P. These precursors are intermediate metabolites used to produce proteins—such as ribosomes and enzymes—responsible for specific cellular functions; and metabolites of interest X which are excreted from the cell. The proteins forming bacterial cells are divided into three classes M, R and Q, associated to the following cellular functions:

- **Class M:** Proteins of the metabolic machinery, responsible for the uptake of nutrients S from the medium, the production of precursor metabolites P, and the synthesis of metabolites of interest X.
- **Class R:** Proteins of the gene expression machinery (such as ribosomes) actively involved in protein biosynthesis (*i.e.* in the production of proteins of classes M, Q and R).
- **Class Q:** Growth rate-independent proteins, such as housekeeping proteins responsible for cell maintenance, and ribosomes not involved in protein synthesis [15].

From a biological perspective, the production of proteins M, R and Q is catalyzed by ribosomal proteins R, and the absorption of S and synthesis of X are both catalyzed by the metabolic proteins M. This catalytic effect is represented in Figure 1 through dashed arrows. Intracellular proteins are produced at a synthesis rate V_R measured



FIGURE 1. Self-replicator model of bacterial growth representing the intracellular micro-chemical reactions behind nutrient uptake, cell growth and metabolite synthesis. Solid arrows represent flow of resources resulting from the microchemical reactions, while dashed arrows indicate a catalyzing effect (*i.e.* the presence of a protein accelerating the synthesis of another protein).

in grams per hour. The synthesis rates of proteins M, R and Q are $r_{\max}(1-u)V_R$, $r_{\max}uV_R$ and $(1-r_{\max})V_R$, respectively; where the parameter r_{\max} is a certain empirical constant imposing a maximum threshold to the rate of production of proteins M and R. The proportion of precursors dedicated to the production of growth rate-independent proteins Q is fixed, while the balance between proteins M and R is decided by the allocation control u. The latter is modelled through a time-varying function $u(t) \in [0, 1]$, where u = 0 means no production of ribosomal proteins R, and u = 1 means no production of metabolic proteins M. The control u can represent the natural allocation used by bacteria, as modelled in [6], as well as the artificially modified allocation modelled in [26], depending on the objective to be analyzed.

2.2. **Dynamical system.** The dynamics of the self-replicator system are described by

$$\begin{cases} \dot{S} = -V_M, \\ \dot{P} = V_M - V_X - V_R, \\ \dot{R} = r_{\max} u V_R, \\ \dot{M} = r_{\max} \left(1 - u\right) V_R, \\ \dot{Q} = (1 - r_{\max}) V_R, \\ \dot{X} = V_X, \end{cases}$$

where the variables S, P, R, M, Q and X represent the masses (in grams) of substrate, precursors metabolites, the gene expression machinery, the metabolic machinery, the growth rate-independent proteins and the metabolites of interest at time t, respectively. V_M , V_R and V_X are the reaction rates of the system (in grams per hour), and u(t) is the allocation control previously defined. We define the volume (in liters) of the bacterial population in the bacterial culture $\mathcal{V}_{in}(t)$ as

$$\mathcal{V}_{in} \doteq \beta (M + R + Q), \tag{1}$$

where β is a constant relating protein density and volume [2]. Definition (1) purposely neglects the mass of precursor metabolites P, which greatly simplifies the computations. The latter assumption is based on the fact that most of the mass in bacterial cells corresponds to proteins M, R and Q, as confirmed in previous studies [6]. This allows to define intracellular concentrations (in grams per liter) with respect to this volume

$$p_{in} \doteq \frac{P}{\mathcal{V}_{in}}, \qquad r_{in} \doteq \frac{R}{\mathcal{V}_{in}}, \qquad m_{in} \doteq \frac{M}{\mathcal{V}_{in}}, \qquad q_{in} \doteq \frac{Q}{\mathcal{V}_{in}}.$$
 (2)

Likewise, we define the extracellular concentrations related to the external volume

$$s_e = \frac{S}{\mathcal{V}_e}, \quad x_e = \frac{X}{\mathcal{V}_e}.$$
 (3)

We define the relative synthesis rates involved in the processes as increasing functions of the concentrations used in each reaction [15], and taking into account the catalytic effect previously described

$$v_M(s_e, m_{in}) \doteq \frac{V_M}{\mathcal{V}_{in}}, \quad v_R(p_{in}, r_{in}) \doteq \frac{V_R}{\mathcal{V}_{in}}, \quad v_X(p_{in}, m_{in}) \doteq \frac{V_X}{\mathcal{V}_{in}}.$$

From (1) and (2), we have that

$$m_{in} + r_{in} + q_{in} = \frac{1}{\beta},\tag{4}$$

which implies that the concentrations m_{in} , r_{in} and q_{in} cannot be bigger than $1/\beta$. We define the growth rate of the bacterial culture μ as

$$\mu \doteq \frac{\dot{\mathcal{V}}_{in}}{\mathcal{V}_{in}} = \beta v_R(p_{in}, r_{in}).$$

Then, the dynamical system can be expressed in terms of the concentrations as

$$\begin{aligned} \dot{s}_{e} &= -v_{M}(s_{e}, m_{in}) \frac{\mathcal{V}_{in}}{\mathcal{V}_{e}}, \\ \dot{p}_{in} &= v_{M}(s_{e}, m_{in}) - v_{X}(p_{in}, m_{in}) - v_{R}(p_{in}, r_{in})(\beta p_{in} + 1), \\ \dot{r}_{in} &= (r_{\max}u - \beta r_{in})v_{R}(p_{in}, r_{in}), \\ \dot{m}_{in} &= (r_{\max}(1 - u) - \beta m_{in})v_{R}(p_{in}, r_{in}), \\ \dot{q}_{in} &= ((1 - r_{\max}) - \beta q_{in})v_{R}(p_{in}, r_{in}), \\ \dot{\mathcal{V}}_{in} &= \beta v_{R}(p_{in}, r_{in})\mathcal{V}_{in}, \\ \dot{x}_{e} &= v_{X}(p_{in}, m_{in})\frac{\mathcal{V}_{in}}{\mathcal{V}_{e}}. \end{aligned}$$

2.3. Kinetics definition. We model the kinetics of the system by supposing that both the synthesis rates of precursors v_M and metabolites v_X are linear in the concentration of metabolic proteins m_{in} , and the protein synthesis rate v_R is linear in the concentration of active ribosomal proteins r_{in} [16]. Thus, they can be expressed as

$$\begin{aligned} v_M(s_e, m_{in}) &= w_M(s_e)m_{in}, \\ v_R(p_{in}, r_{in}) &= w_R(p_{in})r_{in}, \\ v_X(p_{in}, m_{in}) &= \gamma w_R(p_{in})m_{in}, \end{aligned}$$

where $\gamma > 0$ is a proportionality constant, which allows the metabolite synthesis rate to be expressed as $v_X(p_{in}, m_{in}) = \gamma v_R(p_{in}, r_{in}) m_{in}/r_{in}$. Such assumption implies that the bacterial cell has the same affinity to synthesize biomass and metabolites from the precursors, even if the reactions do not consume P in the same proportion. In the particular case of Michaelis-Menten kinetics, this feature is captured by the half-saturation constant [12]. The functions w_I are assumed to have the following properties:

Function $w_I(x) : \mathbb{R}_+ \to \mathbb{R}_+$ is

- Continuously differentiable w.r.t. x,
- Null at the origin: $w_I(0) = 0$,
- Strictly monotonically increasing: $w'_I(x) > 0, \forall x \ge 0$,
- Strictly concave downwards: $w_I''(x) < 0, \forall x \ge 0$,
- Upper bounded: $\lim_{x\to\infty} w_I(x) = k_I > 0.$

For numerical simulations, we resort to the particular case where the functions follow Michalis-Menten kinetics. For that case, we define

$$w_R(p_{in}) \doteq k_R \frac{p_{in}}{K_R + p_{in}}, \quad w_X(p_{in}) \doteq k_X \frac{p_{in}}{K_X + p_{in}}, \quad w_M(s_e) \doteq k_M \frac{s_e}{K_M + s_e},$$

where the values of the constants k_R , K_R , k_X , K_X , k_M and K_M are based on the literature [6, 26]. For the general case introduced in Hypothesis 2.3, we define

$$k_R \doteq \lim_{p_{in} \to \infty} w_R(p_{in}), \quad k_X \doteq \lim_{p_{in} \to \infty} w_X(p_{in}), \quad k_M \doteq \lim_{s_e \to \infty} w_M(s_e).$$

2.4. Mass fraction formulation and non-dimensionalization. We define mass fractions

$$s \doteq \beta s_e, \quad p \doteq \beta p_{in}, \quad r \doteq \frac{\beta}{r_{\max}} r_{in}, \quad m \doteq \frac{\beta}{r_{\max}} m_{in}, \quad q \doteq \beta q_{in}, \quad x \doteq \beta x_e,$$
(5)

where r and m are the mass fractions of the maximal ribosomal fraction r_{max} . Then, given that the transcription of housekeeping proteins in bacterial cells is internally auto-regulated [19], and that the mass fraction of non-translating ribosomal proteins is constant [16], we assume that the mass fraction of growth rate-independent proteins q variates mildly compared to the remaining states, and thus we fix

$$q = 1 - r_{\max},\tag{6}$$

which, replacing in (4), yields

$$m + r = 1.$$

The latter implies that the metabolic fraction can be expressed in terms of the ribosomal fraction as m = 1-r, and so the dynamical equation of m can be removed from the system. Additionally, we see that the quantity βr_{in} represents the mass fraction of translating ribosomal proteins in the cell which, using (4) and (6), has bounds $[0, r_{\text{max}}]$. Thus, its upper bound is given by the difference between the maximal total ribosomal mass fraction and the constant non-translating ribosomal mass fraction. In the literature [25], such values are empirically fixed to 0.5 and 0.07, respectively, and so the parameter r_{max} is here set to 0.43 for the numerical calculations. The biomass fraction of the bacterial culture is defined as

$$\mathcal{V} \doteq \frac{\mathcal{V}_{in}}{\mathcal{V}_e}.\tag{7}$$

We define the non-dimensional time $\hat{t} \doteq k_R r_{max} t$ and the non-dimensional functions

$$\hat{w}_R(p) = \frac{w_R(p_{in})}{k_R}, \quad \hat{w}_X(p) = \frac{w_X(p_{in})}{k_R}, \quad \hat{w}_M(s) = \frac{w_M(s_e)}{k_R}$$

so that $\lim_{p\to\infty} \hat{w}_R(p) = 1$. For the sake of simplicity, let us drop all hats from the current notation. Thus, system becomes

$$\begin{cases} \dot{s} = -w_M(s)(1-r)\mathcal{V}, \\ \dot{p} = w_M(s)(1-r) - \gamma w_R(p)(1-r) - w_R(p)r(p+1), \\ \dot{r} = (u-r)w_R(p)r, \\ \dot{\mathcal{V}} = w_R(p)r\mathcal{V}, \\ \dot{x} = \gamma w_R(p)(1-r)\mathcal{V}. \end{cases}$$
(S)

In this formulation, and using (2), (3), (4), (5) and (7), the total mass in the bioreactor can be expressed in terms of the concentrations as

$$S + P + M + R + Q + X = \frac{\mathcal{V}_e}{\beta} (s + (p+1)\mathcal{V} + x).$$
(8)

3. Model analysis.

8

Lemma 3.1. The set

$$\Gamma = \{ (s, p, r, \mathcal{V}, x) \in \mathbb{R}^5 : s \ge 0, p \ge 0, 1 \ge r \ge 0, \mathcal{V} \ge 0, x \ge 0 \}$$

is positively invariant for the initial value problem.

Proving Lemma 3.1 is standard and can be done by evaluating the vector field of (S) over the boundaries of Γ . Thus, we fix initial conditions

$$s(0) = s_0 > 0, \quad p(0) = p_0 > 0, \quad r(0) = r_0 \in (0, 1), \quad \mathcal{V}(0) = \mathcal{V}_0 > 0, \quad x(0) = 0,$$
 (IC)

where the initial concentration of metabolites x(0) is set to 0 to represent the fact that, at the beginning of the bioprocess, no metabolite has been produced. Some relations are immediate from the dynamics: as $\dot{s} \leq 0$ and $\dot{\mathcal{V}} \geq 0$ for all t, we have

$$s(t) \le s_0, \quad \mathcal{V}(t) \ge \mathcal{V}_0,$$
(9)

representing the fact that the substrate can only be consumed (and not replenished), and the biomass can only grow.

3.1. Total available mass. As typically occurs in batch processes, there is neither inflow nor outflow of mass in the bioreactor, which is reflected in the dynamics of the system though a mass conservation law. We define the constant

$$\Sigma \doteq s_0 + (p_0 + 1)\mathcal{V}_0,$$

representing the initial mass concentration in the system. It can be seen that the total mass concentration

$$z \doteq s + (p+1)\mathcal{V} + x$$

is constant for all t (as $\dot{z} = 0$). This means that

$$s + (p+1)\mathcal{V} + x = \Sigma,$$
 (First integral 1)

for all t. Thus, relation (8) and (First integral 1) show that the total mass in the system is constant and equal to $\mathcal{V}_e \Sigma/\beta$. Variables \mathcal{V} and x are maximal when the remaining variables are equal to 0, and so they are upper bounded. In particular, both $\mathcal{V}(t)$ and x(t) are decreasing w.r.t. s(t) and p(t). As neither s nor p can be negative, we have that

$$\mathcal{V}(t) + x(t) = \Sigma \tag{10}$$

when s(t) = p(t) = 0. This condition means that all the available substrate and precursor metabolites have been depleted and transformed into biomass and metabolites, which is intuitively what one would expect from system (S) for t sufficiently large. Additionally, using (9) and (First integral 1), we can obtain the following result.

Proposition 1. $\mathcal{V}(t) \in [\mathcal{V}_0, \Sigma]$, $x(t) \in [0, \Sigma - \mathcal{V}_0]$ and $p(t) \in [0, p^+]$ for all t, with $p^+ = \Sigma/\mathcal{V}_0 - 1$.

3.2. Infinite-time full depletion. Dynamics (S) shows that, under initial conditions (IC), s(t) and p(t) can only vanish asymptotically, that is, when $t \to \infty$. The latter can be proved by seeing that the derivatives of s and p can be bounded by

$$\dot{s} \ge -w_M(s)\Sigma, \quad \dot{p} \ge -w_R(p)p^+(p^++1+\gamma),$$

which means that, at worst, s and p decay exponentially (as functions $w_i(x)$ can be upper bounded by linear functions $w_i(x) \leq c_i x$), so that s(t) = p(t) = 0 cannot be reached in finite time. Accordingly, we define the depletion of s and p in an infinite-time horizon.

Definition 3.2. System (S) achieves *Full depletion* when all the substrate and the precursors are asymptotically depleted, *i.e.*

$$\lim_{t \to \infty} s(t) = \lim_{t \to \infty} p(t) = 0, \qquad (Full \ depletion)$$

3.3. Asymptotic behaviour. Now, we study the system dynamics for an infinite time $t \to \infty$. First, the case with a constant allocation $u(t) = u^* \in (0, 1)$ for all t is analyzed, and then an extension to a general allocation function is proposed.

3.3.1. Constant allocation u^* .

Theorem 3.3. For any trajectory of system (S) with initial conditions (IC) and constant allocation $u(t) = u^*$, it follows that

$$u^* - r(t))\mathcal{V}(t) = (u^* - r_0)\mathcal{V}_0.$$
 (First integral 2)

Proof. Under a constant allocation $u(t) = u^*$, the dynamics of r becomes

$$\dot{r} = (u^* - r)\mu(p)r.$$

Using dynamics (S), it is possible to see that the quantity $R_u = (u^* - r)\mathcal{V}$ is constant (as $\dot{R}_u = 0$), which yields (First integral 2).

3.3.2. General allocation u(t). Due to the boundedness of \mathcal{V} stated in Lemma 1, and the relation between \mathcal{V} and r shown in (First integral 1), we can see that any constant control u^* yields a bounded ribosomal fraction r. We extend this notion to any function u(t).

Lemma 3.4. For any trajectory of system (S) with initial conditions (IC) and any control u(t), the ribosomal concentration has bounds $r(t) \in [r^-, r^+]$ for all t, with

$$r^{-} \doteq r_0 \frac{\mathcal{V}_0}{\Sigma} > 0, \qquad r^{+} \doteq 1 - (1 - r_0) \frac{\mathcal{V}_0}{\Sigma} < 1.$$

Proof. Let us extend system (S) by defining variables $r_{\text{low}}(t)$ and $r_{\text{up}}(t)$ with dynamics

$$\dot{r}_{\text{low}} = -r_{\text{low}} w_R(p) r \le 0, \quad \dot{r}_{\text{up}} = (1 - r_{\text{up}}) w_R(p) r \ge 0,$$

 $r_{\text{low}}(0) = r_0, \quad r_{\text{up}}(0) = r_0,$

which correspond to the dynamics of r with u = 0 and u = 1 respectively, and which satisfy

$$r_{\rm low}(t) \le r(t) \le r_{\rm up}(t)$$

for all t. The latter can be easily proved by showing that the time-varying differences

$$\Delta_{\text{low}}(t) = r(t) - r_{\text{low}}(t), \quad \Delta_{\text{up}}(t) = r_{\text{up}}(t) - r(t)$$

with dynamics

$$\dot{\Delta}_{\text{low}} = (u - \Delta_{\text{low}})w_R(p)r, \quad \dot{\Delta}_{\text{up}} = (1 - u - \Delta_{\text{up}})w_R(p)r$$

are always non-negative: they satisfy $\Delta_{\text{low}}(0) = \Delta_{\text{up}}(0) = 0$ and are repulsive or (at worst) invariant at 0. Then, based on the same principle used to obtain (First integral 2), we define the quantities $R_{\text{low}} = r_{\text{low}}\mathcal{V}$ and $R_{\text{up}} = (1 - r_{\text{up}})\mathcal{V}$ which are constant (as $\dot{R}_{\text{low}} = \dot{R}_{\text{up}} = 0$), and so

$$r_{\text{low}}(t) = r_0 \frac{\mathcal{V}_0}{\mathcal{V}(t)}, \qquad r_{\text{up}}(t) = 1 - (1 - r_0) \frac{\mathcal{V}_0}{\mathcal{V}(t)},$$

for all t. As $\mathcal{V}_0 \leq \mathcal{V}(t) \leq \Sigma$ for all t, we have

$$r_{\text{low}}(t) \in \left[r_0 \frac{\mathcal{V}_0}{\Sigma}, r_0\right], \qquad r_{\text{up}}(t) \in \left[r_0, 1 - (1 - r_0) \frac{\mathcal{V}_0}{\Sigma}\right]$$

which shows that $r^{-} \leq r(t) \leq r^{+}$ for all t.

Lemma 3.4 states that, for any control u(t), the ribosomal concentration never reaches the bounds r = 0 and r = 1, and thus neither the substrate intake nor the protein synthesis is arrested. Using this fact, it can be proved that any control u(t) produces (*Full depletion*).

Theorem 3.5. Any trajectory of system (S) with initial conditions (IC) and any control u(t) achieves (*Full depletion*).

Proof. Using Lemma 3.4, it is easy to see that

$$\dot{s} \le -w_M(s)(1-r^+)\mathcal{V}_0,$$

which means that s(t) converges to 0 as $t \to \infty$. Then, this means that

$$\dot{p} \le -\gamma w_R(p)(1-r^+) - w_R(p)r^-,$$

and so p(t) also converges to 0 as $t \to \infty$.

4. The biomass maximization case. In this section, we write the problem of maximizing the biomass both for infinite time and finite time in terms of the allocation parameter u. The latter is a mathematical representation of the naturally-evolved resource allocation strategy used by bacteria in nature. Indeed, in biology it is very often assumed that bacteria during exponential growth allocate their internal resources to maximize their growth rate, thus maximizing long-term biomass production [6]. For this particular problem, we assume that no metabolite is produced, as the pathway responsible for its production is artificially engineered, and thus not present in wild-type bacteria. This is simply modeled through $\gamma = 0$. The resulting Wild-Type Bacterial Model is

$$\begin{split} \dot{s} &= -w_M(s)(1-r)\mathcal{V}, \\ \dot{p} &= w_M(s)(1-r) - w_R(p)r(p+1), \\ \dot{r} &= (u-r)w_R(p)r, \\ \dot{\mathcal{V}} &= w_R(p)r\mathcal{V}, \end{split} \tag{WTB-M}$$

4.1. Infinite-time problem.

4.1.1. *Problem formulation*. We first write the biomass maximization problem for an infinite-time horizon. The problem can be expressed as

$$\max_{u(t)} \lim_{t \to \infty} \mathcal{V}(t).$$

Since $\mathcal{V} \in [\mathcal{V}_0, \Sigma]$, applying (*Full depletion*) in (First integral 1) yields the condition

$$\lim_{t \to \infty} \mathcal{V}(t) = \Sigma$$

meaning that, in infinite time, the biomass is maximized for every control u(t). We formalize the latter in the following theorem.

Theorem 4.1. For any trajectory of system (WTB-M) with initial conditions (IC) and any control u(t), the volume $\mathcal{V}(t) \to \max \mathcal{V}(t) = \Sigma$ as $t \to \infty$.

As a consequence, using Lemma 3.3, we have the following result for constant allocations.

Corollary 1. For any trajectory of system (WTB-M) with initial conditions (IC) and constant control $u(t) = u^*$,

$$\lim_{t \to \infty} r(t) = u^* - (u^* - r_0) \frac{\mathcal{V}_0}{\Sigma}$$

This results are illustrated by the numerical simulations shown in next section.

4.1.2. Numerical simulations. Examples of trajectories confirming the analytical results are shown in Figure 2 and Figure 3, where we see that the system approaches (*Full depletion*) asymptotically in every case, thus approaching the maximal biomass value $\mathcal{V}(t) = \Sigma$. Figure 2 shows the resulting trajectories associated to the same initial conditions, when varying the allocation parameter u. On the other hand, Figure 3 illustrates the trajectories for different values of r_0 . Indeed, as Σ does not depend on the resource allocation strategy, all the available mass is transformed into biomass independently of the values of r_0 and u(t).

4.2. Finite-time problem.

4.2.1. Problem formulation. For the Biomass Maximization problem at final time t_f , we write the OCP maximizing the final bacterial volume $\mathcal{V}(t_f)$ with initial conditions (IC):

$$\begin{array}{ll} maximize & \mathcal{V}(t_f),\\ subject \ to & dynamics \ of \ (WTB-M),\\ & initial \ conditions \ (IC),\\ & u(\cdot) \in \mathcal{U}. \end{array} \tag{BM-OCP}$$

For this class of optimal control problem, with no terminal constraints, there are no controllability issues. Additionally, the dynamics is affine in the control, with the latter included in a compact and convex set (a closed interval), and it can be checked that every finite-time trajectory remains bounded. Thus, existence of a solution is guaranteed by Filippov's theorem [1]. Then, for a problem (BM-OCP) with state $\varphi \in \mathbb{R}^n$, PMP ensures that there exist $\lambda^0 \leq 0$ and an absolutely continuous mapping



FIGURE 2. Simulation of (WTB-M) with initial conditions $s_0 = 0.3$, $p_0 = 0.001$, $r_0 = 0.8$, $\mathcal{V}_0 = 0.003$, fixed final time $t_f = 50$ and different allocation functions u.



FIGURE 3. Simulation of (WTB-M) with initial conditions $s_0 = 0.3$, $p_0 = 0.001$, $\mathcal{V}_0 = 0.003$, u = 0.5, fixed final time $t_f = 50$ and different values of r_0 .

 $\lambda(\cdot)$: $[0, t_f] \to \mathbb{R}^n$, with $(\lambda(\cdot), \lambda^0) \neq (0, 0)$, such that the extremal $(\varphi, \lambda, \lambda^0, u)$

satisfies the generalized Hamiltonian system

1

$$\begin{cases} \dot{\varphi} = \frac{\partial}{\partial \lambda} H(\varphi, \lambda, \lambda^{0}, u), \\ \dot{\lambda} = -\frac{\partial}{\partial \varphi} H(\varphi, \lambda, \lambda^{0}, u), \\ H(\varphi, \lambda, \lambda^{0}, u) = \max_{u \in [0, 1]} H(\varphi, \lambda, \lambda^{0}, u), \end{cases}$$
(PMP)

for almost every $t \in [0, t_f]$. We define the adjoint states for this particular case as $\lambda = (\lambda_s, \lambda_p, \lambda_r, \lambda_V)$, and we write the Hamiltonian

$$H = -w_M(s)(1-r)\mathcal{V}\lambda_s + \left(w_M(s)(1-r) - w_R(p)r(p+1)\right)\lambda_p + w_R(p)r\mathcal{V}\lambda_\mathcal{V} + (u-r)w_R(p)r\lambda_r,$$

and the adjoint system as

$$\frac{d\lambda_s}{dt} = w'_M(s)(1-r)\left(\mathcal{V}\lambda_s - \lambda_p\right),$$

$$\frac{d\lambda_p}{dt} = \left(w'_R(p)r(p+1) + w_R(p)r\right)\lambda_p - w'_R(p)r\mathcal{V}\lambda_{\mathcal{V}} - (u-r)w'_R(p)r\lambda_r,$$

$$\frac{d\lambda_r}{dt} = -w_M(s)(\mathcal{V}\lambda_s - \lambda_p) + w_R(p)(p+1)\lambda_p - w_R(p)\mathcal{V}\lambda_{\mathcal{V}} - (u-2r)w_R(p)\lambda_r,$$

$$\frac{d\lambda_v}{dt} = w_M(s)(1-r)\lambda_s - w_R(p)r\lambda_{\mathcal{V}}.$$

Given that there are no terminal conditions on the state, the transversality conditions for the adjoint state are $\lambda(t_f) = (0, 0, 0, -\lambda_0)$, where the presence of λ_0 is due to the cost function being $\mathcal{V}(t_f)$. Since the Hamiltonian is linear in the control u, we rewrite it in the input-affine form $H = H_0 + uH_1$ with

$$H_0 = -w_M(s)(1-r)\mathcal{V}\lambda_s + \left(w_M(s)(1-r) - w_R(p)r(p+1)\right)\lambda_r$$
$$+ w_R(p)r\mathcal{V}\lambda_{\mathcal{V}} - w_R(p)r^2\lambda_r,$$
$$H_1 = w_R(p)r\lambda_r.$$

The constrained optimal control u should maximize the Hamiltonian, so the solution of (BM-OCP) is

$$u(t) = \begin{cases} 0 & if H_1 < 0, \\ 1 & if H_1 > 0, \end{cases}$$
(11)

while $u(t) = u_{\text{sing}}(t)$ is called singular whenever H_1 vanishes on a whole subinterval of $[0, t_f]$. This tells that an optimal control is a (possibly very complicated) concatenation of bangs (u = 0 and u = 1) and singular arcs, depending on the sign of the switching function H_1 . We see that, at final time, the dynamics of λ_r becomes

$$\lambda_r(t_f) = w_R(p(t_f))\mathcal{V}(t_f)\lambda_0 < 0,$$

which, using the fact that $\lambda_r(t_f) = 0$, implies that $\lambda_r > 0$ for a period $[t_f - \epsilon, t_f]$, and thus $H_1 > 0$ for a period $[t_f - \epsilon, t_f]$. As the control u should maximize the Hamiltonian, we have the following result.

Lemma 4.2. There exists ϵ such that the final bang arc of the optimal control solution of (BM-OCP) corresponds to u(t) = 0 for the time interval $[t_f - \epsilon, t_f]$.

A more detailed analysis of the optimal control solution can be done by studying the behavior of singular extremals. The latter is key in describing the structure of the optimal control, as it is typically associated to intermediate values of the control u (i.e. non-bang arcs). The singular control can indeed be expressed as a function of the state and the adjoint state $u_{sing}(t) = f(\varphi, \lambda, \lambda_0)$, where the explicit expression of f can be obtained by successively differentiating the switching function H_1 until the singular control can be solved for.

4.2.2. Singular arcs. A singular arc occurs when H_1 vanishes (as well as its successive derivatives w.r.t. time) on a subinterval $[t_1, t_2] \subset [0, t_f]$. As r is bounded and p cannot vanish in finite time, along the singular arc one has

$$\phi_0 = 0, \quad \text{with } \phi_0 \doteq \lambda_r, \tag{C1}$$

so the Hamiltonian becomes

$$H = -w_M(s)(1-r)\mathcal{V}\lambda_s + (w_M(s)(1-r) - w_R(p)r(p+1))\lambda_p + w_R(p)r\mathcal{V}\lambda_{\mathcal{V}}.$$

We differentiate (C1) and we get

$$\phi_1 = 0$$
, with $\phi_1 \doteq -w_M(s)(\mathcal{V}\lambda_s - \lambda_p) + w_R(p)(p+1)\lambda_p - w_R(p)\mathcal{V}\lambda_{\mathcal{V}}$. (C2)

Then, the Hamiltonian can be expressed as

$$H = -w_M(s)(\mathcal{V}\lambda_s - \lambda_p) - r\phi_1 + (u - r)H_1 = c$$

on the interval $[t_1, t_2]$, and so ϕ_1 becomes

$$\phi_1 = c + w_R(p)((p+1)\lambda_p - \mathcal{V}\lambda_\mathcal{V}).$$

We differentiate (C2) and we get

$$\phi_2 = 0, \quad with \ \phi_2 \doteq \frac{w_R^2(p)}{w_M(s)w_B'(p)} - 1.$$
 (C3)

Given that u does not appear in (C3), and that, based on Assumption (2.3), function $w_M(s)$ is invertible, we have the following result.

Lemma 4.3. Any singular arc is of order two and, along it, s can be expressed in terms of p through the equation $w_M(s) = x(p)$, where $x(p) : \mathbb{R}^+ \to \mathbb{R}^+$ is defined as $x(p) \doteq w_R^2(p)/w_R'(p)$.

We differentiate (C3) and we get

$$\phi_3 = 0, \quad \text{with } \phi_3 \doteq w'_M(s)w_M(s)(1-r)\mathcal{V} + x'(p)(w_M(s)(1-r) - w_R(p)r(p+1)),$$
(C4)

which shows another relation between states. The latter implies:

Lemma 4.4. Along the singular arc, r can be expressed in terms of p and \mathcal{V} as $r = \omega(p, \mathcal{V})$ with $\omega(p, \mathcal{V}) : \mathbb{R}^+ \times \mathbb{R}^+ \to (0, 1)$ defined as

$$\omega(p,\mathcal{V}) \doteq \frac{x(p)(w'_M(s)\mathcal{V} + x'(p))}{x(p)(w'_M(s)\mathcal{V} + x'(p)) + x'(p)w_R(p)(p+1))}$$

Finally, given that $\omega(p, \mathcal{V})$ does not depend on the adjoint state, the following result is straightforward.

Theorem 4.5. The singular optimal control is in feedback form, u = u(p, V).

Proof. Derivating the expression $r = \omega(p, \mathcal{V})$ and solving for u yields

$$u = \frac{1}{w_R(p)r} \frac{d\omega(p, \mathcal{V})}{dt} + r.$$

As ω only depends on p and \mathcal{V} , its derivative only depends on the state. Then, since s and r can be expressed in terms of p and \mathcal{V} , the control u becomes a function of these variables.

4.2.3. Numerical simulations. The optimal trajectories were computed with Bocop [17], which solves the OCP through a direct method. The time discretization algorithm used is Lobato IIIC (implicit, 4-stage, order 6) with 2000 time steps. Figures 4 and 5 show optimal trajectories for the same set of initial conditions and different values of t_f . Using the mass conservation law (First integral 1), the quantities are represented in the plots as fractions of the total mass in the bioreactor $\mathcal{V}_e \Sigma / \beta$. The optimal control u is characterized by the presence of chattering after and before the singular arc, as expected in singular arcs of order two. From a biological point of view, both allocation strategies prioritize the synthesis of proteins of the metabolic machinery M (red in both Figures): the singular arc takes rather small values, and a large proportion of the optimal control corresponds to a bang arc u = 0 at the end of the bioprocess. The latter strategy promotes nutrient uptake, which results in a faster depletion of the substrate. It is interesting to note that, in opposition to previous results in the literature [6, 25], the presence of the turnpike properties [18]is not assured. Intuitively, the turnpike phenomenon would cause the time interval corresponding to the singular arc to increase as the final time t_f increases. However, a quick comparison between Figures 4 and 5 shows that this is not the case, as the increase of t_f only produces a larger 0-bang arc at the end of the bioprocess. This suggests that the duration of the initial phase dedicating a fraction of the resources to ribosomal proteins is fixed and independent of the duration of the batch process. Figure 6 also show an optimal trajectory, but with a different initial ribosomal concentration. According to multiple simulations, this change shows an impact on the initial Fuller arc, that becomes perceptively larger, but has no major effects on the remaining of the process. Additionally, we note that the concentration of precursor metabolites in the bioreactor remains negligible in comparison with the other quantities, a result that matches the biological assumptions done in the modelling section.

Figure 7 illustrates optimal trajectories in the sp-plane for different final times (20, 25, 30 and 40). Each trajectory approaches the singular curve $\phi_2(p,s) = 0$ (obtained from the singular surface) through a Fuller arc, slides along it during a certain time interval, and then follows a trajectory obtained from the u = 0 arc that approaches asymptotically (*Full depletion*). Naturally, the longer the simulation of the process, the closer the final state to (*Full depletion*). These results also confirm the observations previously done: the duration in time of the singular arc is not directly related to the final time t_f . In fact, all the singular arcs start at approximately the same time instant and finish around t = 18. The independence of the initial phase from the duration of the bioprocess is coherent with the fact that bacteria allocate resources in terms of their cellular composition and the environment [6] (which, in this case, is described by the concentration of the bioprocess.



FIGURE 4. Numerical simulation of (BM-OCP) with initial conditions $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$ and $t_f = 30$. Quantities in the right plot are shown as fractions of the total mass in the bioreactor $\mathcal{V}_e \Sigma / \beta$. The final volume $\mathcal{V}(t_f)$ is at 95% of $\mathcal{V}_e \Sigma / \beta$.



FIGURE 5. Numerical simulation of (BM-OCP) with initial conditions $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$ and $t_f = 40$. Quantities in the right plot are shown as fractions of the total mass in the bioreactor $\mathcal{V}_e \Sigma / \beta$. The final volume $\mathcal{V}(t_f)$ is at 99.8% of $\mathcal{V}_e \Sigma / \beta$.

4.2.4. Alternative approach: prescribed performance in minimum time. As confirmed by the finite-time case studied in last section, the associated optimal control problem with fixed final time yields a final volume $\mathcal{V}(t_f)$ that can be viewed as a fraction (between 0 and 1) of the total mass concentration in the system Σ . Indeed, Theorem 4.1 showed that \mathcal{V} can reach its maximum value Σ only when t goes to infinity. Thus, (BM-OCP) can be reformulated so as to achieve a minimal-time tranfer between an initial state (IC) (with biomass $\mathcal{V}(0) = \mathcal{V}_0$) and a final state



FIGURE 6. Numerical simulation of (BM-OCP) with initial conditions $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.3$, $\mathcal{V}_0 = 0.003$ and $t_f = 40$. Quantities in the right plot are shown as fractions of the total mass in the bioreactor $\mathcal{V}_e \Sigma / \beta$. The final volume $\mathcal{V}(t_f)$ is at 99.8% of $\mathcal{V}_e \Sigma / \beta$.



FIGURE 7. Numerical simulation of (BM-OCP) showing different trajectories in the *sp*-plane. Initial conditions are set to $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$. In all cases, the state approaches the singular curve $\phi_2 = 0$ and slides along it.

with terminal constraint

$$\mathcal{V}(t_f) = \eta \Sigma, \tag{TC}$$

for a certain performance parameter $\eta \in [\eta_{\min}, 1)$, where $\eta_{\min} \doteq \mathcal{V}_0/\Sigma$. The reformulated minimal-time OCP with Prescribed Performance writes

$$\begin{array}{ll} \mbox{minimize} & t_f, \\ \mbox{subject to} & \mbox{dynamics of (WTB-M)}, \\ & \mbox{initial conditions (IC)}, & (\mbox{PP-OCP}) \\ & \mbox{terminal constraints (TC)}, \\ & \mbox{u}(\cdot) \in \mathcal{U}. \end{array}$$

A natural question arising from OCPs with terminal constraints is the existence of a solution. In this case, Theorem 4.1 guarantees that any final volume $\mathcal{V}(t_f) = \eta \Sigma$ can be reached in finite time, as long as $\eta \in [\eta_{\min}, 1)$. The latter ensures the existence of the solution for any $\eta \in [\eta_{\min}, 1)$. The study of the solutions of (PP-OCP) can be performed through an analogous PMP approach, with the difference that the Hamiltonian is null for every $t \in [0, t_f]$ due to the free final time t_f , and that there is no terminal constraint on $\lambda_{\mathcal{V}}$ (*i.e.* $\lambda_{\mathcal{V}}(t_f)$ is free). Given the similarity with the previously analyzed case, the computations of such optimal control solution are not explicited here. A numerical solution of the problem is shown in Figure 8.



FIGURE 8. Numerical simulation of (PP-OCP) with initial conditions $s_0 = 0.1$, $p_0 = 0.003$, $r_0 = 0.1$ and $\mathcal{V}_0 = 0.003$. The performance parameter is fixed to $\eta = 0.8$, which is achieved in $t_f = 25.1$. Quantities in the right plot are shown as fractions of the total mass in the bioreactor $\mathcal{V}_e \Sigma/\beta$.

5. The product maximization case. As done in the previous section, we approach the product maximization objective in infinite time and finite time using the full model (S) where $\gamma \in \mathbb{R}^+$.

5.1. Infinite-time problem. The problem of maximizing the product concentration at infinite time is given by the expression

$$\max_{u^*} \lim_{t \to \infty} x(t),$$

which, using (10), can be rewritten as

$$\min_{u^*} \lim_{t \to \infty} \mathcal{V}(t)$$

indicating that maximizing the metabolite concentration at infinite time equates to minimizing the biomass. While, in the previous section, the conditions (*Full depletion*) and (First integral 1) were sufficient to determine the asymptotic behavior of the system, the presence of x in this particular problem does not allow a similar resolution. An alternative approach, in order to better understand the role of cellular composition in the final objective, is to study a simplified version of the problem assuming cellular composition is at steady state (a common property of bacterial cells during exponential growth). We then propose a reduced version of the dynamical system by fixing the ribosomal concentration to a constant value r^* , which reduces the dimension of the model by one. Thus, the metabolite maximization problem is solved in terms of the parameter r^* , which represents a simpler analysis that can potentially provide an insight into the original optimization problem.

5.1.1. Constant ribosomal concentration. The system with Constant Ribosomal Concentration r^* writes

$$\begin{cases} \dot{s} = -w_M(s)(1 - r^*)\mathcal{V}, \\ \dot{p} = w_M(s)(1 - r^*) - \gamma w_R(p)(1 - r^*) - w_R(p)r^*(p+1), \\ \dot{\mathcal{V}} = w_R(p)r^*\mathcal{V}. \end{cases}$$
(CRC-M)

It can be seen that the study of the asymptotic behavior of system (S) applies to (CRC-M) as the latter is a particular case of the original one (S) with $r_0 = u^* = r^*$. We then maximize the final product x^* in terms of the constant ribosomal concentration $r^* \in [r^-, r^+]$. The latter is given by the expression

$$\max_{r^*} \lim_{t \to \infty} x(t)$$

We can see that the quantity

$$z = s + (p+1)\mathcal{V} + \gamma \frac{1-r^*}{r^*}\mathcal{V}$$

is constant. Thus,

$$\mathcal{V}^* + \gamma \frac{1 - r^*}{r^*} (\mathcal{V}^* - \mathcal{V}_0) = \Sigma$$

which, using (10), yields

$$x^* = \gamma \frac{1 - r^*}{r^*} (\mathcal{V}^* - \mathcal{V}_0).$$

Using the fact that $\mathcal{V}^* + x^* = \Sigma$ from (10), we see that x^* is monotone decreasing w.r.t. r^* , and so the ribosomal concentration maximizing the infinite-time metabolite mass is $r^* = r^-$. This is what one would expect intuitively in an infinite-time horizon, as $r^* = r^-$ favours the production of metabolic proteins M, which catalyzes the synthesis of metabolites X without arresting the production of biomass (given by the case $r^* = 0$, which cannot be attained in trajectories starting in Γ). However,

this kind of strategies might perform sub-optimally for the finite horizon case, as not having enough biomass can translate into a slow metabolite synthesis rate. Mathematically, this is represented through the presence of \mathcal{V} in the dynamical equation of x. Similar to previous results [26], a first phase dedicated to bacterial growth can also foster the production of X, which depends directly on the concentration of bacteria in the bioreactor.

5.2. Finite-time problem.

5.2.1. Problem formulation. In this section, we study the metabolite production objective in (S) for a time interval $[0, t_f]$, in which the final concentration of metabolite in the bioreactor $x(t_f)$ is maximized. While the biomass maximization objective $\mathcal{V}(t_f)$ was already studied in model (WTB-M) (representing a wild-type bacteria), it is likely that the presence of the heterologous pathway responsible for the production of x might affect the results already obtained. Thus, the two objectives are compared in model (S) from a numerical perspective. Given a fixed final time $t_f > 0$, the OCP maximizing $c\mathcal{V}(t_f) + (1-c)x(t_f)$ (with c = 0 or c = 1 depending on the objective) with initial conditions (IC) writes

$$\begin{array}{ll} maximize & c\mathcal{V}(t_f) + (1-c)x(t_f),\\ \\ subject \ to & dynamics \ of \ (\mathbf{S}),\\ & initial \ conditions \ (\mathbf{IC}),\\ \\ & and & u(\cdot) \in \mathcal{U}, \end{array} \tag{MP-OCP}$$

One can easily see that, given the dynamics of the system, applying PMP would yield a Hamiltonian linear in the control for both values of c, which means that the solution of (MP-OCP) is similar to that of (BM-OCP), given by expression (11). A numerical analysis of these results is provided in the next section.

5.2.2. Numerical simulations. The optimal trajectories were obtained following the same procedure as in the biomass maximization case. Figures 9, 10 and 11 are solutions of (MP-OCP) where the objective is the final-time product maximization $x(t_f)$ (obtained by fixing c = 0) for different final times t_f (40, 60 and 80, respectively) and with the same set of initial conditions. The metabolite synthesis rate is set to $\gamma = 0.5$. As expected, and similar to the results obtained for (BM-OCP), the optimal control takes the value u = 0 for most of the interval, representing an allocation strategy that promotes the synthesis of proteins of the metabolic machinery M, consequently catalyzing the absorption of nutrients from the medium and the production of x. Additionally, and as it can be seen across Figures 9, 10 and 11, these results do not depend on the final time t_f : the final bang u = 0 of the optimal control is always predominant in the control strategy, and becomes larger as t_f increases. The finite-time numerical results are consistent with the results obtained for the infinite-time case in Section 5.1.1, in which the ribosomal sector of the cell r should be minimized to maximize the production of x. Figure 12 shows an optimal trajectory solution of (MP-OCP) with cost function $\mathcal{V}(t_f)$. In this case, the allocation strategy is described by an initial bang u = 1 followed by a singular arc that takes up most of the optimal solution, with values near to an intermediate strategy u = 0.5; and a short bang u = 0 at the end. Such strategy leads to a

21

bacterial composition much more balanced between ribosomal and enzymatic proteins, in opposition to the metabolite production case (with c = 0 for (MP-OCP)), where most of the bacterial proteins were dedicated to the metabolic machinery. The latter behavior illustrates a natural trade-off between two opposed strategies: maximizing the number of ribosomes to prioritize the synthesis of macromolecules over the production of x and, at the same time, maximizing the enzymatic activity in order to consume the substrate in the medium as fast as possible, towards (*Full depletion*).

Under the hypothesis that the mechanisms behind the allocation of cellular resources in bacteria have been optimized to outgrow competitors, it is coherent to think that a genetically modified bacteria (e.g. able to synthesize the metabolite X) would also maximize biomass. Thus, these internal mechanisms would produce a cellular composition profile similar to the one shown in Figure 12. Indeed, this is expected to happen even for artificially engineered specimens, as, in microorganisms, natural selection occurs very rapidly (even for time windows in the order of hours) due to the strong genetic variability of bacteria and their astonishingly high doubling rate. Then, interfering with this strategy so as to obtain allocations maximizing the production of metabolites in the bioreactor (as the ones shown in Figures 9, 10 and 11) can be accomplished by externally shutting down the production of ribosomes at a certain time instant, which can be triggered by well-known biotechnological control techniques such as growth arrest [9].



FIGURE 9. Solution of (MP-OCP) for the metabolite maximization case $x(t_f)$, with $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$ and $t_f = 60$. The final product concentration $x(t_f)$ is at 35% of the total mass in the bioreactor $\mathcal{V}_e \Sigma/\beta$, while the final volume $\mathcal{V}(t_f)$ is only at 16%.

6. **Discussion.** This paper presented a mathematical study of bacterial resource allocation in batch processing, and its applications to biomass and metabolite production. A dynamical model considering the production of a value-added chemical compound is proposed, and a study of the asymptotic behavior of the system based on mass conservation laws shows that, under all possible resource allocation



FIGURE 10. Solution of (MP-OCP) for the metabolite maximization case $x(t_f)$, with $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$ and $t_f = 60$. The final product concentration $x(t_f)$ is at 62% of the total mass in the bioreactor $\mathcal{V}_e \Sigma / \beta$, while the final volume $\mathcal{V}(t_f)$ is only at 23%.



FIGURE 11. Solution of (MP-OCP) for the metabolite maximization case $x(t_f)$, with $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$ and $t_f = 60$. The final product concentration $x(t_f)$ is at 74% of the total mass in the bioreactor $\mathcal{V}_e \Sigma/\beta$, while the final volume $\mathcal{V}(t_f)$ is only at 19%.

strategies, all the substrate in the medium is consumed. Then, the particular case of a wild-type bacteria with no metabolite production is analyzed, showing that the optimal allocation propitious for biomass maximization—and thus, competitors outgrowing—is accomplished through a rather low value of the optimal control u,



FIGURE 12. Solution of (MP-OCP) for the biomass maximization case $\mathcal{V}(t_f)$, with $s_0 = 0.1$, $p_0 = 0.001$, $r_0 = 0.1$, $\mathcal{V}_0 = 0.003$ and $t_f = 60$. The final product concentration $x(t_f)$ is at 27% of the total mass in the bioreactor $\mathcal{V}_e \Sigma / \beta$, while the final volume $\mathcal{V}(t_f)$ is at 56%.

which yields a very high m/r ratio (i.e. the ratio of enzymatic to ribosomal mass fractions) in the cell throughout the bioprocess. Paradoxically, for the metabolite production case, this kind of strategies would rather maximize the production of x, while for maximizing the biomass under the presence of the heterologous pathway, a more balanced cellular composition is required. Overall, results show that optimal allocation can be accomplished through a two-phase control: a first phase of pure bacterial growth dedicated to produce as many ribosomal proteins as possible; followed by a production phase where the remainder of the feedstock is used to synthesize the compound of interest. While the first phase matches the natural bacterial behavior, the second one requires human intervention to arrest the production of ribosomes (by externally setting u = 0). The obtained two-phased external control profile is in close agreement with well-known metabolic engineering techniques used in microbial cell factory [26].

The analysis presented in this paper raises interesting questions both from mathematical and biological points of view. For instance, it would be worthwhile to further study the potential presence (or absence) of the turnpike phenomenon in the optimal control solutions. Depending on the complexity of the OCP, it is often possible to obtain an analytical proof of the exponential convergence of the singular arc to the solution of the static OCP [25], and to find an explicit link between the length of the singular arc and the duration of the bioprocess. From a biological perspective, including additional substrates could broaden the approach to represent other well-studied phenomena observed in bacterial growth. For example, under the presence of different nutrients in the culture, bacteria tend to favor (i.e. consume first) those that are easier to metabolize, which is a phenomenon known as diauxic growth. This behavior has been extensively studied from an optimal control viewpoint, but without taking into consideration cellular composition. Formulating more comprehensive dynamical models and studying their associated OCPs can be instrumental in understanding natural allocation strategies in microbial growth, and in engineering synthetic control schemes targeting industrial objectives.

Acknowledgments. The authors thank Hidde de Jong (Microcosme team, Inria Grenoble – Rhône-Alpes) for many discussions on the biological context. We also appreciate the help of Sacha Psalmon and Baptiste Schall, from Polytech Nice Sophia, for the numerical simulations and the production of the online example in the control toolbox gallery.¹

REFERENCES

- A. A. Agrachev and Y. Sachkov, Control Theory from the Geometric Viewpoint, vol. 87, Springer Science & Business Media, 2013.
- [2] M. Basan, M. Zhu, X. Dai, M. Warren, D. Sévin, Y.-P. Wang and T. Hwa, Inflating bacterial cells by increased protein synthesis, *Molecular systems biology*, **11** (2015), 836.
- [3] V. Borisov, Fuller's phenomenon, Journal of Mathematical Sciences, 100 (2000), 2311–2354.
- [4] E. Cinquemani, F. Mairet, I. Yegorov, H. de Jong and J.-L. Gouzé, Optimal control of bacterial growth for metabolite production: The role of timing and costs of control, in 2019 18th European Control Conference (ECC), IEEE, 2019, 2657–2662.
- [5] H. de Jong, J. Geiselmann and D. Ropers, Resource reallocation in bacteria by reengineering the gene expression machinery, *Trends in microbiology*, 25 (2017), 480–493.
- [6] N. Giordano, F. Mairet, J.-L. Gouzé, J. Geiselmann and H. De Jong, Dynamical allocation of cellular resources as an optimal control problem: novel insights into microbial growth strategies, *PLoS computational biology*, **12** (2016), e1004802.
- [7] S. Hui, J. M. Silverman, S. S. Chen, D. W. Erickson, M. Basan, J. Wang, T. Hwa and J. R. Williamson, Quantitative proteomic analysis reveals a simple strategy of global resource allocation in bacteria, *Molecular systems biology*, **11** (2015), 784.
- [8] L. Huo, J. J. Hug, C. Fu, X. Bian, Y. Zhang and R. Müller, Heterologous expression of bacterial natural product biosynthetic pathways, *Natural Product Reports*, **36** (2019), 1412– 1436.
- [9] J. Izard, C. D. G. Balderas, D. Ropers, S. Lacour, X. Song, Y. Yang, A. B. Lindner, J. Geiselmann and H. De Jong, A synthetic growth switch based on controlled expression of RNA polymerase, *Molecular systems biology*, **11** (2015), 840.
- [10] G. Jeanne, A. Goelzer, S. Tebbani, D. Dumur and V. Fromion, Dynamical resource allocation models for bioreactor optimization, *IFAC-PapersOnLine*, **51** (2018), 20–23.
- [11] F. Mairet, J.-L. Gouzé and H. de Jong, Optimal proteome allocation and the temperature dependence of microbial growth laws, NPJ systems biology and applications, 7 (2021), 1–11.
- [12] L. Michaelis, M. L. Menten et al., Die kinetik der invertinwirkung, Biochem. z, 49 (1913), 352.
- [13] D. Molenaar, R. Van Berlo, D. De Ridder and B. Teusink, Shifts in growth strategies reflect tradeoffs in cellular economics, *Molecular systems biology*, 5 (2009), 323.
- [14] L. S. Pontryagin, Mathematical theory of optimal processes, Routledge, 2018.
- [15] M. Scott, S. Klumpp, E. Mateescu and T. Hwa, Emergence of robust growth laws from optimal regulation of ribosome synthesis, *Molecular Systems Biology*, **10** (2014), 747.
- [16] M. Scott, C. W. Gunderson, E. M. Mateescu, Z. Zhang and T. Hwa, Interdependence of cell growth and gene expression: origins and consequences, *Science*, **330** (2010), 1099–1102.
- [17] I. S. Team Commands, Bocop: an open source toolbox for optimal control, http://bocop.org, 2017.
- [18] E. Trélat and E. Zuazua, The turnpike property in finite-dimensional nonlinear optimal control, Journal of Differential Equations, 258 (2015), 81–114.
- [19] A. Y. Weiße, D. A. Oyarzún, V. Danos and P. S. Swain, Mechanistic links between cellular trade-offs, gene expression, and growth, *Proceedings of the National Academy of Sciences*, 112 (2015), E1038–E1047.
- [20] A. G. Yabo, Optimal resource allocation in bacterial growth: theoretical study and applications to metabolite production, PhD thesis, Université Côte d'Azur, 2021.

24

¹ct.gitlabpages.inria.fr/gallery

- [21] A. G. Yabo, J.-B. Caillau and J.-L. Gouzé, Singular regimes for the maximization of metabolite production, in 2019 IEEE 58th Conference on Decision and Control (CDC), IEEE, 2019, 31–36.
- [22] A. G. Yabo, J.-B. Caillau and J.-L. Gouzé, Optimal bacterial resource allocation: metabolite production in continuous bioreactors, *Mathematical Biosciences and Engineering*, 17 (2020), 7074–7100.
- [23] A. G. Yabo, J.-B. Caillau and J.-L. Gouzé, Optimal allocation of bacterial resources in fedbatch reactors, in 2022 European control conference (ECC), IEEE, 2022, Accepted for publication.
- [24] A. G. Yabo and J.-L. Gouzé, Optimizing bacterial resource allocation: metabolite production in continuous bioreactors, *IFAC-PapersOnLine*, 53 (2020), 16753–16758.
- [25] A. G. Yabo, J.-B. Caillau, J.-L. Gouzé, H. de Jong and F. Mairet, Dynamical analysis and optimization of a generalized resource allocation model of microbial growth, SIAM Journal on Applied Dynamical Systems, 21 (2022), 137–165.
- [26] I. Yegorov, F. Mairet, H. De Jong and J.-L. Gouzé, Optimal control of bacterial growth for the maximization of metabolite production, *Journal of mathematical biology*, 78 (2019), 985–1032.

Received xxxx 20xx; revised xxxx 20xx; early access xxxx 20xx.

E-mail address: agustin.yabo@inrae.fr E-mail address: jean-baptiste.caillau@univ-cotedazur.fr E-mail address: jean-luc.gouze@inria.fr