



HAL
open science

Ensemble forecasts in reproducing kernel Hilbert space family: dynamical systems in Wonderland

Berenger Hug, Etienne Mémin, Gilles Tissot

► **To cite this version:**

Berenger Hug, Etienne Mémin, Gilles Tissot. Ensemble forecasts in reproducing kernel Hilbert space family: dynamical systems in Wonderland. 2022. hal-03740500

HAL Id: hal-03740500

<https://inria.hal.science/hal-03740500>

Preprint submitted on 30 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ensemble forecasts in reproducing kernel Hilbert space family: dynamical systems in Wonderland

Bérenger Hug¹, Étienne Mémin¹, and Gilles Tissot¹

¹Inria/IRMAR Campus de Beaulieu 35042 Rennes Cedex

July 29, 2022

Abstract

A methodological framework for ensemble-based estimation and simulation of high dimensional dynamical systems such as the oceanic or atmospheric flows is proposed. To that end, the dynamical system is embedded in a family of reproducing kernel Hilbert spaces with kernel functions driven by the dynamics. This family is nicknamed Wonderland for its appealing properties. In Wonderland the Koopman and Perron-Frobenius operators are unitary and uniformly continuous. This property warrants they can be expressed in exponential series of diagonalizable bounded infinitesimal generators. Access to Lyapunov exponents and to exact ensemble based expressions of the tangent linear dynamics are directly available as well. Wonderland enables us the devise of strikingly simple ensemble data assimilation methods for trajectory reconstructions in terms of constant-in-time linear combinations of trajectory samples. Such an embarrassingly simple strategy is made possible through a fully justified superposition principle ensuing from several fundamental theorems.

1 Introduction

Forecasting the state of geophysical fluids has become of paramount importance for our daily life either at short time scale, for weather prediction, or at long time scale, for climate study. There is in particular a strong need to provide reliable likely scenarios for probabilistic forecasts and uncertainty quantification. To that end, methods based on linear combinations of an ensemble of numerical simulations are becoming more and more ubiquitous. Ensemble forecasting and data assimilation techniques, where the linear coefficients are further constrained through partial observations of the system and Gaussian stochastic filtering strategies, are good examples developed in meteorological centers for weather forecast operational routine use. However, although efficient in practice, this linear superposition principle of solutions remains theoretically questionable for nonlinear dynamics.

Very recently, such techniques have been further coupled with machine learning to characterize surrogate dynamical models from long time-series of observations. This includes neural networks, analog forecasting, reservoir computing, kernel methods to name only a few of the works of this wide emerging research effort [12, 15, 34, 43, 49, 52, 51, 53, 65]. To characterize a surrogate model of the dynamics, these techniques are built upon principles such as delay coordinates embedding, empirical basis functions, or sparse linear representations. The difficulty here is to get sufficiently long data series to represent the manifold associated to a given observable of the system as well as to exhibit an intrinsic representation of the whole system that does not depend on parameters such as the discrete time-stepping, the integration time window, or the initial conditions. Looking for such intrinsic representation often leads to focus (at least theoretically) on reversible ergodic systems.

Spectral representation of the Koopman operator [40] or of its infinitesimal generator constitutes one of the most versatile approaches to achieve this objective as they enable in theory to extract intrinsic eigenfunctions of the system dynamics. Koopman operator or its dual, the Perron-Frobenius operator, have attracted a great deal of attention in the operator-theoretic branch of ergodic theory [30] as well as in data science for the study of data driven modeling of dynamical systems. Several techniques, starting from [28] and [46], have been proposed to estimate such spectral representation. Most of them rely either on longtime harmonic averages [46, 17, 25, 24] or on finite-dimensional approximations of the Koopman operator such as the dynamic modes representation and its extensions [54, 58, 61, 63, 64, 41, 27, 19, 50, 22, 6] or on Galerkin approximation and delay embedding [13, 32, 31]. These methods enable performing spectral projection from long time series of measured observables of the system. The practical linear nature of the *compositional* Koopman operator, defined for any observable, f , of a dynamical system $\Phi_t(x)$ with initial condition x , as $(U_t f)(x) := f \circ \Phi_t(x)$, is nevertheless hindered by its infinite dimension. Due to this, it often exhibits a spectrum with a continuous component [47] and is not amenable to diagonalization nor to any sparsity assumption. Finite dimensional approximations and the direct use of matrix algebra numerical methods raises immediate questions on the pertinence of the associated estimation. To tackle these difficulties several techniques combining regularization, compactification of the generators, and reproducing kernel Hilbert space (RKHS) to work in spaces of smooth functions have been recently proposed [25, 24, 26]. They are possibly associated with forecast strategies [33, 1, 18] and transfer operator estimations [38].

Working with RKHS brings convenient regular basis functions together with approximation convergence guaranties. However, the dynamical systems need still to be ergodic, and invertible to ensure simple point spectrum and an associated system of orthogonal basis eigenfunctions. Measure invariance and invertibility warrants that the associated Koopman/Perron-Frobenius operators are dual unitary operators with an exponential expansion expression akin to Fourier series [30]. Note also that state-of-the-art estimation techniques based on Dynamic Mode Decom-

position or Hankel matrix [3, 16] require very long time series to converge to actual Koopman eigenvalues [66]. This is all the more detrimental for high-dimensional state-spaces. Recent reviews on modern Koopman analysis applications and their related algorithmic developments from measurements time series can be found in [14, 11, 55].

In this study we will explore the problem from a different angle. Instead of working with long time series of data, we will work with a set of realizations of a given dynamical system. The objective will be to learn this dynamics locally in the phase space to synthesize in a fast way new trajectories of the system through a theoretically well justified superposition principle. Attempts to use ensemble methods to estimate the transition matrix representation of a dynamical system in view of performing data assimilation has been explored in [57] within an ensemble least squares framework. To perform an efficient estimation inherently bound to the dynamics, we propose to embed the ensemble members in a set of time-evolving RKHS defining a family of spaces. This setting is designed to deal with large-scale systems such as oceanic or meteorological flows, for which it is out of the question to explore the whole attractor (if any), neither to run very long time simulations calibrated on real-world data of a likely transient system due to climate change. The present work relies on the fact that the kernel functions, usually called feature maps, between the native space and the RKHS family are transported by the dynamical system. This creates, at any time, an isometry between the evolving RKHS and the RKHS at a given initial time. Instead of being attached to a system state, the kernel is now fully associated with the dynamics and new ensemble members embedded in the RKHS family at a given time can be very simply forecast at a further time. This framework displays striking properties – which affectionately makes us give to the RKHS family the nickname of Wonderland. For invertible dynamical systems the Koopman and Perron-Frobenius operators in Wonderland are unitary, like in L^2 . However, they are in addition uniformly continuous, with bounded generators, and diagonalizable. As such they can be rigorously expanded in exponential form and their eigenfunctions provide an intrinsic orthonormal basis system related to the dynamics. This set of analytical properties immediately brings practical techniques to estimate Koopman eigenfunctions or essential features of the dynamics. Evaluations of these Koopman eigenfunctions at the ensemble members are directly available. Finite-time Lyapunov exponents associated with each Koopman eigenfunction are easily accessible on the RKHS family as well and can be used to get practical predictability time of the system. Expressions relating Koopman eigenfunctions to the tangent linear dynamics and its adjoint are also immediately available for data assimilation. These expressions, ensuing from the fundamental results listed previously, correspond in their simpler forms to the ensemble-based finite difference approximations intensively used in ensemble methods. Beyond that, Wonderland leads finally to a theoretically well grounded superposition principle (for nonlinear dynamics) enabling the devise of embarrassingly simple methods for trajectory reconstructions in terms of constant-in-time linear combinations of trajectory samples.

The embedding of the trajectories in a time-evolving RKHS which follows the dynamical system can be related to concepts of local kernels [10], embedding in dynamical coordinates [7] or embedding of changing data over the parameter space as described in [21]. In our case, we further exploit the relations with the Koopman operator together with an associated isometry, which allows us to define simple practical schemes for the setting of the time-evolving kernels and the modal information that can be extracted from them.

This paper is organized as follows. We start briefly recalling the definition and some properties of RKHS in the next section. These definitions are complemented by A which details and adapts several known results to our functional setting. The embedding of the Koopman and Perron-Frobenius operators in Wonderand are then presented in sections 3 and 4. This is followed by the enunciation of our main results. The main theorems and properties are demonstrated in section 4 and in the appendixes. Numerical results for a quasi-geostrophic barotropic ocean model in an idealistic north Atlantic configuration will be presented in section 5 to support the pertinence of the proposed approach.

2 Reproducing kernel Hilbert spaces

A RKHS, \mathcal{H} , is a Hilbert space of smooth complex functions $f : E \mapsto \mathbb{C}$ defined over a non empty abstract set E on which a positive definite kernel and a kernel-based inner product, $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ can be defined. Throughout this work, we will consider the set E to be a topological compact set as we will work with an integral compact operator from which a convenient functional description can be set. RKHS possess remarkable properties, which make their use very appealing in statistical machine learning applications and interpolation problems [9, 23]. The kernels from which they are defined have a so called “reproducing property”.

Definition 1 (Reproducing kernel) Let \mathcal{H} be a Hilbert space of \mathbb{C} -valued functions defined on a non-empty compact topological space E . A map $k : E \times E \mapsto \mathbb{C}$ is called a reproducing kernel of \mathcal{H} if it satisfies the following principal properties: $\forall x \in E$,

- membership of the evaluation function $k(\cdot, x) \in \mathcal{H}$,
- reproducing property $\forall f \in \mathcal{H}, \langle f, k(\cdot, x) \rangle_{\mathcal{H}} = f(x)$.

The last property, provides an expression of the kernel as $k(x, y) = \overline{k(y, x)} = \langle k(\cdot, y), k(\cdot, x) \rangle_{\mathcal{H}}$, which is Hermitian – with $\bar{\bullet}$ denoting complex conjugate –, positive definite and associated with a continuous evaluation function $\delta_x f = \langle f, k(\cdot, x) \rangle_{\mathcal{H}} = f(x)$. The continuity of the Dirac evaluation operator is indeed sometimes taken as a definition of RKHS. By the Moore-Aronszajn theorem [4], the kernel k defines uniquely the RKHS, \mathcal{H} , and vice versa. The set spanned by the feature maps $\text{Vect}\{k(\cdot, x), x \in$

$E\}$, is dense in $(\mathcal{H}, \|\cdot\|_{\mathcal{H}})$. We note also that useful kernel closure properties enable to define kernels through operations such as addition, Schur product, and function composition [9]. Besides, RKHS can be meaningfully characterized through integral operators, leading to an isometry with $L^2_{\mathbb{C}}(E, \nu)$ the space of square integrable functions defined on a compact metric space E with finite measure ν [23].

Integral kernel operators

Let $k : E \times E \mapsto \mathbb{C}$ be $C^1(E \times E)$, Hermitian, and positive definite, with continuous derivative on both arguments, and let the map $\mathcal{L}_k : L^2_{\mathbb{C}}(E, \nu) \mapsto L^2_{\mathbb{C}}(E, \nu)$ be defined as:

$$(\mathcal{L}_k f)(x) = \int_E k(x, y) f(y) \nu(dy). \quad (1)$$

This operator, which must be understood within the composition with the continuous inclusion $i : C^0(E, \mathbb{C}) \hookrightarrow L^2_{\mathbb{C}}(E, \nu)$, can be shown to be well defined, positive definite, compact and self-adjoint [23]. The range of this operator is assumed to be dense in $L^2_{\mathbb{C}}(E, \nu)$. From Mercer's theorem [39], the feature maps $k(\cdot, x)$ span a RKHS defined through the eigenpairs $(\mu_i, \varphi_i)_{i \in \mathbb{N}}$ of the kernel k :

$$\mathcal{H} := \left\{ f \in L^2_{\mathbb{C}}(E, \nu), f = \sum_{i=0}^{\infty} a_i \varphi_i : \sum_i \frac{|a_i|^2}{\mu_i} < \infty \right\}, \quad (2)$$

with no null eigenvalues since we have assumed that the kernel range is dense in $L^2_{\mathbb{C}}(E, \nu)$. The rank of the kernel (number of – non-zero – eigenvalues) corresponds to the dimension of \mathcal{H} , and will potentially be infinite in this work, as E will be a subspace of functions. The RKHS \mathcal{H} is a space of smooth functions with decreasing coefficients. In fact, for all $u \in E$, there exists a constant $C > 0$ for which $\|\partial_u f\|_{L^2_{\mathbb{C}}(E, \nu)} \leq C \|f\|_{\mathcal{H}}$ (Theorem 7 – A.2). We may also define uniquely a square-root symmetric isometric bijective operator $\mathcal{L}_k^{1/2}$ between $L^2_{\mathbb{C}}(E, \nu)$ and \mathcal{H} (A.3). This operator enables to go from $L^2_{\mathbb{C}}(E, \nu)$ to \mathcal{H} by increasing the functions regularity while its inverse lowers the function regularity by bringing them back to $L^2_{\mathbb{C}}(E, \nu)$. Both operators are bounded. The injection $j : \mathcal{H} \mapsto L^2_{\mathbb{C}}(E, \nu)$ is continuous and compact (Theorem 8 – A.3) and $j(\mathcal{H})$ is dense in $L^2_{\mathbb{C}}(E, \nu)$ (Prop. 11 – A.3). These statements are precisely recalled in A.

3 Dynamical systems on a RKHS family

We consider an invertible nonsingular dynamical system $X(t) = \Phi_t(X_0)$, defined from a continuous flow, Φ_t on a compact invariant metric phase space manifold, Ω , (i.e. $\Phi_t^{-1}(\Omega) = \Omega, \forall t \in \mathbb{R}^+$) of time evolving vector functions over a spatial support Ω_x . The functions $X : \mathbb{R}^+ \times \Omega_x \mapsto \mathbb{R}^d$ with $X \in C^p, p \geq 1$, are solutions of the following d -dimensional differential system:

$$\begin{cases} \partial_t X(t, x) = \mathcal{M}(X(t, x)), \text{ with } X(t) \in \Omega, \forall t > 0, \\ X(0, x) = X_0(x). \end{cases} \quad (3)$$

The differential operator \mathcal{M} is assumed C^1 , and in particular $\sup_{X \in \Omega} \partial_X \mathcal{M}(X) < \infty$ (since Ω is compact). We consider the measure space (Ω_x, d) where d is the Lebesgue measure on Ω_x . We note as $L^2(\Omega_x, \mathbb{R}^d)$ the space of the square-integrable functions on (Ω_x, d) and $L^2(\Omega_x, \mathbb{R}^d) := \{f = (f_1, \dots, f_d) : \Omega_x \mapsto \mathbb{R}^d : \text{for all } 1 \leq i \leq d \ f_i \in L^2(\Omega_x, \mathbb{R})\}$ the vector space of square integrable functions on Ω_x . We note $\|\cdot\|_{L^2}$ the norm associated with $L^2(\Omega_x, \mathbb{R}^d)$ and given by $\|f\|_{L^2}^2 := \sum_{i=1}^d \|f_i\|_{L^2(\Omega_x, \mathbb{R})}^2$ for all $f \in L^2(\Omega_x, \mathbb{R}^d)$. The set Ω is included in $L^2(\Omega_x, \mathbb{R}^d)$. The system (3) is assumed to admit a finite invariant measure ν on Ω (note that from invertibility the measure is also nonsingular with $\nu(\Phi_t^{-1}(A)) = 0, \forall A \in \Omega$ such that $\nu(A) = 0$). The system's observables are square integrable measurable complex functions with respect to measure ν . They belong to the Hilbert space $L^2_{\mathbb{C}}(\Omega, \nu)$ with the inner product $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}$ given for f and $g \in L^2_{\mathbb{C}}(\Omega, \nu)$ by

$$\langle f, g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} := \int_{\Omega} f(y) \overline{g(y)} \nu(dy).$$

Depending on the context $X(t)$ or X_t will denote either an element of Ω or the function $X(t) : \Omega_x \rightarrow \mathbb{R}^d$ such that $x \mapsto X(t, x)$.

In this work we place ourselves in a situation where n realizations (called ensemble members) of this dynamical system, $\{X_t^{(i)}, i = 1, \dots, n\}$, generated from different initial conditions up to time t , are available. For all time $t \geq 0$, we denote by $\Omega_t \subset L^2(\Omega_x, \mathbb{R}^d)$ the space spanned by the realizations $X_t^{(i)}$ with $1 \leq i \leq n$. Defining at each time t , from this ensemble, a positive Hermitian kernel $k_{\Omega_t} : \Omega_t \times \Omega_t$, there exists a unique associated RKHS \mathcal{H}_t . In the following for the sake of concision the kernels k_{Ω_t} will be denoted by k_t to refer to the dependence on the set Ω_t . For all $t \geq 0$, we will use the notation $X_t = \Phi_t(X_0)$ and $(\mathcal{H}_t, \langle \cdot, \cdot \rangle_{\mathcal{H}_t}, \|\cdot\|_{\mathcal{H}_t})$ for the RKHS associated with the kernel k_t defined on $\Omega_t \times \Omega_t$. The kernels are assumed to be $C^2(\Omega_t \times \Omega_t)$ and as a consequence as shown in appendix A, the associated feature maps have derivatives in \mathcal{H}_t . The RKHS \mathcal{H}_t for all time $t \geq 0$ forms a family of Hilbert spaces of complex functions, each of them equipped with their own inner product $k_t(Y_t, X_t)$, for all functions $X_t, Y_t \in \mathcal{H}_t$. At any time, a measurable function of the system state, usually often referred to as an observable, $f(X_t)$, belonging to the RKHS \mathcal{H}_t can be described as a linear combination of the feature maps $\{k_t(\cdot, X_t^{(i)}), i = 1, \dots, n, t \in [0, T]\}$ and any observable trajectory $(f(X_t))_{t \geq 0}$ can be locally described as the linear combinations of time varying features maps of $(\mathcal{H}_t)_{t \geq 0}$. As it will be shown, the features maps of this RKHS family can be expressed on a time evolving orthonormal systems of basis functions, connected to each other through an exponential form and given by the eigenfunctions of the infinitesimal generator of a ‘‘Koopman-like’’ operator defined on the RKHS family. The RKHS family $\mathcal{W} = (\mathcal{H}_t)_{t \geq 0}$ is here nicknamed Wonderland because of its very appealing properties. In the following, we shall present a summary of the mathematical results associated to the Koopman operator in Wonderland.

4 Koopman operator in Wonderland

So far we did not yet give any precise definition of the kernels associated to the RKHS family \mathcal{H}_t . In order to define them from a known *a priori* initial kernel, we first introduce an evolution operator operating on the feature maps. An isometric property of this operator on the RKHS family, will then enable us to fully define the kernels along time. In the setting we are dealing with, the kernels k_t associated to the RKHS \mathcal{H}_t are defined from an operator $\mathcal{U}_t : \mathcal{H}_0 \mapsto \mathcal{H}_t$ such that

$$\mathcal{U}_t k_0(\cdot, X_0) = k_t(\cdot, \Phi_t(X_0)), \quad (4)$$

which transports the kernel feature maps on the RKHS family by composition with the system's dynamics. This operator, and more specifically its infinitesimal operator, will enable us to define the feature maps of \mathcal{H}_t from an initial feature maps on \mathcal{H}_0 . As will be fully detailed in section 4.1, we will see that the operator \mathcal{U}_t is indeed directly related to the restriction on \mathcal{H}_t of the Koopman operator U_t on a bigger RKHS space, \mathcal{H} , associated to a fixed kernel defined on the whole phase space, Ω . For that reason \mathcal{U}_t is referred to in the following as the Koopman operator in Wonderland. The global kernel k (respectively the associated RKHS) is tightly bound to the time evolving kernels k_t (respectively $(\mathcal{H}_t)_{t \geq 0}$). The feature map evolution expression of \mathcal{U}_t inherits from the relations between kernels k and k_t and ensues in particular from a dual form of Koopman operator U_t when applied on the feature maps of the global RKHS space, \mathcal{H} (Remark 1):

$$U_t [k(\cdot, X)] = k(\Phi_t(\cdot), X) = k(\cdot, \Phi_t(X)).$$

The Koopman operator U_t on \mathcal{H} , and its adjoint the Perron-Frobenius P_t exhibits some remarkable properties. The operators U_t and P_t are unitary in $L^2_{\mathbb{C}}(\Omega, \nu)$ (Prop. 1) and have the striking property to be uniformly continuous in $L^2_{\mathbb{C}}(\Omega, \nu)$ (i.e. with bounded generators – Theorem 2). As such, they can be expanded in a uniformly converging exponential series. Nevertheless, it must be outlined that the fixed kernel $k(x, y)$ and consequently \mathcal{H} are in practice only partially accessible as it is defined on the whole manifold of the dynamics and such expansion cannot be directly used. As a matter of fact, this would require a huge number of points obtained either from a great number of short-time trajectories or from very long time numerical integrations. A local representation of the RKHS family \mathcal{W} is on the other hand much easier to infer in practice through the time evolution of ensembles Ω_t and kernels k_t . As we shall see it the operator, \mathcal{U}_t , on \mathcal{W} inherits a lot of the properties of U_t and in particular, a related form of exponential series expansion (Theorem 1).

The Koopman operator in Wonderland, \mathcal{U}_t , defines an isometry from \mathcal{H}_0 to \mathcal{H}_t (Theorem 3)

$$\begin{aligned} \left\langle k_t(\cdot, \Phi_t(Y_0)), k_t(\cdot, \Phi_t(X_0)) \right\rangle_{\mathcal{H}_t} &= \left\langle k_0(\cdot, Y_0), k_0(\cdot, X_0) \right\rangle_{\mathcal{H}_0}, \\ k_t(\Phi_t(X_0), \Phi_t(Y_0)) &= k_0(X_0, Y_0). \end{aligned} \quad (5)$$

Informally, this isometry, can be guessed from definition (4) and a unitarity character of \mathcal{U}_t (Theorem 3, Prop.6) inherited from the unitarity in $L^2_{\mathbb{C}}(\Omega, \nu)$ of the Koopman operator and of its adjoint, the Perron-Frobenius operator. This property is of major practical interest as it allows us to define the kernels of the Wonderland RKHS family from a given initial kernel fixed by the user. The kernels remain constant along the system trajectories. Alternatively, an explicit form of the feature maps can be obtained from an adjoint transport equation associated to the infinitesimal generator of the Koopman operator in Wonderland. Nevertheless, the isometry is far more straightforward to use to set the kernel evolution. This isometry will reveal also very useful for the data assimilation and trajectories reconstruction problems investigated in section 5. Strikingly, we have even more than this kernel isometry. An evolution operator $A_{U,t} : \mathcal{H}_t \rightarrow L^2_{\mathbb{C}}(\Omega_t, \nu)$, associated to the infinitesimal generator of Koopman operator U_t can also be defined as

$$A_{U,t} k_t(\cdot, X_t) := \partial_{\mathcal{H}(\cdot)} k_t(\cdot, X_t), \quad (6)$$

where $\partial_u k_t(\cdot, X_t)$ stands for the directional derivative along $u \in \Omega$ of function $k(\cdot, X_t)$. This operator that will be shown to be bounded (Prop.7) and skew-symmetric (Prop.8) for the inner product of $L^2_{\mathbb{C}}(\Omega, \nu)$ plays the role of an infinitesimal generator on \mathcal{W} and enables us expressing an exponential expansion of \mathcal{U}_t .

Theorem 1 (Wonderland spectral representation) *For a measure preserving invertible dynamical system of the form (3), assuming a C^2 initial kernel, the evolution operator $A_{U,t} : \mathcal{H}_t \mapsto L^2_{\mathbb{C}}(\Omega_t, \nu)$ defined in (6), and which corresponds to the infinitesimal generator of the Koopman operator in the Wonderland RKHS family $(\mathcal{H}_t)_{t \geq 0}$ (4) can be diagonalized, at any time $t \geq 0$, by an orthonormal basis $(\psi_\ell^t)_\ell$ of \mathcal{H}_t such that*

$$A_{U,t} \psi_\ell^t(X_t) = \lambda_\ell j \circ \psi_\ell^t(X_t),$$

where j is the injection $j : \mathcal{H}_t \mapsto L^2_{\mathbb{C}}(\Omega, \nu)$. We have in addition the following relation between the orthonormal basis systems along time:

$$\forall t \geq 0 \quad \psi_\ell^t(X_t) = \exp(t \lambda_\ell) \psi_\ell^0(X_0), \quad (7)$$

with $X_t = \Phi_t(X_0)$. Furthermore, the purely imaginary eigenvalues $(\lambda_\ell)_\ell$ do not depend on time.

This theorem, which constitutes our main result, provides us a time-evolving system of orthonormal bases of the RKHS family. It brings us the capability to express any observable of the system in terms of bases that are intrinsically linked to the dynamics and related to each other by an exponential relation. The eigenvalues of $A_{U,t}$ are purely imaginary since this operator is skew-symmetric in $L^2_{\mathbb{C}}(\Omega_t, \nu)$. Remarkably, the eigenvalues of each $A_{U,t}$ do not depend on time and are connected with the same covariant eigenfunctions (in the sense of (7)). These eigenfunctions correspond to restrictions of eigenfunctions of the infinitesimal generator of

Koopman/Perron-Frobenius operators defined on \mathcal{H} . The proofs of these results are fully detailed in the next section.

The kernel isometry (5) (Theorem 3) and the Koopman spectral representation in Wonderland (Theorem 1) constitute fundamental results enabling us to build very simple ensemble-based trajectory reconstructions for new initial conditions without the requirement of resimulating the dynamical system. Amazingly, the Wonderland family of RKHS together with the Koopman isometry allows to define a system's trajectory as a constant-in-time linear combination of the time varying RKHS feature maps. Several of such reconstruction techniques, based on this fully justified superposition principle, will be shown in section 5. We will in particular demonstrate the potential of this framework with a simple and very cheap data assimilation technique in the context of very scarce data in space and time. In the next section we demonstrate the different properties related to Wonderland.

4.1 Proofs on the properties of the Koopman operator in Wonderland and of the Wonderland spectral theorem

As explained in the previous section the RKHS family, $(\mathcal{H}_t)_{t \geq 0}$, does not have a good topology to work with. We need first to start defining a “big” set with a better topology and which encompasses all the RKHS \mathcal{H}_t . On this big encompassing set, we shall then define a Koopman operator enabling us to study properly the Koopman operator in Wonderland.

4.1.1 Construction of the “big” encompassing set \mathcal{H}

The phase space Ω corresponds to the set generated by the value of the dynamical system at all time t . We note hence that Ω_t is a subset of Ω . Let us define a symmetric positive definite map $k : \Omega \times \Omega$ from the kernel $k_t : \Omega_t \times \Omega_t$ such that, for all $X_t = \Phi_t(X) \in \Omega, Y \in \Omega$

$$k(X_t, Y) = k_t(\Phi_t(X), \Phi_t(Y)), \quad (8)$$

and so as its restriction $k \upharpoonright_{\Omega_t \times \Omega_t} = k_t$. The extension k of kernel k_t , could be formulated for each points X_s of the trajectory X_t , and in particular for the initial time:

$$k(X_t, Y) = k_0(X, \Phi_t^{-1}(Y)). \quad (9)$$

As both expressions (8) and (9) should be equal, we obtain directly the isometric property (5) of kernels k_t . Kernel k inherits the regularity condition of k_t and is $C^2(\Omega \times \Omega)$ as well. By the Moore-Aronszajn theorem, there exists a unique RKHS called $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}}, \|\cdot\|_{\mathcal{H}})$ with kernel k . The RKHS \mathcal{H} can be connected to each RKHS of the family \mathcal{W} through extension and restriction operators denoted E_t and R_t respectively, and defined as follows. For all $t \geq 0$, let

$$E_t : \begin{array}{ccc} \mathcal{H}_t & \mapsto & \mathcal{H} \\ k_t(\cdot, X_t) & \mapsto & k(\cdot, X_t) \end{array} \quad (10)$$

and extend this definition by linearity on $\text{Vect}\{k_t(\cdot, X_t) : X_t \in \Omega_t\}$. Then, by density the function $E_t(f)$ is defined for all $f \in \mathcal{H}_t$. The restriction ,

$$\begin{aligned} R_t : \quad \mathcal{H} &\mapsto \mathcal{H}_t \\ k(\cdot, X_t) &\mapsto k(\cdot, X_t) \upharpoonright_{\Omega_t} = k_t(\cdot, X_t) \end{aligned} \quad (11)$$

is defined similarly for $g \in \mathcal{H}^0 = \text{Vect}\{k(\cdot, X) : X \in \Omega\}$ and extended by density in \mathcal{H} by the Moore-Aronszajn theorem [4]. The expression of $R_t(f)$ for $f \in \mathcal{H}$ is further specified in remark 17 (B). The extension map is built in such a way that each RKHS \mathcal{H}_t of the family is included in the “big” encompassing RKHS \mathcal{H} , as any functions of \mathcal{H}_t can be evaluated through kernel k , and expressed as a linear combination of the feature maps $k(\cdot, X_t)$. Let us stress again that for the type of flows targeted here, \mathcal{H} is in practice totally inaccessible, as opposed to \mathcal{H}_t .

In B, we list several useful properties of the restriction and extension operators. Namely, E_t and R_t are both isometries (Prop.12); they form an adjoint pair (Prop.13); the restriction is continuous in $L^2_{\mathbb{C}}(\Omega_t, \nu)$ (Prop.14). We define now the Koopman operator on the encompassing RKHS \mathcal{H} .

4.1.2 Koopman operators on \mathcal{H}

For all $t \geq 0$, we consider the Koopman operator on \mathcal{H} defined by $U_t(f) := f \circ \Phi_t$, for all $f \in \mathcal{H}$. Since \mathcal{H} is dense in $(L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$ by proposition 11 (A.3), the operator U_t can be continuously extended on $L^2_{\mathbb{C}}(\Omega, \nu)$ and to avoid notations inflation we keep noting U_t this extension. We first study $U_t : L^2_{\mathbb{C}}(\Omega, \nu) \mapsto L^2_{\mathbb{C}}(\Omega, \nu)$ with the $L^2_{\mathbb{C}}(\Omega, \nu)$ topology. The family $(U_t)_{t \geq 0}$ is a strongly continuous semi-group on $(L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$ since $t \mapsto \Phi_t(\cdot)$ is continuous on \mathbb{R}^+ . As the feature maps are functions of \mathcal{H} , it can be noticed that for all $X \in \Omega$

$$U_t[k(\cdot, X)] = k(\Phi_t(\cdot), X). \quad (12)$$

This corresponds to a natural expression of the Koopman operator for any functions $g \in \mathcal{H}^0$, extended then by density to \mathcal{H} . Yet another useful equivalent expression of the Koopman operator is available for the feature maps. As a matter of fact, since k is a Hermitian function, we have that for all X and $Y \in \Omega$, $k(\cdot, X)(Y) = \overline{k(\cdot, Y)}(X)$ and consequently by remark 11 (A.1), and the definition of extension k of k_t we have

$$U_t[k(\cdot, X)](Y) = k(\Phi_t(Y), X) = k_t(\Phi_t(Y), \Phi_t(X)), \quad (13)$$

and

$$k_t(\Phi_t(Y), \Phi_t(X)) = \overline{k_t(\Phi_t(X), \Phi_t(Y))} = \overline{k(\Phi_t(X), Y)}, \quad (14)$$

which leads to

$$U_t \overline{k(\cdot, Y)}(X) = k(Y, \Phi_t(X)). \quad (15)$$

and hence

$$U_t[k(\cdot, X)](Y) = k(Y, \Phi_t(X)). \quad (16)$$

As the previous equality is true for all $Y \in \Omega$, this implies that for all $X \in \Omega$

$$U_t[k(\cdot, X)] = k(\cdot, \Phi_t(X)). \quad (17)$$

This dual formulation of the kernel expression of the Koopman operator is intrinsically linked to the definition of the kernels k_t and to the fact that their associated feature maps are evaluated at the same time. This expression will be also recovered later on from the expression of the infinitesimal generator of the Koopman operator U_t . This dual expression will be of central interest in the following as it enables us to formulate the time evolution of the feature maps in terms of the Koopman operator U_t at any time $t \geq 0$.

Remark 1 (Transport of the kernel k) For all $X \in \Omega$ and $t \geq 0$, by definition of the Koopman operators on the feature maps and (17), $U_t[k(\cdot, X)]$ has two expressions and we obtain

$$U_t[k(\cdot, X)] = k(\Phi_t(\cdot), X) = k(\cdot, \Phi_t(X)).$$

The next remark provides a useful commutation property between U_t and the kernel integral operator \mathcal{L}_k defined in equation (1) or of its unique symmetric square-root $\mathcal{L}_k^{1/2}$ defined from the square-root of the kernel eigenvalues (see A.3 for a precise definition).

Remark 2 (Commutation between $\mathcal{L}_k^{1/2}$ (or \mathcal{L}_k) and U_t) For all $X \in \Omega$, we have

$$\mathcal{L}_k^{1/2} \circ U_t[k(\cdot, X)] = U_t \circ \mathcal{L}_k^{1/2}[k(\cdot, X)].$$

PROOF By (12), we have $\mathcal{L}_k^{1/2} \circ U_t[k(\cdot, X)] = \mathcal{L}_k^{1/2}[k(\Phi_t(\cdot), X)]$. The map $\mathcal{L}_k^{1/2} \circ U_t[k(\cdot, X)]$ can also be seen as the map $[\mathcal{L}_k^{1/2} k(\cdot, X)](\Phi_t(\cdot))$, and we obtain for all $Y \in \Omega$

$$(\mathcal{L}_k^{1/2} \circ U_t[k(\cdot, X)])(Y) = [\mathcal{L}_k^{1/2} k(\cdot, X)](\Phi_t(Y)) = U_t[\mathcal{L}_k^{1/2} k(\cdot, X)](Y),$$

by definition of the Koopman operator since $\mathcal{L}_k^{1/2} k(\cdot, X) \in \mathcal{H}$. \square

This commutation property that ensues directly from the compositional nature of the Koopman operator allows us to write immediately the equality

$$\mathcal{L}_k \circ U_t[k(\cdot, X)] = U_t \circ \mathcal{L}_k[k(\cdot, X)]. \quad (18)$$

By linearity these properties extend to all functions of \mathcal{H} . Let us note that the function $\mathcal{L}_k^{-1/2} k(\cdot, X)$, for all $X \in \Omega$, does not necessarily belong to \mathcal{H} , therefore this proof cannot be applied to $\mathcal{L}_k^{-1/2}$.

The next proposition shows the Koopman operator defined on RKHS \mathcal{H} is unitary in $L^2_{\mathbb{C}}(\Omega, \nu)$, which is a classical property of the Koopman operator in L^2 for measure preserving invertible system.

Proposition 1 (Unitarity of the Koopman operator in $L^2_{\mathbb{C}}(\Omega, \nu)$) *The map $U_t : (L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)}) \mapsto (L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$ is unitary for all $t \geq 0$.*

PROOF Since the dynamical system $(\Phi_t)_{t \geq 0}$ is measure preserving, (i.e. $\forall A \in \Omega, \nu(A) = \nu(\Phi_t^{-1}(A))$), with $\Phi_t^{-1}(A)$ denoting the pre-image of Φ_t and left invariant (i.e. $\Phi_t(\Omega) = \Omega$, with $\text{Ran}(\Phi_t) = \Omega$), the Koopman operator U_t is an isometry: for all f and $g \in \mathcal{H}$

$$\langle U_t(f), U_t(g) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle f, g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}.$$

By density, this property is extended for f and $g \in L^2_{\mathbb{C}}(\Omega, \nu)$. Besides, it can be noticed that $U_t(\mathcal{H}^0)$, with $\mathcal{H}^0 = \text{Vect}\{k(\cdot, X) : X \in \Omega\}$, is a subset of the range of $U_t : L^2_{\mathbb{C}}(\Omega, \nu) \mapsto L^2_{\mathbb{C}}(\Omega, \nu)$, $U_t(\mathcal{H}^0) = \text{Vect}\{k(\cdot, \Phi_t(X)) : X \in \Omega\}$ and we have the following inclusions

$$U_t(\mathcal{H}^0) \subset \text{Ran}(U_t) \subset L^2_{\mathbb{C}}(\Omega, \nu).$$

We show now that $U_t(\mathcal{H}^0)$ is dense in $(L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$. Let $f \in L^2_{\mathbb{C}}(\Omega, \nu)$ be such that $\langle f, k(\cdot, \Phi_t(X)) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = 0$ for all $X \in \Omega$, or in a more compact form $(\mathcal{L}_k f)[\Phi_t(X)] = 0$. We infer that

$$0 = \langle \mathcal{L}_k^{1/2} \circ f \circ \Phi_t, \mathcal{L}_k^{1/2} \circ f \circ \Phi_t \rangle_{\mathcal{H}} = \langle f \circ \Phi_t, f \circ \Phi_t \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \|f\|_{L^2_{\mathbb{C}}(\Omega, \nu)}^2,$$

consequently $f = 0$. By a consequence of Hahn-Banach theorem this result shows that the range of U_t is dense in $L^2_{\mathbb{C}}(\Omega, \nu)$ and Proposition 1 is proved. \square

Proposition 1, shows that the Koopman operator U_t is invertible and that its inverse in $L^2_{\mathbb{C}}(\Omega, \nu)$ is U_t^* . Denoting by P_t the operator defined by $P_t := U_t^*$, for all f and $g \in L^2_{\mathbb{C}}(\Omega, \nu)$ we have $\langle U_t(f), g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle f, P_t(g) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}$. The family $(P_t)_t$ is a strongly continuous semi-group on $(L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$. This operator is referred to as the Perron-Frobenius operator. For all $t \geq 0$, the Perron-Frobenius operator P_t verifies for all $X \in \Omega$

$$P_t[k(\cdot, \Phi_t(X))] = k(\cdot, X). \quad (19)$$

As previously stated the Koopman operator U_t is an isometry in $L^2_{\mathbb{C}}(\Omega, \nu)$. The next result asserts that U_t is also an isometry in \mathcal{H} .

Proposition 2 (Isometric relation of the kernel) *For all X and $Y \in \Omega$, we have $k(\Phi_t(X), \Phi_t(Y)) = k(X, Y)$.*

PROOF By remark 2, we see the function $\mathcal{L}_k[U_t k(\cdot, \Phi_t(Y))]$ of \mathcal{H} as $[\mathcal{L}_k k(\cdot, \Phi_t(Y))](\Phi_t(\cdot))$. By (17) and (58), we have for all $X \in \Omega$

$$\begin{aligned} [\mathcal{L}_k k(\cdot, \Phi_t(Y))](\Phi_t(X)) &= \langle \mathcal{L}_k k(\cdot, \Phi_t(Y)), k(\cdot, \Phi_t(X)) \rangle_{\mathcal{H}} \\ &= \langle \mathcal{L}_k^{1/2} \circ U_t[k(\cdot, Y)], \mathcal{L}_k^{-1/2} \circ U_t[k(\cdot, X)] \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}. \end{aligned}$$

As $\mathcal{L}_k^{1/2}$ is self-adjoint on $L_{\mathbb{C}}^2(\Omega, \nu)$, upon applying proposition 1 we obtain

$$\begin{aligned} [\mathcal{L}_k k(\cdot, \Phi_t(Y))](\Phi_t(X)) &= \langle U_t[k(\cdot, Y)], U_t[k(\cdot, X)] \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \\ &= \langle k(\cdot, Y), k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}. \end{aligned}$$

In the same way, we have also

$$\begin{aligned} \langle k(\cdot, Y), k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} &= \langle \mathcal{L}_k^{1/2} k(\cdot, Y), \mathcal{L}_k^{-1/2} k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \\ &= \langle \mathcal{L}_k k(\cdot, Y), k(\cdot, X) \rangle_{\mathcal{H}}, \end{aligned}$$

and we obtain

$$[\mathcal{L}_k k(\cdot, \Phi_t(Y))](\Phi_t(X)) = [\mathcal{L}_k k(\cdot, Y)](X).$$

Hence $\mathcal{L}_k[U_t k(\cdot, \Phi_t(Y))](X) = \mathcal{L}_k k(\cdot, Y)(X)$ for all $X \in \Omega$ and as \mathcal{L}_k is injective, we have finally $k(\Phi_t(X), \Phi_t(Y)) = k(X, Y)$ for all X and $Y \in \Omega$. \square

Remark 1 on the application of the operator U_t to $k(\cdot, X)$, with $X \in \Omega$, enables us to apply the flow, Φ_t , on the first or second variable of $k(\cdot, \cdot)$. Consequently, for all $f \in \mathcal{H}$ and $X \in \Omega$, we have on the one hand $\langle f, k(\cdot, \Phi_t(X)) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle f, U_t k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}$ and on the other hand, considering now $U_t k(\cdot, X)$ as $k(\Phi_t(\cdot), X)$ and by definition of P_t we obtain $\langle f, U_t k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle P_t(f), k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}$. Hence to sum up, we get that for all $f \in \mathcal{H}$, $X \in \Omega$ and $t \geq 0$

$$\langle f, k(\cdot, \Phi_t(X)) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle P_t(f), k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}. \quad (20)$$

Interpreting the Perron-Frobenius in the usual sense as a transfer operator of density, this equation provides an expression of the transported density through the evaluation with the feature maps $k(\cdot, \Phi_t(X))$. This expression is in practice much easier to compute than the usual one, thanks to the kernel isometry (Proposition 2).

In order to further exhibit several analytical results on the Koopman operator $(U_t)_{t \geq 0}$ in $L_{\mathbb{C}}^2(\Omega, \nu)$, we introduce in the following its infinitesimal generator.

4.1.3 Koopman infinitesimal generator

We will note as $(A_U, \mathcal{D}(A_U))$ the infinitesimal generator of the strongly continuous semigroup $(U_t)_{t \geq 0}$ on $L_{\mathbb{C}}^2(\Omega, \nu)$ and its domain. As the Koopman and Perron-Frobenius operators are adjoint in $L_{\mathbb{C}}^2(\Omega, \nu)$ their infinitesimal generators are also adjoint of each other with possibly their own domain. The following lemma characterizes first the Perron-Frobenius infinitesimal generator A_P and its domain $\mathcal{D}(A_P)$.

Lemma 1 (Perron-Frobenius infinitesimal generator) *The Perron-Frobenius infinitesimal generator is the unbounded operator, $(A_P, \mathcal{D}(A_P))$ defined by*

$$\begin{aligned} \mathcal{D}(A_P) &= \{f \in L_{\mathbb{C}}^2(\Omega, \nu) : x \mapsto \partial_{\mathcal{M}(x)} f(x) \in L_{\mathbb{C}}^2(\Omega, \nu)\} \\ &\text{and } A_P f := -\partial_{\mathcal{M}(\cdot)} f(\cdot) \text{ for } f \in \mathcal{D}(A_P), \end{aligned}$$

where \mathcal{M} denotes the differential tendency of the system dynamics (3) and $\partial_u f$ stands for the directional derivative of f along $u \in \Omega$.

PROOF See C.

For all $t \geq 0$, P_t is the adjoint of U_t in $L^2_{\mathbb{C}}(\Omega, \nu)$. The Koopman infinitesimal generator in $L^2_{\mathbb{C}}(\Omega, \nu)$ is consequently given by

$$\mathcal{D}(A_U) = \mathcal{D}(A_P^*) \quad \text{and} \quad A_U = A_P^*. \quad (21)$$

The Koopman and Frobenius-Perron operators are unitary in $L^2_{\mathbb{C}}(\Omega, \nu)$, as $\mathcal{D}(A_U)$ is dense in $L^2_{\mathbb{C}}(\Omega, \nu)$, by Stone's theorem, their infinitesimal generators are therefore skew symmetric for $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}$ and we have

$$\mathcal{D}(A_P^*) = \mathcal{D}(A_P) \quad \text{and} \quad A_P^* = -A_P, \quad (22)$$

$$\mathcal{D}(A_U^*) = \mathcal{D}(A_U) \quad \text{and} \quad A_U^* = -A_U. \quad (23)$$

The adjoint has to be understood in the topology of $L^2_{\mathbb{C}}(\Omega, \nu)$. The two next propositions – well known in $L^2_{\mathbb{C}}(\Omega, \nu)$ for invertible measure preserving systems –, summarize (21), (22) and (23).

Proposition 3 (Koopman infinitesimal generator) *The Koopman infinitesimal generator of $(U_t)_t$ in $L^2_{\mathbb{C}}(\Omega, \nu)$ is the unbounded operator, $(A_U, \mathcal{D}(A_U))$, defined by*

$$\begin{aligned} \mathcal{D}(A_U) &= \{f \in L^2_{\mathbb{C}}(\Omega, \nu) : x \mapsto \partial_{\mathcal{M}(x)} f(x) \in L^2_{\mathbb{C}}(\Omega, \nu)\} \\ &\quad \text{and } A_U f = \partial_{\mathcal{M}(\cdot)} f(\cdot) \quad \text{for } f \in \mathcal{D}(A_U). \end{aligned}$$

Proposition 4 (Skew symmetry of the generators) *The Koopman infinitesimal generator A_U is skew-symmetric in $L^2_{\mathbb{C}}(\Omega, \nu)$.*

We set now a continuity property of the Koopman infinitesimal generator on a subspace of $\mathcal{D}(A_U)$. Let $f \in \mathcal{H}$, by theorem 6 (A.2), $x \mapsto \partial_{\mathcal{M}(x)} f(x) \in L^2_{\mathbb{C}}(\Omega, \nu)$, and through Cauchy-Schwarz inequality we get that $f \in \mathcal{D}(A_U)$ since the real valued function \mathcal{M} belongs to $L^2_{\mathbb{C}}(\Omega, \nu)$ and consequently $\mathcal{H} \subset \mathcal{D}(A_U)$.

Theorem 2 (Continuity of the Koopman generator on \mathcal{H}) *The restriction of the Koopman infinitesimal generator*

$$A_U : (\mathcal{H}, \|\cdot\|_{\mathcal{H}}) \mapsto (L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$$

is continuous.

PROOF For all $f \in \mathcal{H}$ by the Cauchy-Schwarz inequality and theorem 7 (A.2) we have

$$\|A_U f\|_{L^2_{\mathbb{C}}(\Omega, \nu)} = \|\partial_{\mathcal{M}(\cdot)} f(\cdot)\|_{L^2_{\mathbb{C}}(\Omega, \nu)} \leq C \|f\|_{\mathcal{H}}.$$

□

Through the dual expression of the Koopman operator (remark 1) the infinitesimal generator A_U provides a dynamical system specifying the time evolution of the feature maps as, for all $X \in \Omega$, the feature map $k(\cdot, \Phi_t(X))$ verifies

$$\partial_t U_t k(\cdot, X) = \partial_t k(\cdot, \Phi_t(X)) = A_U k(\cdot, \Phi_t(X)). \quad (24)$$

We have the following useful differentiation formulae.

Proposition 5 (Differentiation formulae) *For all $f \in \mathcal{H}$ (with $f = E_t g$, $g \in \mathcal{H}_t$) and $X_t \in \Omega_t$ we have*

$$\langle \mathcal{L}_k \circ \partial_t f, k(\cdot, X_t) \rangle_{\mathcal{H}} = -\langle f, \mathcal{L}_k \circ \partial_t k(\cdot, X_t) \rangle_{\mathcal{H}}.$$

PROOF As $\mathcal{L}_k \circ A_U f$ belongs to \mathcal{H} , we have $\mathcal{L}_k \circ A_U f(X_t) = \mathcal{L}_k \circ \partial_t f(X_t)$ and

$$\langle \mathcal{L}_k \circ \partial_t f, k(\cdot, X_t) \rangle_{\mathcal{H}} = \langle \mathcal{L}_k \circ A_U f, k(\cdot, X_t) \rangle_{\mathcal{H}} = \langle \mathcal{L}_k^{1/2} \circ A_U f, \mathcal{L}_k^{-1/2} k(\cdot, X_t) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}.$$

As $\mathcal{L}_k^{1/2}$ is self-adjoint in $L^2_{\mathbb{C}}(\Omega, \nu)$ and A_U is skew-symmetric in $L^2_{\mathbb{C}}(\Omega, \nu)$, we obtain

$$\begin{aligned} \langle \mathcal{L}_k \circ \partial_t f, k(\cdot, X_t) \rangle_{\mathcal{H}} &= \langle A_U f, k(\cdot, X_t) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \\ &= -\langle f, A_U k(\cdot, X_t) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \\ &= -\langle f, \partial_t k(\cdot, X_t) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}. \end{aligned}$$

As $\langle f, \partial_t k(\cdot, X_t) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle f, \mathcal{L}_k \circ \partial_t k(\cdot, X_t) \rangle_{\mathcal{H}}$, this complete the proof. \square

As already outlined, the whole phase space, Ω , and the global embedding RKHS, \mathcal{H} , defined on it are both completely inaccessible for high dimensional state spaces. Instead of seeking to reconstruct this global RKHS, we will work in the following with time-varying ‘‘local’’ RKHS spaces built from a small (w.r.t. the phase space dimension) ensemble of initial conditions. The set of these time-varying spaces forms the Wonderland family. To express the time evolution of the features maps associated to the Wonderland RKHS family we now define an appropriate expression of the Koopman operator on this family.

4.1.4 Derivation of Koopman operator expression in \mathcal{W}

To fully specify the Koopman operators in Wonderland, we rely on the family of extension, restriction mappings $(E_t)_{t \geq 0}$ and $(R_t)_{t \geq 0}$ (10, 11) relating the ‘‘big’’ encompassing RKHS \mathcal{H} to the family of time-evolving RKHS \mathcal{H}_t and use the Koopman operators $U_t : L^2_{\mathbb{C}}(\Omega, \nu) \mapsto L^2_{\mathbb{C}}(\Omega, \nu)$ for $t \geq 0$.

From now on, we note $X_t := \Phi_t(X_0)$ for all $X_0 \in \Omega_0$ with X_t belonging to Ω_t . We define the Koopman operator in Wonderland by $\mathcal{U}_t := R_t \circ U_t \circ E_0$ for all $t \geq 0$. For all $Y_0 \in \Omega_0$ and $X_t \in \Omega_t$, we have

$$\mathcal{U}_t [k_0(\cdot, Y_0)](X_t) = R_t \circ U_t [k(\cdot, Y_0)](X_t) = R_t k(\cdot, Y_t)(X_t) = k_t(X_t, Y_t). \quad (25)$$

The Koopman operator in Wonderland, $\mathcal{U}_t : \mathcal{H}_0 \mapsto \mathcal{H}_t$, transports the kernel feature maps on the RKHS family by composition with the system's dynamics. It inherits some of the nice properties of the Koopman operator defined on the encompassing global RKHS. As shown by the following theorem, proposition 2 remains valid for the family of kernels $(k_t)_{t \geq 0}$ and \mathcal{U}_t is still unitary in the sense of the following theorem.

Theorem 3 (Koopman RKHS isometry) *The Koopman operator on the RKHS family defines an isometry from \mathcal{H}_0 to \mathcal{H}_t : for all X_0 and $Y_0 \in \Omega_0$*

$$\begin{aligned} \langle \mathcal{U}_t k_0(\cdot, Y_0), \mathcal{U}_t k_0(\cdot, X_0) \rangle_{\mathcal{H}_t} &= \langle k_0(\cdot, Y_0), k_0(\cdot, X_0) \rangle_{\mathcal{H}_0}, \\ k_t(\Phi_t(X_0), \Phi_t(Y_0)) &= k_0(X_0, Y_0). \end{aligned}$$

The range of $\mathcal{U}_t : \mathcal{H}_0 \mapsto \mathcal{H}_t$ is dense in \mathcal{H}_t .

PROOF Upon applying equations (62 of Prop. 12, B), and (17) with the definition of E_0 , we have

$$\begin{aligned} \langle \mathcal{U}_t k_0(\cdot, Y_0), \mathcal{U}_t k_0(\cdot, X_0) \rangle_{\mathcal{H}_t} &= \langle U_t k(\cdot, Y_0), U_t k(\cdot, X_0) \rangle_{\mathcal{H}} \\ &= \langle k(\cdot, Y_t), k(\cdot, X_t) \rangle_{\mathcal{H}} = k_t(X_t, Y_t). \end{aligned}$$

With proposition 2 we obtain

$$\langle \mathcal{U}_t k_0(\cdot, Y_0), \mathcal{U}_t k_0(\cdot, X_0) \rangle_{\mathcal{H}_t} = k(X_0, Y_0) = \langle k(\cdot, Y_0), k(\cdot, X_0) \rangle_{\mathcal{H}}.$$

By definition of R_0 and again (62), we conclude that

$$\langle \mathcal{U}_t k_0(\cdot, Y_0), \mathcal{U}_t k_0(\cdot, X_0) \rangle_{\mathcal{H}_t} = \langle k_0(\cdot, Y_0), k_0(\cdot, X_0) \rangle_{\mathcal{H}_0}.$$

Besides, the image of $\text{Vect}\{k_0(\cdot, Y_0) : Y_0 \in \Omega_0\}$ under \mathcal{U}_t is the set $\text{Vect}\{k_t(\cdot, Y_t) : Y_t \in \Omega_t\}$ which is dense in $(\mathcal{H}_t, \|\cdot\|_{\mathcal{H}_t})$. The range of $\mathcal{U}_t : \mathcal{H}_0 \mapsto \mathcal{H}_t$ is thus dense in \mathcal{H}_t and the operator \mathcal{U}_t is, in that sense, unitary. \square

Let us now determine the adjoint of the Koopman operator in Wonderland. Let $\mathcal{P}_t := R_0 \circ P_t \circ E_t$ for all $t \geq 0$ and let $X_t \in \Omega_t$. With the same arguments as for \mathcal{U}_t and (19), we have for all $Y_t \in \Omega_t$ and $X_0 \in \Omega_0$

$$\mathcal{P}_t [k_t(\cdot, Y_t)](X_0) = k_0(X_0, Y_0). \quad (26)$$

The mapping $\mathcal{P}_t : \mathcal{H}_t \mapsto \mathcal{H}_0$ with $t \geq 0$ constitute the Perron-Frobenius family of operators in Wonderland. The mapping $\mathcal{P}_t : \mathcal{H}_t \mapsto \mathcal{H}_0$ is unitary for the RKHS family topology (isometry from \mathcal{H}_t to \mathcal{H}_0 and the range of \mathcal{P}_t is dense in \mathcal{H}_0). The next proposition justifies that \mathcal{U}_t and \mathcal{P}_t have inverse action on the feature maps.

Proposition 6 (Koopman Perron-Frobenius duality) *For all $X_0 \in \Omega_0$ and $Y_t \in \Omega_t$, we have*

$$\langle \mathcal{U}_t k_0(\cdot, X_0), k_t(\cdot, Y_t) \rangle_{\mathcal{H}_t} = \langle k_0(\cdot, X_0), \mathcal{P}_t [k_t(\cdot, Y_t)] \rangle_{\mathcal{H}_0}.$$

PROOF By the equations (25) and (26), we have

$$\langle \mathcal{U}_t k_0(\cdot, X_0), k_t(\cdot, Y_t) \rangle_{\mathcal{H}_t} = k_t(Y_t, X_t) \text{ and } \langle k_0(\cdot, X_0), \mathcal{P}_t[k_t(\cdot, Y_t)] \rangle_{\mathcal{H}_0} = k_0(Y_0, X_0),$$

and applying theorem 3 completes the proof. \square

In order to derive the Koopman and Perron-Frobenius operators' spectral representation in Wonderland, we exhibit now two propagation operators that will allow us to express the evolution of the Wonderland's feature maps.

4.1.5 Wonderland spectral representation

We specify hereafter a family of operator $A_{U,t}$, related to the Koopman infinitesimal generator A_U , and that give rise to an evolution equation on Ω_t akin to (24). They will play, in that sense, the role of infinitesimal generators on the RKHS family.

For all $t \geq 0$, let $A_{U,t}$ be defined by $A_{U,t} := R_t \circ A_U \circ E_t$ with $E_t : \mathcal{H}_t \mapsto \mathcal{H}$ and $R_t : L_{\mathbb{C}}^2(\Omega, \nu) \mapsto L_{\mathbb{C}}^2(\Omega_t, \nu)$ (64 B).

Proposition 7 (Continuity of $A_{U,t}$) *The mapping $A_{U,t} : (\mathcal{H}_t, \|\cdot\|_{\mathcal{H}_t}) \mapsto (L_{\mathbb{C}}^2(\Omega_t, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega_t, \nu)})$ is well-defined and continuous.*

PROOF Upon applying (61 of Prop. 12, B), the continuity of A_U (Theorem 2) and of R_t (Proposition 14, B) we obtain the proof of the proposition. \square

In a very similar way as the infinitesimal generator on \mathcal{H} , the operator $A_{U,t}$ can be understood as an evolution equation of the feature maps defined on Ω_t and associated to \mathcal{H}_t . For $X_t \in \Omega_t$, by proposition 3 it can be noticed that

$$A_{U,t} k_t(\cdot, X_t) = R_t \circ A_U [k(\cdot, X_t)] = R_t [\partial_{\mathcal{M}(\cdot)} k(\cdot, X_t)]$$

and in particular on Ω_t

$$A_{U,t} k_t(\cdot, X_t) = \partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t). \quad (27)$$

Remark 3 Operator $A_{U,t}$ is alike the Koopman infinitesimal generator A_U . As a matter of fact by proposition 3 and theorem 2 we have for all $X \in \Omega$

$$A_U k(\cdot, X) = \partial_{\mathcal{M}(\cdot)} k(\cdot, X),$$

and for all $X_t \in \Omega_t$

$$A_{U,t} k_t(\cdot, X_t) = \partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t).$$

The domains of both operators are nevertheless different.

Remark 4 Operators $A_{U,t}$ can be understood as an evolution operator for the feature maps. As a matter of fact through (24) and the definition of $A_{U,t}$, we have, for all $X \in \Omega_{t_0}$

$$\partial_t k(\cdot, \Phi_t(X)) = A_{U,t} k_t(\cdot, \Phi_t(X)). \quad (28)$$

Through the above remark the operator $A_{U,t}$ inherits the properties of operator A_U defined on the global encompassing RKHS \mathcal{H} . As shown in the next proposition it remains in particular skew-symmetric.

Proposition 8 (Skew-symmetry of $A_{U,t}$) *The operator $A_{U,t}$ is skew-symmetric in $L^2_{\mathbb{C}}(\Omega_t, \nu)$: for all f and $g \in \mathcal{H}_t$*

$$\langle A_{U,t} f, g \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)} = -\langle f, A_{U,t} g \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)}.$$

For the need of this proof only, we introduce the following notation. For a function h defined on Ω_t , we note \widetilde{h} the extension on Ω defined by

$$\widetilde{h} := \begin{cases} h & \text{on } \Omega_t \\ 0 & \text{else.} \end{cases}$$

PROOF As $g = E_t(g)$ holds on Ω_t , we have

$$\langle A_{U,t} f, g \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)} = \langle A_U \circ E_t(f) \upharpoonright_{\Omega_t}, E_t(g) \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)} = \langle A_U \circ \widetilde{E_t(f)}, E_t(g) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}$$

By proposition 4 and since $E_t(f) \upharpoonright_{\Omega_t} = f$ on Ω_t we infer that

$$\begin{aligned} \langle A_{U,t} f, g \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)} &= -\langle E_t(f), A_U \circ \widetilde{E_t(g)} \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \\ &= -\langle f, A_U \circ E_t(g) \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)} \\ &= -\langle f, A_{U,t}(g) \rangle_{L^2_{\mathbb{C}}(\Omega_t, \nu)}. \end{aligned}$$

□

We are now ready to prove the Wonderland spectral representation theorem, which states that the bounded operator $A_{U,t} : \mathcal{H}_t \mapsto L^2_{\mathbb{C}}(\Omega_t, \nu)$ is diagonalizable for all $t \geq 0$.

4.1.6 Proof of Theorem 1 [Wonderland spectral representation]

The full proof, organized in two main steps, is thoroughly detailed in D. For this proof, we consider the infinitesimal generator $A_U : \mathcal{H} \mapsto L^2_{\mathbb{C}}(\Omega, \nu)$, which is connected to each $A_{U,t}$ for $t \geq 0$ through the restriction operator R_t . In the first step, the diagonalization of operator $A_U : \mathcal{H} \mapsto L^2_{\mathbb{C}}(\Omega, \nu)$ in $L^2_{\mathbb{C}}(\Omega, \nu)$ is first performed. To that end we introduce an intermediate (approximating) operator denoted \widetilde{A}_U , directly related to A_U and whose inverse is shown to be compact and self-adjoint. The second step of the proof consists in deducing the diagonalization of each $A_{U,t}$ from the diagonalization of A_U obtained at step 1. These two steps are thoroughly presented in D.

4.2 Tangent linear dynamics

A result of practical interest concerns the establishment of a rigorous ensemble expression of the tangent linear dynamics operator. Recall that we note $X_t = \Phi_t(X_0)$ for all $t \geq 0$. We define $\delta X(t, x) := \Phi_t[X_0(x) + \delta X_0(x)] - \Phi_t[X_0(x)]$ for all $t \geq 0$ and almost $x \in \Omega_x$, and where $\delta X_0(x)$ is a perturbation of the initial condition at point x . The function $\delta X(t, \cdot) \in L^2(\Omega_x, \mathbb{R}^d)$ is the perturbation of the flow at time t with respect to the initial condition X_0 . We have for almost any $x \in \Omega_x$

$$\begin{aligned} \Phi_t(X_0 + \delta X_0) - \Phi_t(X_0) &= \int_0^t \mathcal{M}[\Phi_s(X_0 + \delta X_0)] - \mathcal{M}[\Phi_s(X_0)] ds + o(\|\delta X_0\|) \\ &= \int_0^t d(\mathcal{M} \circ \Phi_s(X_0)) \delta X_0 ds + o(\|\delta X_0\|) \end{aligned}$$

and we obtain also that

$$\begin{aligned} \Phi_t(X_0 + \delta X_0) - \Phi_t(X_0) &= \int_0^t d\mathcal{M}[\Phi_s(X_0)] [d\Phi_s(X_0) \delta X_0] ds + o(\|\delta X_0\|) \\ &= \int_0^t \partial_X \mathcal{M}[\Phi_s(X_0)] \delta X(s, \cdot) ds + o(\|\delta X_0\|). \end{aligned}$$

The variation of the flow verifies $\delta X(t, \cdot) = \int_0^t \partial_X \mathcal{M}[X_s] \delta X(s, \cdot) ds$ almost everywhere on Ω_x . Recall that \mathcal{M} is assumed C^1 , in particular $\sup_{X \in \Omega} \partial_X \mathcal{M}(X) < \infty$ (since Ω is compact). The function $\delta X(t, \cdot)$ belongs to $L^2(\Omega_x, \mathbb{R}^d)$ and verifies for almost $x \in \Omega_x$ and all $t \geq 0$

$$\partial_t \delta X(t, x) = \partial_X \mathcal{M}[X_t] \delta X(t, x). \quad (29)$$

Each component $[\delta X(t, \cdot)]_i$ of $\delta X(t, \cdot)$ belongs to $L^2(\Omega_x, \mathbb{R})$. Let $[g_x^\varepsilon(\delta X_t)]_i : \Omega \mapsto \mathbb{C}$ be the representation of $[\delta X(t, \cdot)]_i$ in $L^2_{\mathbb{C}}(\Omega, \nu)$ defined for all $x \in \Omega_x$ by

$$[g_x^\varepsilon(\delta X_t)]_i := [\delta X_t(x)]_i.$$

For any $x \in \Omega_x$, the family of functions $g_x^\varepsilon(X_t)$ correspond to vector observables of the dynamical system and we may immediately write for all $t \geq 0$

$$\partial_t g_x^\varepsilon(\delta X_t) = \partial_X \mathcal{M}(X_t) g_x^\varepsilon(\delta X_t). \quad (30)$$

We are now ready to exhibit a kernel expression of the tangent linear operator $\partial_X \mathcal{M}(X_t)$. For all $1 \leq i \leq d$, the function $\mathcal{L}_k [g_x^\varepsilon]_i \in \mathcal{H}$ verifies for all $\delta X_t \in \Omega$

$$\langle \mathcal{L}_k \circ \partial_t \mathcal{L}_k [g_x^\varepsilon]_i, k(\cdot, \delta X_t) \rangle_{\mathcal{H}} = \mathcal{L}_k (\partial_t \mathcal{L}_k [g_x^\varepsilon]_i) (\delta X_t). \quad (31)$$

Upon applying the differentiation formulae of proposition 5 and the evolution equation (24), we have

$$\begin{aligned} \langle \mathcal{L}_k \circ \partial_t \mathcal{L}_k [g_x^\varepsilon]_i, k(\cdot, \delta X_t) \rangle_{\mathcal{H}} &= -\langle \mathcal{L}_k [g_x^\varepsilon]_i, \mathcal{L}_k \circ \partial_t k(\cdot, \delta X_t) \rangle_{\mathcal{H}} \\ &= -\langle \mathcal{L}_k [g_x^\varepsilon]_i, \mathcal{L}_k [A_U k(\cdot, \delta X_t)] \rangle_{\mathcal{H}} \\ &= -\langle \mathcal{L}_k^{1/2} [g_x^\varepsilon]_i, \mathcal{L}_k^{1/2} [A_U k(\cdot, \delta X_t)] \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}. \end{aligned}$$

The operator $\mathcal{L}_k^{1/2}$ being self-adjoint in $L_{\mathbb{C}}^2(\Omega, \nu)$ and A_U skew-symmetric for the inner product of $L_{\mathbb{C}}^2(\Omega, \nu)$, we have

$$\begin{aligned} \langle \mathcal{L}_k \circ \partial_t \mathcal{L}_k [g_x^\varepsilon]_i, k(\cdot, \delta X_t) \rangle_{\mathcal{H}} &= \langle A_U \mathcal{L}_k [g_x^\varepsilon]_i, k(\cdot, \delta X_t) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \\ &= \langle \mathcal{L}_k^{1/2} A_U \mathcal{L}_k [g_x^\varepsilon]_i, \mathcal{L}_k^{-1/2} k(\cdot, \delta X_t) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \\ &= \langle \mathcal{L}_k A_U \mathcal{L}_k [g_x^\varepsilon]_i, k(\cdot, \delta X_t) \rangle_{\mathcal{H}} \\ &= \mathcal{L}_k A_U \mathcal{L}_k [g_x^\varepsilon]_i(\delta X_t). \end{aligned}$$

Combining the right-hand side of the above expression with (31), we have

$$\mathcal{L}_k(\partial_t \mathcal{L}_k [g_x^\varepsilon]_i)(\delta X_t) = \mathcal{L}_k(A_U \mathcal{L}_k [g_x^\varepsilon]_i)(\delta X_t). \quad (32)$$

As \mathcal{L}_k is injective, the kernel of the tangent linear operator in the RKHS \mathcal{H} reads:

$$(\partial_t \mathcal{L}_k [g_x^\varepsilon]_i)(\delta X_t) = A_U \mathcal{L}_k [g_x^\varepsilon]_i(\delta X_t).$$

As Ω is bounded, the function $t \mapsto \delta X(t, \cdot)$ belongs to $L^\infty([0, T])$ and therefore $\delta X_t \mapsto g_x^\varepsilon(\delta X_t)$ belongs to $L^\infty([0, T], L^2(\Omega, \nu))$. By (30), we have also that $\delta X_t \mapsto \partial_t g_x^\varepsilon(\delta X_t)$ belongs to $L^\infty([0, T], L^2(\Omega, \nu))$ since $\sup_{Y \in \Omega} \partial_X \mathcal{M}(Y) < \infty$. We have therefore on Ω

$$\partial_t \mathcal{L}_k [g_x^\varepsilon]_i(\delta X_t) = \int_{\Omega} k(\delta X_t, z) \partial_t [g_x^\varepsilon]_i(z) \nu(dz) = \mathcal{L}_k(\partial_t [g_x^\varepsilon]_i)(\delta X_t),$$

and thus,

$$\partial_t [g_x^\varepsilon]_i(\delta X_t) = \mathcal{L}_k^{-1} A_U \mathcal{L}_k [g_x^\varepsilon]_i(\delta X_t).$$

By the commutation property of remark 2, together with (30) and as $\mathcal{L}_k^{1/2} [g_x^\varepsilon]_i \in \mathcal{H}$, the following equalities hold for all $\delta X_t \in \Omega$

$$\begin{aligned} \partial_t [g_x^\varepsilon]_i(\delta X_t) &= \mathcal{L}_k^{-1/2} A_U \mathcal{L}_k^{1/2} [g_x^\varepsilon]_i(\delta X_t), \\ \partial_X \mathcal{M}(X_t) g_x^\varepsilon(\delta X_t) &= \mathcal{L}_k^{-1/2} A_U \mathcal{L}_k^{1/2} g_x^\varepsilon(\delta X_t). \end{aligned} \quad (33)$$

Let's specify now the kernel expression of the tangent linear dynamics in Ω_s for all $s \geq 0$. The function $\mathcal{L}_k^{1/2} [g_x^\varepsilon]_i|_{\Omega_s}$ belongs to \mathcal{H}_s . We have as well $\mathcal{L}_k^{1/2}(\partial_t g_x^\varepsilon) = \mathcal{L}_{k_s}^{1/2}(\partial_t g_x^\varepsilon)$ on Ω_s . But, for all Y_s and $X'_s \in \Omega_s$ we have that $A_U k(\cdot, X'_s)(Y_s) = A_{U,s} k_s(\cdot, X'_s)(Y_s)$. In particular, we get $A_U \mathcal{L}_k^{1/2} [g_x^\varepsilon]_i(Y_s) = A_{U,s} \mathcal{L}_{k_s}^{1/2} [g_x^\varepsilon]_i(Y_s)$. We obtain hence the kernel expression for all $Y_s \in \Omega_s$

$$\partial_X \mathcal{M}(X_s) g_x^\varepsilon(Y_s) = \mathcal{L}_{k_s}^{-1/2} A_{U,s} \mathcal{L}_{k_s}^{1/2} g_x^\varepsilon(Y_s). \quad (34)$$

Note that the domains of the kernel expressions of the tangent linear (33) and (34) are different. The right-hand side of (34) provides a convenient kernel expression of the tangent linear operator, enabling us to evaluate the tangent linear dynamics from an ensemble of feature maps. The adjoint of the tangent linear dynamics is straightforwardly given by:

$$\partial_X \mathcal{M}^*(X_s) g_x^\varepsilon(Y_s) = -\mathcal{L}_{k_s}^{-1/2} A_{U,s} \mathcal{L}_{k_s}^{1/2} g_x^\varepsilon(Y_s). \quad (35)$$

Remark 5 (Projection observables) The point observable functions g_x^ε used above can be extended to other functions defined from basis $(\Psi_j)_{j \geq 0}$ of $L^2(\Omega_x, \mathbb{R}^d)$, with

$$g_x^\Psi X_t := \sum_j \langle X_t, \Psi_j \rangle_{L^2(\Omega_x)} \Psi_j(x),$$

$$g_x^\Psi : \Omega \mapsto \sum_j \langle \cdot, \Psi_j \rangle_{L^2(\Omega_x)} \Psi_j(x) \in \mathbb{C}.$$

Remark 6 (Dependence on X_t) It should be noted that in (34) and (35) the right-hand side does not depend on X_s whereas the tangent linear operator or its adjoint on the left-hand side does. Function X_s indicates around which function of Ω_s the nonlinear system is linearized. The infinitesimal generator of the Koopman operator, representing the dynamics' linear tangent operator on the RKHS family, depends necessarily also on this function. This dependence is here implicit and induced by the considered sampled functions used to define the RKHS \mathcal{H}_t . If the set of members $(X_t^{(i)})_{i \geq 1}$ are centered around a particular function Y_s , the infinitesimal generators of the Koopman operator can be interpreted as a representation of the tangent linear operator around function Y_s . In ensemble methods, Y_s , is in general taken as the ensemble mean, and $(X_s^{(i)})_{i \geq 1}$ is an ensemble of of time-dependent perturbations around this mean.

Remark 7 (More regularity on δX_t) For all $1 \leq i \leq d$, if we suppose that the function $[g_x^\varepsilon]_i$ belongs to $\mathcal{H} \subset L^2_{\mathbb{C}}(\Omega, \nu)$, the proof can be simplified and (34) is replaced by

$$\partial_X \mathcal{M}(X_s) g_x^\varepsilon(\delta X_s) = A_{U,s} g_x^\varepsilon(\delta X_s). \quad (36)$$

It can be pointed out that the expression above corresponds to the approximation of the tangent linear dynamics used in ensemble method if we work in a finite dimensional space such that $\Omega \subset \mathbb{R}^n$ and assume that $k_t(\cdot, X_t)$ is defined as $(p-1)^{-1/2} \langle (X_t - \bar{X}_t), \cdot - \bar{X}_t \rangle_{\mathbb{R}^n}$, with \bar{X}_t the empirical ensemble mean. With that definition, we have $A_{U,t} k_t(\cdot, X^{(i)}) = \sum_j \nabla_{X_j} k_t(\cdot, X_t^{(i)}) \mathcal{M}(\cdot)_j = \langle (\delta_{X_t^{(i)}} - \frac{1}{p} \sum_\ell \delta_{X_t^{(\ell)}}) \cdot \mathcal{M}(\cdot), \cdot - \bar{X}_t \rangle_{\mathbb{R}^n}$ that reads $(p-1)^{-1/2} \langle \mathcal{M}(X_t^{(i)}) - \frac{1}{p} \sum_j \mathcal{M}(X_t^{(j)}), \cdot - \bar{X}_t \rangle_{\mathbb{R}^n}$, for $i = 1, \dots, p$ and for which, when associated to the \mathbb{R}^n Euclidean inner product on a resolution grid of size n , the left-hand side of this latter expression corresponds to the so-called $(p \times n)$ anomaly matrix built from p ensemble members of the dynamical system. The tangent linear approximation provided by ensemble methods can be thus immediately interpreted as a particular instance of feature maps together with a given choice of specific inner product to define the reproducing kernel. Keeping a finite dimensional approximation but working without assuming that the functions $[g_x^\varepsilon]$ belong to \mathcal{H} , and thus with now expression (33) for the ensemble tangent linear expression, corresponds to the case in which a localization procedure identified to the square root operator $\mathcal{L}_{k_t}^{1/2}$ has been considered. These two choices embed the problem within a particular RKHS family of functions. The relation between

the tangent linear dynamics and the anomaly matrix is in our case exact and does not correspond to a finite difference approximation as classically presented in ensemble methods. Wonderland family can be seen as a way of linearizing locally a nonlinear system in a convenient sequence of spaces of smooth functions.

4.3 Finite time Lyapunov exponents

The kernel of the Koopman operator provides also a direct access to the finite time Lyapunov exponents. Recalling from (30) that for any punctual observable g_x , as defined previously, we have

$$g_x(\delta X_t) = \int_0^t \partial_X \mathcal{M}(X_s) g_x(\delta X_s) ds + o(\|\delta X_0\|). \quad (37)$$

With the expression of the tangent linear operator in terms of the Koopman infinitesimal generator (33) on Ω_s , we have then

$$g_x(\delta X_t) = \int_0^t \mathcal{L}_{k_s}^{-1/2} A_{U,s} \mathcal{L}_{k_s}^{1/2} g_x(\delta X_s) ds. \quad (38)$$

For all $s \in [0, t]$, we consider a perturbation $g_x(\delta X_s) = \mathcal{L}_{k_s}^{-1/2} \Psi_\ell^s(\delta X_s)$ along a Koopman generator eigenfunction associated to the eigenvalue of maximal modulus $|\lambda_\ell|$. By theorem 1, we have

$$\partial_t g_x(\delta X_s) = \lambda_\ell \mathcal{L}_{k_s}^{-1/2} \Psi_\ell^s(\delta X_s) = \lambda_\ell g_x(\delta X_s).$$

Therefore we get

$$|g_x(\delta X_t)| = e^{|\lambda_\ell|t} |g_x(\delta X_0)|, \quad (39)$$

and the finite time Lyapunov exponent is consequently defined as

$$\sigma = |\lambda_\ell|. \quad (40)$$

For regular perturbations $g_x \in \mathcal{H}_s$ with unitary perturbation $g_x(\delta X_s) = \Psi_\ell^s(\delta X_s)$, the derivation is even simpler as we obtain from remark 7

$$\partial_t g_x(\delta X_s) = A_{U,t} g_x(\delta X_s),$$

which yields directly to expression (39) and to the same expression for the Lyapunov exponent.

The modulus of the larger Koopman eigenvalue in Wonderland provides thus an estimate of the Lyapunov exponent. It can be outlined that the computation of Lyapunov exponents for large scale systems is computationally very demanding as it requires the construction of the linear tangent dynamics operator and the solution of an eigenvalue problem of very big dimension. The construction of the exact numerical tangent linear operator is in general a tedious task when expressed in $L^2(\Omega_x)$, as in equation (29). The ensemble based method provided by our formalism is on the contrary very simple by expressing the tangent linear operator in

$L_{\mathbb{C}}^2(\Omega_t, \nu)$, as in equation (30). It can be noticed that, by this change of norm in the definition, the computed values are not the same.

Three distinct values can then be defined for practical computations. First, the Lyapunov spectrum expressed in $L_{\mathbb{C}}^2(\Omega_t, \nu)$ can be determined by computing the singular values of $\mathcal{L}_{k_t}^{-1/2} A_{U,t} \mathcal{L}_{k_t}^{1/2}$. The time integral is dropped since the evaluations are constant along trajectories. It can be viewed as an advantage of working in $L_{\mathbb{C}}^2(\Omega_t, \nu)$ instead of $L^2(\Omega_x)$ as performed classically. The time independence is due to the fact that the Koopman operator is intrinsic to the dynamical system. However from a numerical point of view as the computation is performed in practice through an ensemble with a limited number of members, the learned spectrum is representative only of the local dynamics at the time ($t = t_0$) at which the kernel has been evaluated. As an alternative, modal Lyapunov exponents can be defined by the square root of the first singular values of $\mathcal{L}_{k_t}^{-1/2} \Psi_{\ell} |\lambda_{\ell}|^2 \Psi_{\ell}^* K_t^{-1} \mathcal{L}_{k_t}^{1/2} = \mathcal{L}_{k_t}^{-1/2} \Psi_{\ell} |\lambda_{\ell}|^2 \Psi_{\ell}^* \mathcal{L}_{k_t}^{-1/2}$ with $K_t(i, j) = k_t(X_t^{(i)}, X_t^{(j)})$. We call these singular values the *Koopman modal Lyapunov exponents* (KMLE). Finally, as a third option, equation (40) can simply be considered to evaluate modal exponents. We can notice that the two modal Lyapunov exponents definitions are very similar; the former being expressed in $L_{\mathbb{C}}^2(\Omega_t, \nu)$ and the latter in \mathcal{H}_t . An example of the estimation of the two types of modal exponents will be presented for a quasi-geostrophic dynamics in the numerical section.

4.4 Practical considerations

Let us stress again that in practice, we only have access to the mappings k_t and $A_{U,t}$ with $t \geq 0$. The mapping k and A_U are completely inaccessible for high-dimensional systems as they require the complete knowledge of the phase space or at least of a long enough orbit with a density assumption in the whole phase space. This last assumption is associated to strong requirements of the dynamical system and is not necessarily valid for a given time series of a particular observable. Instead of working with an infinite (dense) trajectory, Theorem 1 enables us to estimate the eigenvalues and eigenfunctions of the Koopman operator locally in the ‘‘Wonderland’’ RKHS family, which can locally conveniently be accessed from an ensemble of finite time trajectories. As it will be described in the following, operators $A_{U,t}$, can be discretized as an ensemble matrix – itself related, as we saw it previously, to the tangent linear dynamics operator. This matrix is then diagonalized to get access to Koopman eigenvalues and their associated eigenfunctions. In theory, the diagonalization of $A_{U,t}$ needs to be performed only once, at a given time, to access the Koopman eigenpairs $(\psi_{\ell}^t)_{\ell}$ and $(\lambda_{\ell})_{\ell}$. However, the exponential relation between distinct instants allows us also to consider averaging strategies to eventually robustify the estimation in practice. This capability will be exploited in the numerical section for the data assimilation of time series.

Diagonalization in practice

For all $t \geq 0$, let m_t be the kernel expression of the operator $A_{U,t}$ given by $m_t(X_t, Y_t) := A_{U,t}^*[k_t(\cdot, Y_t)](X_t)$ for all X_t and $Y_t \in \Omega_t$. By proposition 8, we have $m_t(X_t, Y_t) = -A_{U,t}[k_t(\cdot, Y_t)](X_t)$.

Let us denote by $\{X_t^{(i)} : 1 \leq i \leq n\}$ an ensemble of members generated by the dynamical system and by $\{k_t(\cdot, X_t^{(i)}) : 1 \leq i \leq n\}$ the n associated feature maps. For all time $t \geq 0$, these n feature maps enable us to build a kernel expression of operator $A_{U,t}$ as the $n \times n$ matrix $\mathbb{M}_t = \left(\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) \right)_{1 \leq i, j \leq n}$ with:

$$\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) := -A_{U,t}[k_t(\cdot, X_t^{(j)})](X_t^{(i)}) = [-\partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t^{(j)})](X_t^{(i)}).$$

As shown in the following, this matrix enables us to access to the Koopman generator eigenvalues and to the evaluation of the eigenfunctions at the ensemble members.

By definition of $A_{U,t}$, we have $A_{U,t}[k_t(\cdot, X_t^{(j)})](X_t^{(i)}) = A_U[k(\cdot, X_t^{(j)})](X_t^{(i)})$. Otherwise, as $-\mathcal{L}_k^{1/2} \circ A_U[k(\cdot, X_t^{(j)})]$ belongs to \mathcal{H} , by the proof (Step 1) (D) of Theorem 1 and denoting $(\psi_\ell)_\ell$ an orthonormal basis of \mathcal{H} sets from the eigenfunctions of A_U , we get

$$\begin{aligned} -\mathcal{L}_k^{1/2} \circ A_U[k(\cdot, X_t^{(j)})](X_t^{(i)}) &= -\sum_{\ell=0}^{\infty} \langle -\mathcal{L}_k^{1/2} \circ A_U[k(\cdot, X_t^{(j)})], \psi_\ell \rangle_{\mathcal{H}} \psi_\ell(X_t^{(i)}), \\ &= -\sum_{\ell=0}^{\infty} \langle A_U k(\cdot, X_t^{(j)}), \mathcal{L}_k^{-1/2} \psi_\ell \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \psi_\ell(X_t^{(i)}). \end{aligned}$$

Noting β_ℓ the eigenvalues of \mathcal{L}_k , we have also (step 1 D theorem 1) $\mathcal{L}_k \psi_\ell(X_t^{(j)}) = \beta_\ell \psi_\ell(X_t^{(j)})$ and

$$-\mathcal{L}_k^{1/2} \circ A_U[k(\cdot, X_t^{(j)})](X_t^{(i)}) = -\sum_{\ell=0}^{\infty} \beta_\ell^{-1/2} \langle A_U k(\cdot, X_t^{(j)}), \psi_\ell \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \psi_\ell(X_t^{(i)}).$$

By the skew symmetry of the generators (proposition 4) we have that

$$\begin{aligned} -\mathcal{L}_k^{1/2} \circ A_U[k(\cdot, X_t^{(j)})](X_t^{(i)}) &= \sum_{\ell=0}^{\infty} \lambda_\ell \beta_\ell^{-1/2} \langle k(\cdot, X_t^{(j)}), \psi_\ell \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \psi_\ell(X_t^{(i)}) \\ &= \sum_{\ell=0}^{\infty} \lambda_\ell \beta_\ell^{-1/2} \overline{\mathcal{L}_k \psi_\ell(X_t^{(j)})} \psi_\ell(X_t^{(i)}), \end{aligned}$$

which leads to

$$-\mathcal{L}_k^{1/2} \circ A_U[k(\cdot, X_t^{(j)})](X_t^{(i)}) = \sum_{\ell=0}^{\infty} \lambda_\ell \beta_\ell^{1/2} \overline{\psi_\ell(X_t^{(j)})} \psi_\ell(X_t^{(i)}),$$

and by continuity of $\mathcal{L}_k^{-1/2}$

$$\begin{aligned} -A_U[k(\cdot, X_t^{(j)})](X_t^{(i)}) &= \sum_{\ell=0}^{\infty} \lambda_{\ell} \beta_{\ell}^{1/2} \overline{\psi_{\ell}(X_t^{(j)})} \mathcal{L}_k^{-1/2} \psi_{\ell}(X_t^{(i)}) \\ &= \sum_{\ell=0}^{\infty} \lambda_{\ell} \overline{\psi_{\ell}(X_t^{(j)})} \psi_{\ell}(X_t^{(i)}). \end{aligned}$$

By the restriction expression (remark 17, B), we finally obtain the following equality for all $t \geq 0$

$$\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) = \sum_{\ell=0}^{\infty} \lambda_{\ell} \psi_{\ell}^t(X_t^{(i)}) \overline{\psi_{\ell}^t(X_t^{(j)})},$$

which shows that the diagonalization of \mathbb{M}_t provides a set of the Koopman generator eigenpairs in Wonderland.

In practice, the skew-symmetric matrix $\mathbb{M}_t := (\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}))_{1 \leq i, j \leq n}$ is assembled from the definition of $A_{U,t}$ and a given choice of the kernel. As explained in the previous section this matrix corresponds to a kernel expression of $A_{U,t}$ with

$$\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) := -A_{U,t}[k_t(\cdot, X_t^{(j)})](X_t^{(i)}) = [-\partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t^{(j)})](X_t^{(i)}).$$

This matrix can be interpreted in the RKHS setting as resulting from the matrix multiplication:

$$\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) := \sum_{\ell} \mathbb{F}_t(j, \ell) K_t(\ell, i),$$

with $\mathbb{F}_t(i, \ell) = -A_{U,t}[k_t(Z^{\ell}, X_t^{(j)})]$. This indeed corresponds to a discretization of the kernel expression of operator $A_{U,t}$ through the empirical Dirac measure. Numerically, instead of working with matrix \mathbb{M}_t (i.e. the evaluation of $A_{U,t}k_t(\cdot, X)$ at several discrete points), we will work directly with matrix $\mathbb{F}_t = \mathbb{M}_t K_t^{-1}$. This has the advantage of directly working with an implicit discretization of operator $A_{U,t}$, and to relax somewhat its dependency on the kernel choice. The skew-symmetric matrix $\mathbb{M}_t K_t^{-1}$ is then diagonalized through a direct numerical procedure (using LAPACK library and working numerically on the anti-symmetric part of $\mathbb{M}_t K_t^{-1}$) and can be written $\mathbb{M}_t K_t^{-1} = V_t \Lambda V_t^T$, with V_t a unitary matrix and Λ a diagonal matrix. The matrix V_t gathers eigenvectors V_j^t of \mathbb{F} , which as a discretization of $A_{U,t}k_t(\cdot, X)$ gives access to the values of the Koopman eigenfunctions $\psi_j^t = \mathcal{L}_k^{1/2} V_j^t$ at the n ensemble members points $X_t^{(i)}$. The matrix Λ is composed of Koopman eigenvalues with conjugate pairs of pure imaginary eigenvalues.

As previously mentioned this diagonalization can be performed at a single time or at several instants accompanied with an averaging procedure. Theorem 1 and (7), give access to the eigenvectors evaluation along a trajectory for all time instants.

We provide below as examples, expressions of the evaluation of $\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) = \left(-\partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t^{(j)})\right) \left(X_t^{(i)}\right)$ for the empirical covariance kernel (k_E) and the Gaussian kernel (k_G). The empirical covariance kernel is defined through the kernel isometry property as

$$k_t(X_t^{(i)}, X_t^{(j)}) = \left\langle X_0^{(i)}, X_0^{(j)} \right\rangle_{\Omega_x},$$

with $(\cdot, \cdot)_{\Omega_x}$ the inner product of $L^2(\Omega_x, \mathbb{R}^d)$ and where Ω_x denotes the physical domain of the considered dynamics. We obtain

$$\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) = \left\langle \frac{\partial}{\partial t} X_0^{(j)}, X_0^{(i)} \right\rangle_{\Omega_x}. \quad (41)$$

In this expression, we see that the time derivative of the ensemble members at the initial time is required. Similarly, the Gaussian kernel is defined as

$$k_t(X_t^{(i)}, X_t^{(j)}) = \exp\left(-\frac{1}{\ell^2} \left\| X_0^{(i)} - X_0^{(j)} \right\|_{\Omega_x}^2\right),$$

This leads to

$$\mathbb{M}_t(X_t^{(i)}, X_t^{(j)}) = -\frac{2}{\ell^2} \left\langle \frac{\partial}{\partial t} X_0^{(j)}, \left(X_0^{(i)} - X_0^{(j)}\right) \right\rangle_{\Omega_x} \exp\left(-\frac{1}{\ell^2} \left\| X_0^{(i)} - X_0^{(j)} \right\|_{\Omega_x}^2\right). \quad (42)$$

Thanks to the isometry property, this matrix needs to be evaluated only at a single time.

5 Application to ensemble methods

5.1 Quasi-geostrophic model

Based on a set of p ensemble members, the ability of the proposed method to predict a trajectory associated with a new initial condition is demonstrated on a quasi-geostrophic barotropic model of a double-gyre [35] in a “rectangular-box ocean”. This idealistic model of North-Atlantic large-scale circulation considers the transport of the ageostrophic state variable components by a velocity in geostrophic balance. The barotropic vorticity equation written for the potential vorticity q and the stream function ψ in $\Omega_x = ([0 : L] \times [-L : L])$, with $L = 1$, is

$$\begin{cases} \frac{\partial q}{\partial t} + J(\psi, q) = f - (\delta/L)^5 \nabla^4 \omega \\ \Delta \psi = \omega \quad ; \quad q = R_o \omega + y_c, \end{cases} \quad (43)$$

where ω is the vorticity and $x_c = (x_c, y_c)$ are the latitude and the longitude increasing eastward and northward, respectively, while $J(f, g) = \partial_x f \partial_y g - \partial_x g \partial_y f$ stands for the Jacobian (with x and y denoting horizontal and vertical coordinates of \mathbb{R}^2). The dimensionless Rossby number $R_o = 0.0036$ measures the ratio between inertial

forces and the Coriolis force, and a steady wind-forcing is defined as $f = \sin(\pi y_c)$. The parameter $\delta/L = 0.032$ controls the hyper-viscosity model, with δ the Munk boundary layer width. Scales smaller than δ will not be resolved. Finally, we note q' the ensemble anomaly of the potential vorticity. Numerical details and links with dimensional variables associated with a mid-latitude ocean basins, such as North Atlantic, are given in E.

5.2 Reconstructions using RKHS

In this study we focus only on two specific kernels: the empirical covariance kernel (k_E), and the Gaussian Kernel (k_G). They are defined respectively by

$$k_E(X_t^{(i)}, X_t^{(j)}) = \left\langle q_t'^{(i)}(\cdot, t_0), q_t'^{(j)}(\cdot, t_0) \right\rangle_{\Omega_x},$$

and


$$k_G(X_t^{(i)}, X_t^{(j)}) = \exp \frac{1}{\ell^2} \left\| q_t^{(i)}(\cdot, t_0) - q_t^{(j)}(\cdot, t_0) \right\|_{\Omega_x}^2,$$

where $\langle \cdot, \cdot \rangle_{\Omega_x}$ stands for the standard L^2 inner product in Ω_x . The former has the advantage to be simple, and to be directly related to standard ensemble methods employed in data assimilation. However, it is strongly rank deficient and introduces spurious correlations in the physical domain Ω_x between two far apart locations. Localization techniques [37] have been introduced to tackle this problem and may be interpreted in our setting as embeddings in specific RKHS. This interpretation may open new computationally efficient generalisations related for instance to localization techniques in the model space as explained after remark 7. The parallel done between operators $\mathcal{L}_{k_t}^{1/2}$ and localization procedure would in particular immediately enables to set up time-varying and flow dependant localization techniques, which are difficult to built. Gaussian kernels are widely used in machine learning for its high regularity and non-compact support in $L^2(\Omega, \nu)$ [48]. It can be also interpreted as a way to perform localization. In this study, compared to other tested – but not presented – kernels (polynomial and exponential for instance), it led to better performances.


Once the kernel matrix $K_{ij} = k_t(X_t^{(i)}, X_t^{(j)})$ for $(i, j) \in [1 \dots p] \times [1 \dots p]$ (equal for all t – kernels isometry, Theorem 3) is evaluated, a very simple reconstruction of the whole trajectory can be performed. Denoting X_0' a new ensemble member initial condition, the linear combination vector β associated to the reconstructed trajectory $\hat{X}_t' = \sum_{i=1}^p \beta_i X_t^{(i)}$ can be estimated by kernel regression from the evaluation $k(X_0', X_0^j)$, for $j = [1 \dots p]$, and the reproducing property as

$$\beta = K^{-1} k(X_0', X_0^{[1 \dots p]}). \quad (44)$$

Strikingly, Koopman isometry implies that these coefficients are constant along time, and brings hence a direct reconstruction of the whole trajectory.



(a) Lyapunov times spectrum expressed in $L_{\mathbb{C}}^2(\Omega_t, \nu)$ for the Gaussian kernel k_G and the empirical covariance k_E .



(b) Modal Lyapunov times associated with individual Koopman modes. Comparison with expression in $L_{\mathbb{C}}^2(\Omega_t, \nu)$ and in \mathcal{H}_t .

Figure 1: Lyapunov times computed using the Gaussian kernel k_G and the empirical covariance kernel k_E .

5.3 Evaluation of the Koopman eigenfunctions

Performing the \mathcal{H}_t inner product of equation (6) with the ensemble members' feature maps leads to the infinitesimal generator matrix

$$\mathbb{M}(X_t^{(i)}, X_t^{(j)}) = \left(-\partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t^{(j)}) \right) \left(X_t^{(i)} \right). \quad (45)$$

As explained previously eigen-elements of $\mathbb{M}K^{-1}$, noted $(\lambda_\ell, V_\ell(X_t^{(i)}))$, give access to the exact evaluation of Koopman eigenfunctions on the RKHS family at the ensemble members $\psi_\ell^t(X_t^{(i)}) = \mathcal{L}_k^{1/2} V_\ell(X_t^{(i)})$, and to the associated eigenvalues λ_ℓ . We note that $\partial_{\mathcal{M}(\cdot)} k_t(\cdot, X_t^{(j)})$ is obtained by differentiating the kernel expression (see equations 41, 42 for the two selected kernels). Skew-symmetry of $\mathbb{M}K^{-1}$, inherited from the skew-symmetry of $A_{U,t}$, is only approached numerically. Evaluations are thus enhanced by considering only the skew-symmetric part of the matrix $\mathbb{M}K^{-1}$, which is dominant in the numerical evaluation of (45).

5.4 Lyapunov times

The Lyapunov times evaluated using the two kernels k_E and k_G are compared. Three methods to compute the Lyapunov exponents σ are detailed in section 4.3. The associated Lyapunov times are defined by $T = \sigma^{-1} \ln C$, with, $C = 10^3$.

The Lyapunov times associated with exponents expressed in $L_{\mathbb{C}}^2(\Omega_t, \nu)$ are shown in figure 1(a). The spectrum associated with all singular values, truncated such that $\sigma > 10^{-5}$, is displayed. Due to this truncation, very slow modes are not well captured. We can see that the Gaussian kernel k_G estimates in average larger Lyapunov times.

In figure 1(b), the Koopman modal Lyapunov times are shown as a function of the absolute value of the Koopman eigenvalue. The Lyapunov times expressed with the RKHS norm equation (40) is directly $T_\ell = 1/|\lambda_\ell| \ln C$, which explains the




Figure 2: Comparison between the reference potential vorticity of a test ensemble member at $t = 0.07$ (corresponding to 3.7 days of prediction) (a) and the predicted reconstruction using the Gaussian kernel ($\ell = 1$) (c). The best possible reconstruction using a linear combination of ensemble members with respect to the L^2 norm is shown in (b).

perfect decaying slope in logarithmic scale. We observe that the computation of the modal exponents in $L^2_{\mathbb{C}}(\Omega_t, \nu)$ leads simply to a vertical shift of the curves due to the scaling factor induced by the change of norm. We can notice that with the Gaussian kernel k_G , slower Koopman eigenfunctions are evaluated compared to k_E , which leads to longer Lyapunov times. This explains longer predictions, which will be presented in section 5.5. As it will be seen, the modal exponents and their associated Lyapunov times will reveal very useful, as they will allow us to filter out sequentially the contributions of the Koopman eigenfunctions beyond their predictability time.

5.5 Reconstructions

The ability of trajectories reconstruction is illustrated in figure 2, where a full trajectory is estimated from the embedding of a new initial condition in the RKHS family. We compare the anomaly (figure 2(a)) of a given new ensemble member at an advanced time (3.7 days), with reconstructions using the Gaussian kernel k_G (figure 2(c)). Figure 2(b) shows the best possible reconstruction as a linear combination of ensemble members, with regards to the $L^2(\Omega_x)$ -norm of the projection error.

Reconstruction errors are quantified by averaging the square of the $L^2(\Omega_x)$ -norm over the whole test ensemble composed of 100 new members (i.e. not used in the training phase). The standard deviations of the error are also displayed on this figure. The time evolution of this error is presented in figure 3. It is compared to the projection error defined as the minimal linear combinations of ensemble members w.r.t. $L^2(\Omega_x)$ -norm. The reconstruction error performed by the ensemble average is also shown. This error is obviously high for small times, but estimations are difficult to be more precise than the ensemble average estimation once the pre-




Figure 3: Mean estimation error using an empirical covariance kernel (short red dashes), a Gaussian kernel (green dash-dots), the reconstructed Gaussian kernel using all Koopman eigenfunctions (orange squares) and the Lyapunov filtered kernel (blue dot-dashes). “Ensemble average” (long black dashes) is the estimation error of the ensemble average. The projection error (purple solid line) corresponds to the optimal time-varying linear combination of ensemble members w.r.t. the $L^2(\Omega_x)$ -norm. The vertical width of the stripes regions over the mean error is equal to the standard deviation of the error. One dimensionless time t corresponds to 92 days. Errors are averaged over the 100 members of the test ensemble.

dictability time of the system has passed. The reconstructions performed with k_E , grow and overtake the ensemble average reconstruction around $t = 0.2$ (i.e. 18.4 days). On this basis, the use of k_G ($\ell = 1.0$) leads to better results. Indeed, despite a slight reconstruction penalty at the initial time due to a stronger smoothing (see figure 2), the reconstruction errors become quickly significantly lower, and the error never reaches the ensemble anomaly error. This small loss of performance in $L^2(\Omega_x)$ -norm at the initial time allows to have a gain in robustness in terms of long-time prediction.

The reconstructions based on (44) are valid for all times. Koopman isometry warrants the use of the same linear combination for the whole trajectory. Nevertheless, in practice the ensemble are of limited (small) size, and the eigenpairs of the infinitesimal generator close to zero (representing almost stationary functions) are difficult to estimate accurately. Modal predictability time associated to Lyapunov exponents (39) can be used to robustify the forecast of new initial conditions. The kernel can be expanded in the Koopman eigenfunctions basis as

$$k(X_t^{(i)}, X_t^{(j)}) = \sum_{j=1}^{\infty} \psi_\ell^t(X_t^{(i)}) \overline{\psi_\ell^t(X_t^{(j)})} = \sum_{j=1}^{\infty} \psi_\ell^0(X_0^{(i)}) \overline{\psi_\ell^0(X_0^{(j)})}. \quad (46)$$

With the numerical procedure proposed here, the p evaluated Koopman eigenfunctions allow to perform an exact reconstruction of the kernel. An appealing potential of equation (46) for faithful predictions consists in filtering the kernel by switching-off components that have passed their predictability time. We set this time from the finite-time Lyapunov exponents (39) as: $T_\ell = |\lambda_\ell|^{-1} \ln C$, with, $C = 10^3$. The filtered kernel $\tilde{k}(X_t^{(i)}, X_t^{(j)})$ and its associated $p \times p$ kernel matrix \tilde{K} allows us to

define reconstructions, with the modified linear coefficients $\tilde{\beta} = \tilde{K}^+ \tilde{K} \beta$. The superscript \bullet^+ stands for the Moore–Penrose pseudo-inverse. It can be remarked that the operator $\tilde{K}^+ \tilde{K}$ is a projector performing filtering directly on the vector of coefficients β .

In figure 3, it is assessed that the kernel is exactly reconstructed by the Koopman eigenfunctions (superimposition of the green dash-dots curve and the orange squares). When the kernel is Lyapunov filtered, the reconstruction almost fits at short time the previous predictions (superimposition of the blue dot-dashes curve, the green dash-dots curve and the orange squares). For longer time it significantly reduces the error and clearly separates the spread of the ensemble from the ensemble average estimation. For very long times (not shown in the figure), filtering acts on all modes, which reduces to the ensemble average estimation. This strategy takes advantage of the modal predictability time. Slow eigenfunctions are used for long time predictions, and fast ones only for short time forecasts. As a final remark, we note that very fast modes (with large Koopman eigenvalues modulus) does not seem to contribute significantly to the reconstruction, since at short times the filtered and full reconstructions match.

5.6 Data assimilation

A possible application of the proposed procedure is ensemble-based data assimilation. In the context of satellite data, ensemble optimal interpolation (EnOI) is widely used to reconstruct spatial fields [42, 29]. It is for instance employed in operational gridded altimetry reconstructions [5]. EnOI consists in expressing at a given time the solution as a linear combination of ensemble members, which minimises an objective cost function. As shown in [62, 36], incorporating information from other times together with an adequate dynamical model significantly improves the estimations. This is obviously at the price of forecasting the whole ensemble by the dynamics.

Koopman isometry provided by Wonderland allows us to incorporate time-series of observations without any new simulation of the dynamics. The spatial reconstruction is embarrassingly simply given as the constant-in-time best linear combination of ensemble members trajectories.

Based on the same ensemble as in the previous sections, we propose two demonstration tests considering *i*) a time-series of extremely sparse (two satellite swaths like) observations *ii*) an even sparser case with the same swaths but at a single time only. To that end, we consider a synthetic observation mimicking two along tracks measurements of an orbiting satellite, idealized by two vertical lines. In the first case these lines are assumed to be known every 0.9 days $\Delta t = 0.01$, while in the second case only day 5.5 is available ($t = 0.06$). The observations are perturbed by a centered Gaussian white noise with the value of the selected standard deviation equal to 10% of the signal root-mean-square. These 2 benchmarks are carried out for each of the 100 members of the test ensemble. As previously, the members of the test ensemble have not been used in the learning phase.

In this numerical experiment, we search for the constant in time coefficients β (thanks to isometry), which minimises

$$\mathcal{J} = \frac{1}{2T} \int_0^T \|\mathbb{H}(\hat{X}_t) - \mathcal{Y}\|_{R^{-1}}^2 dt + \frac{\alpha^2}{2} \|\beta\|^2 \text{ s.t. } \mathbb{H}(\hat{X}_t) = \sum_{i=1}^p \beta_i \mathbb{H}(X_t^{(i)}), \quad (47)$$

where \mathbb{H} is the observation operator associated with the satellite measurements \mathcal{Y} , \hat{X}_t is the estimated state. The observation operator is assumed to have its range in the RKHS family and hence provides observables belonging to the RKHS family, which can be interpolated using the Wonderland kernels. It can be noticed that the expression of the constraint is due to kernel interpolation, and not to some linear property of \mathbb{H} . This treatment is similar to the one performed for instance in discrete empirical interpolation (DEIM) [20]. In the case of a single observation, the time integral is dropped, and the cost function corresponds to a standard EnOI cost function. The matrix R is the observation covariance matrix, defined by the time average of the empirical covariance, localized by performing a Schur product with matrix $\exp(-r_{ij}^2/\ell_{loc}^2)$, where r_{ij} is the distance between measurements i and j , and $\ell_{loc} = 0.05$. The regularization parameter has been set to $\alpha = 10$. For this problem we selected the simple empirical kernel, defined from a non localized potential vorticity matrix as feature maps of the ensemble. Working with the isometry property justifies the estimation of constant in time coefficients for the linear combination. This simplistic method could be easily extended by introducing more elaborated kernels, Koopman eigenfunctions representation to project any system's observable and forecast them, as well as the Lyapunov filtering scheme presented in section (5.5).

Figure 4 shows the potential vorticity trajectory (panel a), the single-time observation (panel b), the trajectory reconstruction associated to this single observation (panel c) and the one obtained from the swaths time-series (panel d). Three consecutive times spaced of 0.02 (1.8 days) are displayed. We can see that despite extremely sparse measurements, a single observation already provides a fairly good estimation, together with forward and backward in time forecasts based only on the estimated linear combination of the ensemble members (i.e. without any simulation). As expected, a time-series of swaths improves further the performances.

In figure 5, the average of the errors obtained over all 100 members of the test ensemble are plotted. As a reference, the best possible reconstruction with respect to the L^2 norm with constant in time linear coefficients β is shown, as well as the estimation by the ensemble average. The error performed with the time-series of satellite observations is almost optimal in terms of average and standard deviation. We note that for the second test-case, near the observation time, the error is lower than the one obtained for the whole time-series. Compared to the time series of observations it may be interpreted as an over-fitting at the observation time since only a single observation is used to infer the ensemble members' coefficients. As expected, this error grows, but reasonably, as the estimation is propagated forward and backward in time.




Figure 4: Estimation of potential vorticity at times $t = 0.04$ (top), $t = 0.06$ (middle) and $t = 0.08$ (bottom). Panel (a): True anomaly. Panel (b): Observation associated with the single-time swath observation. Panel (c): Estimation knowing the two satellite swaths only at day 5.5 ($t = 0.06$). Panel (d): Estimation knowing the two swaths every 0.9 days ($\Delta t = 0.01$).




Figure 5: Estimation errors averaged over the whole 100 members of the test ensemble. The best reconstruction error with constant coefficients (red), estimation by the ensemble average (black), estimation with the time-series of satellite swaths observations (yellow) and only at day 5.5 ($t = 0.06$) (blue).

6 Concluding remarks

We proposed a theoretical setting to estimate the Koopman eigen-pairs of compact dynamical systems in embedding the system in a RKHS family, called Wonderland. Beyond the Koopman operator spectral representation, we showed that Wonderland provides access to finite time Lyapunov exponents, as well as to the tangent linear dynamics. These features enabled us to propose very simple strategies for data assimilation of very sparse observations in time and space. Obviously more efficient data assimilation techniques could be used instead of the simple regression strategy explored here. As Wonderland boils down to a linear setting it should be perfectly adapted to ensemble Kalman filters. An extension of the filter proposed in [34] could for instance be investigated. This technique combines, similarly to our framework, an RKHS to specify an evolution equation and an ensemble data assimilation strategy for inference of unknown trajectories. The main difference lies in the fact that here we propose a sequence of time-evolving RKHS to represent the evolution operator. The dynamics manifold can be by this way gradually unveiled. This study provides a theorem-based setting to estimate the Koopman eigenvalues and eigenfunctions, as well as exact ensemble-based representation of the tangent linear dynamics and of the flow Lyapunov exponents. This work relies on the assumption of an invertible deterministic dynamical system defined on a compact manifold. In future studies, we will focus on the relaxation of the invertibility, the compactness assumptions, and on the generalisation to stochastic fluid flow dynamics with transport noise models [45, 8]. Let's conclude by playing around with Lewis Carroll's novel: "Who are you?" said the Caterpillar to Alice. "Well, I barely don't know, being so many different sizes in a day is very confusing" answered Alice. "It isn't! Duplicate yourself! Shrink and expand repeatedly! And gather all your differences to reveal the stable harmonic parts of yourself." ended the Caterpillar.

Acknowledgments

The authors acknowledge the support of the ERC project 856408-STUOD and the French National program LEFE (Les Enveloppes Fluides et l'Environnement)

A Reproducing kernel Hilbert spaces

In the following appendices we provide complementary elements for the detailed proofs of the results listed in the paper. We first recall some classical theorems on RKHS and present their adaptations to our functional setting. In particular we present useful results on differentiability of RKHS functions and reproducing kernels. In the following $L^2(\Omega_x)$ stands for the Hilbert space of Lebesgue measurable \mathbb{R}^d -valued of square-integrable functions on the set $\Omega_x \subset \mathbb{R}^p$ $p = 2, 3$ endowed at

a fixed time, t , with the inner product

$$\langle f, g \rangle_{L^2(\Omega_x)} = \int_{\Omega_x} f(y) g(y) dy.$$

Let $E \subset L^2(\Omega_x)$ denote a compact set equipped with the norm of $L^2(\Omega_x)$, by the Aronszajn Theorem, [4], the reproducing kernel Hilbert spaces (RKHS) of \mathbb{C} -valued functions defined on E are uniquely characterized through a reproducing kernel $k : E \times E \mapsto \mathbb{C}$. This kernel is a Hermitian, positive definite function. In the following we will assumed furthermore it is $C^2(E \times E)$, which means here it is twice differentiable with respect to its two variables. The derivative with respect to the first and second variable in the direction $v \in E$ are denoted $\partial_v^{(1)}k$, and $\partial_v^{(2)}k$, respectively. The reproducing kernel defines an evaluation function $k(\cdot, x) \in \mathcal{H}$ with a reproducing property (Def. 1):

$$\forall x \in E, \forall f \in \mathcal{H}, \langle f, k(\cdot, x) \rangle_{\mathcal{H}} = f(x).$$

The RKHS \mathcal{H} corresponds to a space of smooth functions with a strong uniform convergence property. The elements of \mathcal{H} are in addition continuous functions and we have for all $f \in \mathcal{H}$ that

$$\forall x \in E \quad |f(x)| \leq \sup_{y \in E} k(y, y)^{1/2} \|f\|_{\mathcal{H}}. \quad (48)$$

A.1 Functional description of \mathcal{H}

We recall in the following the Mercer's theorem which gives a functional description of the RKHS \mathcal{H} and of its associated reproducing kernel k .

Let $L^2_{\mathbb{C}}(E, \mu)$ be the space of measurable complex valued functions defined on the compact set E for which the square modulus is integrable with respect to measure μ , $L^2_{\mathbb{C}}(E, \mu)$ is a Hilbert space equipped with the inner product $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(E, \mu)}$ given for f and $g \in L^2_{\mathbb{C}}(E, \mu)$ by

$$\langle f, g \rangle_{L^2_{\mathbb{C}}(E, \mu)} := \int_E f(y) \overline{g(y)} \mu(dy),$$

so that it is linear in its first argument and antilinear in its second. Let us first define $\mathcal{L}_k : L^2_{\mathbb{C}}(E, \mu) \mapsto L^2_{\mathbb{C}}(E, \mu)$ the integral operator with kernel $k \in C^1(E \times E)$ as:

$$(\mathcal{L}_k f)(x) = \int_E k(x, y) f(y) \mu(dy). \quad (49)$$

We give below a useful remark relating the integral operator \mathcal{L}_k and the inner product $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(E, \mu)}$.

Remark 8 (Links between \mathcal{L}_k and $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(E, \mu)}$) As k is Hermitian, it can be noticed that for all $x \in E$, $(\mathcal{L}_k f)(x) = \int_E f(y) k(y, x) \mu(dy)$, and we have thus $(\mathcal{L}_k f)(x) = \langle f, k(\cdot, x) \rangle_{L^2_{\mathbb{C}}(E, \mu)}$.

This kernel integral operator can be shown to be well-defined and compact (by Arzelà–Ascoli theorem). It is immediate to see that as k is positive definite and Hermitian, \mathcal{L}_k is positive and self-adjoint on $L^2_{\mathbb{C}}(E, \mu)$. By the spectral theorem of compact self-adjoint operators, there exists an orthonormal basis $(\varphi_i)_i$ of $L^2_{\mathbb{C}}(E, \mu)$ consisting of eigenvectors of \mathcal{L}_k with corresponding decreasing real eigenvalues $(\mu_i)_i : \mu_i \geq \mu_{i+1} \geq 0$ for all integers i . These results are proved in [23]. We assume in the following that $\mu_i \neq 0$ for all i and \mathcal{L}_k is injective. This denseness assumption could be relaxed (see remark 15) but simplifies greatly the presentation. The eigenpairs $(\mu_i, \varphi_i)_{i \in \mathbb{N}}$ enables to formulate the following theorems (with proofs available for instance in [23] and [39]). Elaborating on remark 8 and the kernel eigenfunctions (often referred to as Mercer’s eigenfunctions due to theorem 4) we get immediately relations between the eigenfunction evaluation and the $L^2(E, \mu)$ inner product.

Remark 9 (Links between $(\varphi_i)_{i \geq 0}$ and $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(E, \mu)}$) For all $x \in E$, by remark 8, we note that

$$\begin{aligned} \langle \varphi_i, k(\cdot, x) \rangle_{L^2_{\mathbb{C}}(E, \mu)} &= \mathcal{L}_k(\varphi_i)(x) = \mu_i \varphi_i(x) \quad \text{and} \\ \langle k(\cdot, x), \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)} &= \overline{\langle \varphi_i, k(\cdot, x) \rangle_{L^2_{\mathbb{C}}(E, \mu)}} = \mu_i \overline{\varphi_i(x)}. \end{aligned}$$

Remark 10 As $k \in C^1(E \times E)$, the range of \mathcal{L}_k is included in $C^1(E)$ and the eigenfunctions of \mathcal{L}_k are $C^1(E)$.

Theorem 4 (Mercer decomposition) For all x and $y \in E$, $k(x, y) = \sum_{i=0}^{\infty} \mu_i \varphi_i(x) \overline{\varphi_i(y)}$ where the convergence is absolute (for each $x, y \in E \times E$) and uniform on $E \times E$.

In particular, the series $\sum \mu_i$ is convergent as

$$\int_E k(y, y) \mu(dy) = \int_E \sum_{i=0}^{\infty} \mu_i \varphi_i(y) \overline{\varphi_i(y)} \mu(dy) = \sum_{i=0}^{\infty} \mu_i < \infty. \quad (50)$$

Theorem 5 (Mercer representation of RKHS) The RKHS $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}}, \|\cdot\|_{\mathcal{H}})$ is the Hilbert space characterized by

$$\mathcal{H} = \left\{ f \in L^2_{\mathbb{C}}(E, \mu), f = \sum_{i=0}^{\infty} a_i \varphi_i : \sum_i \frac{|a_i|^2}{\mu_i} < \infty \right\},$$

$$\langle f, g \rangle_{\mathcal{H}} = \sum_{i=0}^{\infty} \frac{1}{\mu_i} \langle f, \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)} \overline{\langle g, \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)}}$$

$$\text{and } \|f\|_{\mathcal{H}}^2 = \sum_{i=0}^{\infty} \frac{|\langle f, \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)}|^2}{\mu_i}.$$

The inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ is sesqui-linear (linear in its first argument and antilinear in its second). The proof of Mercer's theorem warrants that the linear space spanned by $\{k(\cdot, x) : x \in E\}$ is dense in $L^2_{\mathbb{C}}(E, \mu)$. If $f = \sum_{i=0}^{\infty} a_i \varphi_i \in \mathcal{H}$ then the series converges absolutely and uniformly to f .

The following remark shows that the feature map's conjugate as well as its real and imaginary parts belong to \mathcal{H} .

Remark 11 (Consequence of the Mercer representation) For all $y \in E$, the functions $\overline{k(\cdot, y)}$, $Re[k(\cdot, y)]$ and $Im[k(\cdot, y)]$ belong to \mathcal{H} . By theorems 5 and 4, we have indeed that

$$\sum_{i=0}^{\infty} \frac{1}{\mu_i} |\langle \overline{k(\cdot, y)}, \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)}|^2 = \sum_{i=0}^{\infty} \frac{1}{\mu_i} |\mathcal{L}_k \varphi_i(y)|^2 = \sum_{i=0}^{\infty} \mu_i |\varphi_i(y)|^2 < \infty, \quad (51)$$

and $\overline{k(\cdot, y)} \in \mathcal{H}$. By operation, $Re[k(\cdot, y)]$ and $Im[k(\cdot, y)]$ belong thus to \mathcal{H} .

Remark 12 (Comparison of the norms \mathcal{H} and $L^2_{\mathbb{C}}(E, \mu)$) For all $f \in \mathcal{H}$, by theorem 5 we have for all $x \in E$, the inequality

$$f(x) \leq |f(x)| = \left| \sum_{i=0}^{\infty} a_i \varphi_i(x) \right| \leq \left(\sum_{i=0}^{\infty} \frac{|a_i|^2}{\mu_i} \right)^{1/2} \left(\sum_{i=0}^{\infty} \mu_i |\varphi_i(x)|^2 \right)^{1/2},$$

with theorem 4 we obtain

$$\|f\|_{L^2_{\mathbb{C}}(E, \mu)} \leq \left(\int_E |k(x, x)|^2 \mu(dx) \right)^{1/2} \|f\|_{\mathcal{H}}. \quad (52)$$

Remark 13 For all $f \in L^2_{\mathbb{C}}(E, \mu)$, $\mathcal{L}_k(f)$ belongs to \mathcal{H} , in fact we have

$$\sum_{i=0}^{\infty} \frac{1}{\mu_i} \left| \langle \mathcal{L}_k(f), \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 = \sum_{i=0}^{\infty} \mu_i \left| \langle f, \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 \leq \left(\sum_{i=0}^{\infty} \mu_i \right) \|f\|_{L^2_{\mathbb{C}}(E, \mu)}^2 < \infty.$$

Moreover, by remark 12 the mapping $\mathcal{L}_k : \mathcal{H} \rightarrow \mathcal{H}$ is continuous : for all $f \in \mathcal{H}$ we have

$$\|\mathcal{L}_k(f)\|_{\mathcal{H}}^2 \leq \left(\sum_{i=0}^{\infty} \mu_i \right) \|f\|_{L^2_{\mathbb{C}}(E, \mu)}^2 \leq C \|f\|_{\mathcal{H}}^2.$$

With a $C^2(E \times E)$ regularity condition on the kernel, we will prove that the RKHS \mathcal{H} is a subspace of smoother functions than $C^0(E, \mathbb{C})$ (theorem 6). In the following appendix we discuss differentiability property of the feature maps.

A.2 Differentiability of feature maps and RKHS functions \mathcal{H}

In this appendix we extend results given in [59, 60] for $E \subset \mathbb{R}^d$ to the case where E is a compact subset of a function space. In the following E designates a compact set included in $L^2(\Omega_x, \mathbb{R}^d)$, the space of measurable functions that are square integrable on Ω_x . Denoting by $\partial_u^{(2)} k(\cdot, y)$ the kernel functional derivative in the direction u with respect to the second variable y – while the upper-script (1) indicates a directional derivative in the first variable –, we have first the following lemma on the feature maps derivative.

Lemma 2 (Feature maps differentiability) *The kernel derivatives belong to \mathcal{H} . More precisely, let $k \in C^2(E \times E)$, then for all $y \in E$ and $u \in E$, we have $\partial_u^{(2)} k(\cdot, y) \in \mathcal{H}$.*

PROOF Let y and $u \in E$. Let $(h_n)_{n \geq 0}$ be a sequence of $\mathbb{C} \setminus \{0\}$ which converges to 0 and such that for all n , $y + h_n u \in E$. For all integers n , let

$$d_n(\cdot, y) := k(\cdot, y + h_n u) - k(\cdot, y) \text{ and } t_n(\cdot, y) := \frac{d_n(\cdot, y)}{h_n}.$$

We first notice that $(t_n(\cdot, y))_{n \geq 0}$ converges pointwise to $\partial_u^{(2)} k(\cdot, y)$ since k is $C^2(E \times E)$. Then, we now show that $(t_n(\cdot, y))_n$ is a Cauchy sequence of $(\mathcal{H}, \|\cdot\|_{\mathcal{H}})$. For all integers p and q , we have

$$\|t_p(\cdot, y) - t_q(\cdot, y)\|_{\mathcal{H}}^2 = \|t_p(\cdot, y)\|_{\mathcal{H}}^2 + \|t_q(\cdot, y)\|_{\mathcal{H}}^2 - 2\text{Re}[\langle t_p(\cdot, y), t_q(\cdot, y) \rangle_{\mathcal{H}}]. \quad (53)$$

By the reproducing property of \mathcal{H} and as k is $C^2(E \times E)$, we have for all $x \in E$ and for some p and q large enough,

$$\begin{aligned} \langle t_p(\cdot, y), t_q(\cdot, x) \rangle_{\mathcal{H}} &= \frac{1}{h_p h_q} \langle k(\cdot, y + h_p u) - k(\cdot, y), k_q(\cdot, x + h_q u) - k(\cdot, x) \rangle_{\mathcal{H}} \\ &= \frac{1}{h_p h_q} (k(x + h_q u, y + h_p u) - k(x, y + h_p u) - k(x + h_q u, y) + k(x, y)) \\ &= \frac{1}{h_p h_q} \left(\partial_u^{(1)} k(x, y + h_p u) \times h_q - \partial_u^{(1)} k(x, y) \times h_q + o(h_q) \right) \\ &= \partial_u^{(2)} \partial_u^{(1)} k(x, y) + o_{p,q}(1) \end{aligned}$$

We have also for some p and q large enough (with $y = x$) and $\varepsilon > 0$,

$$\left| \langle t_p(\cdot, y), t_q(\cdot, y) \rangle_{\mathcal{H}} - \partial_u^{(2)} \partial_u^{(1)} k(y, y) \right| \leq \varepsilon,$$

and by (53) we obtain hence $\|t_p(\cdot, y) - t_q(\cdot, y)\|_{\mathcal{H}}^2 \leq 4\varepsilon$, which means $(t_n(\cdot, y))_n$ is a Cauchy sequence in \mathcal{H} . The sequence $(t_n(\cdot, y))_n$ converges to a function $t(\cdot, y)$ in \mathcal{H} and by (48) the convergence is pointwise. Consequently, we obtain also $\partial_u^{(2)} k(\cdot, y) = t(\cdot, y) \in \mathcal{H}$. \square

As the sequence $(t_n(\cdot, x))_n$ converges to $\partial_u^{(2)} k(\cdot, x)$ in \mathcal{H} , we obtain also

$$\begin{aligned} \left\langle f, \partial_u^{(2)} k(\cdot, x) \right\rangle_{\mathcal{H}} &= \lim_{p \rightarrow \infty} \left\langle f, \frac{k(\cdot, x + h_p) - k(\cdot, x)}{h_p} \right\rangle_{\mathcal{H}} \\ &= \lim_{p \rightarrow \infty} \frac{f(x + h_p u) - f(x)}{h_p} = \partial_u f(x). \end{aligned} \quad (54)$$

We provide in theorem 6 an expression of the directional derivative along direction $u \in E$ of an RKHS function f . To that end we first introduce useful remarks together with an intermediate lemma.

As k is $C^1(E \times E)$, the Mercer eigenfunctions φ_i are $C^1(E, \mathbb{C})$. Similarly to remark 9, we get a computational rule relating the kernel operator derivatives and the inner product $\langle \cdot, \cdot \rangle_{L^2_{\mathbb{C}}(E, \mu)}$. For all $y \in E$ and $u \in E$, we note that

$$\begin{aligned} \langle \partial_u^{(2)} k(\cdot, y), \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)} &= \int_E \partial_u^{(2)} k(x, y) \overline{\varphi_i(x)} \mu(dx) \\ &= \partial_u \int_E \overline{k(y, x)} \overline{\varphi_i(x)} \mu(dx) \\ &= \partial_u \overline{\mathcal{L}_k(\varphi_i)(y)} = \mu_i \partial_u \overline{\varphi_i(y)} \end{aligned}$$

We obtain for all $y \in E$ and $u \in E$ that

$$\langle \partial_u^{(2)} k(\cdot, y), \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)} = \mu_i \partial_u \overline{\varphi_i(y)} \text{ and } \langle \varphi_i, \partial_u^{(2)} k(\cdot, y) \rangle_{L^2_{\mathbb{C}}(E, \mu)} = \mu_i \partial_u \varphi_i(y). \quad (55)$$

These remarks allow us to prove the next lemma.

Lemma 3 *The series $\sum \mu_i |\partial_u \varphi_i(y)|^2$ converges uniformly on E .*

PROOF The functions $\mu_i \varphi_i$ are $C^1(E, \mathbb{C})$. By definition of $\|\cdot\|_{\mathcal{H}}$ and the previous remark, we have

$$\|\partial_u^{(2)} k(\cdot, y)\|_{\mathcal{H}}^2 = \sum_{i=0}^{\infty} \frac{|\langle \partial_u^{(2)} k(\cdot, y), \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)}|^2}{\mu_i} = \sum_{i=0}^{\infty} \mu_i |\partial_u \varphi_i(y)|^2.$$

The series $\sum \mu_i |\partial_u \varphi_i(y)|^2$ converges pointwise to the continuous function $\|\partial_u^{(2)} k(\cdot, y)\|_{\mathcal{H}}^2$. By Dini's theorem (the sequence of partial sums is monotonically increasing as μ_i is positive), the convergence is also uniform. \square

Theorem 6 (Mercer's decomposition of derivatives) *Let $u \in E$. For all $f = \sum_{i=0}^{\infty} a_i \varphi_i \in \mathcal{H}$ (with $a_i = \langle f, \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)}$), the directional derivative of the function f in the direction u exists and we have*

$$\partial_u f = \sum_{i=0}^{\infty} a_i \partial_u(\varphi_i),$$

where the convergence is uniform on E .

PROOF (THEOREM 6) Let $f = \sum_{i=0}^{\infty} a_i \varphi_i \in \mathcal{H}$. For all i , the function $a_i \varphi_i$ belongs to $C^1(E, \mathbb{C})$ and the series $\sum_i a_i \varphi_i$ converges pointwise to f on E . As for the convergence of the derivative series, we have, by Cauchy-Schwarz inequality, that for all $n \in \mathbb{N}$ and $y \in E$,

$$\left| \sum_{i=n}^{\infty} a_i \partial_u(\varphi_i)(y) \right| \leq \left(\sum_{i=n}^{\infty} \frac{|a_i|^2}{\mu_i} \right)^{1/2} \left(\sum_{i=n}^{\infty} \mu_i |\partial_u(\varphi_i)(y)|^2 \right)^{1/2}.$$

By lemma 3, the term $\sum_{i=n}^{\infty} \mu_i |\partial_u(\varphi_i)(y)|^2$ converges uniformly on E to 0. Since $f \in \mathcal{H}$, we obtain also that the series $\sum a_i \partial_u(\varphi_i)$ converges uniformly on E . Then, through term by term differentiation the result is proved. \square

We remark immediately that as functions of the RKHS the kernel feature maps $k(\cdot, x)$ can be differentiated to obtain a term by term Mercer's type decomposition of the kernel derivatives.

Proposition 9 (Mercer Kernel derivatives decomposition) *For all $x, y \in E$ and for all $u, w \in E$, we have*

$$\partial_u^{(1)} \partial_w^{(2)} k(x, y) = \sum_{i=0}^{\infty} \mu_i \partial_u \varphi_i(x) \overline{\partial_w \varphi_i(y)} \quad (56)$$

where the convergence is uniform on $E \times E$.

PROOF The functions $(x, y) \mapsto \mu_i \varphi_i(x) \overline{\varphi_i(y)}$ are $C^2(E \times E)$. For all u and $w \in E$, we apply theorem 6: the series $\sum \mu_i \partial_u \varphi_i(x) \overline{\varphi_i(y)}$ converges uniformly to $\partial_u k(x, y)$ and $\sum \mu_i \varphi_i(x) \overline{\partial_w \varphi_i(y)}$ converges uniformly to $\partial_w k(x, y)$ on $E \times E$. For all $w \in E$, theorem 6 can be applied to the function $x \mapsto \partial_w^{(2)} k(x, y) \in \mathcal{H}$ and for all $u \in E$ and all $x, y \in E$,

$$\partial_u^{(1)} \partial_w^{(2)} k(x, y) = \sum_{i=0}^{\infty} a_{i,w}(y) \partial_u \varphi_i(x),$$

with $a_{i,w}(y) := \langle \partial_w^{(2)} k(\cdot, y), \varphi_i \rangle_{L^2_{\mathbb{C}}(E, \mu)}$. By (55), we obtain that $a_{i,w}(y) = \mu_i \overline{\partial_w \varphi_i(y)}$ and

$$\partial_u^{(1)} \partial_w^{(2)} k(x, y) = \sum_{i=0}^{\infty} \mu_i \partial_u \varphi_i(x) \overline{\partial_w \varphi_i(y)}.$$

Finally, by Cauchy-Schwarz, we have that for all $N \in \mathbb{N}$

$$\left| \sum_{i=0}^N \mu_i \partial_u \varphi_i(x) \overline{\partial_w \varphi_i(y)} \right|^2 \leq \left(\sum_{i=0}^N \mu_i |\partial_u \varphi_i(x)|^2 \right) \left(\sum_{i=0}^N \mu_i |\partial_w \varphi_i(y)|^2 \right),$$

and by lemma 3 the convergence of $\sum \mu_i \partial_u \varphi_i(x) \overline{\partial_w \varphi_i(y)}$ to $\partial_u^{(1)} \partial_w^{(2)} k(x, y)$ is uniform on $E \times E$. \square

These results can be easily extended to higher order derivatives provided a kernel with enough regularity. They correspond, in slightly different forms, to results given in [59, 60] for \mathbb{R}^d . Note that the existence of such higher order derivatives pertains to differentiability conditions of the Mercer eigenfunctions, $(\varphi_i)_{i \in \mathbb{N}}$, associated with kernel k .

By theorem 6, the functions of \mathcal{H} admit a directional derivative along any vector of E . The $L^2_{\mathbb{C}}(E, \mu)$ norm of the directional derivatives of functions of \mathcal{H} can then be controlled by the RKHS norm.

Theorem 7 (Continuity in $L^2_{\mathbb{C}}(E, \mu)$ of RKHS derivatives) *There exists a constant $C > 0$ for which:*

$$\|\partial_u f\|_{L^2_{\mathbb{C}}(E, \mu)} \leq C \|f\|_{\mathcal{H}} \text{ for all } f \in \mathcal{H} \text{ and } u \in E.$$

PROOF Let $u \in E$. We prove the existence of $C > 0$ such that $\|\partial_u f\|_{L^2_{\mathbb{C}}(E, \mu)} \leq C \|f\|_{\mathcal{H}}$ for all $f = \sum_{i=0}^{\infty} a_i \varphi_i \in \mathcal{H}$.

By theorem 6, we have

$$\partial_u f = \sum_{i=0}^{\infty} a_i \partial_u \varphi_i = \sum_{i=0}^{\infty} a_i \sum_{\ell=0}^{\infty} \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \varphi_{\ell} = \sum_{\ell=0}^{\infty} \left(\sum_{i=0}^{\infty} a_i \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right) \varphi_{\ell}.$$

With Cauchy-Schwarz inequality we infer that

$$\begin{aligned} \sum_{\ell=0}^{\infty} \left| \sum_{i=0}^{\infty} a_i \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 &\leq \\ &\sum_{\ell=0}^{\infty} \left(\sum_{i=0}^{\infty} \frac{|a_i|^2}{\mu_i} \right) \left(\sum_{i=0}^{\infty} \mu_i \left| \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 \right) \leq \\ &\|f\|_{\mathcal{H}}^2 \left(\sum_{\ell=0}^{\infty} \sum_{i=0}^{\infty} \mu_i \left| \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 \right). \end{aligned}$$

The last sum is finite as we have, using Fubini's theorem,

$$\begin{aligned} \sum_{\ell=0}^{\infty} \sum_{i=0}^{\infty} \mu_i \left| \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 &= \sum_{i=0}^{\infty} \mu_i \sum_{\ell=0}^{\infty} \left| \langle \partial_u(\varphi_i), \varphi_{\ell} \rangle_{L^2_{\mathbb{C}}(E, \mu)} \right|^2 \leq \\ &\sum_{i=0}^{\infty} \mu_i \int_E |\partial_u(\varphi_i)(x)|^2 \mu(dx). \end{aligned}$$

Upon applying the monotone convergence theorem (since $\mu_i > 0$) and proposition 9 we obtain

$$\begin{aligned} \sum_{i=0}^{\infty} \mu_i \int_E |\partial_u(\varphi_i)(x)|^2 \mu(dx) &= \int_E \sum_{i=0}^{\infty} \mu_i |\partial_u(\varphi_i)(x)|^2 \mu(dx) \\ &= \int_E \partial_u^{(2)} \partial_u^{(1)} k(x, x) \mu(dx), \end{aligned}$$

and $\left(\int_E \partial_u^{(2)} \partial_u^{(1)} k(x,x) \mu(dx)\right)^{1/2} \leq \left(\mu(E) \sup_{x \in E} |\partial^2 k(x,x)| \sup_{u \in E} |u|\right)^{1/2} < \infty$. We denote by C the last constant. \square

A.3 Isometry between $L_{\mathbb{C}}^2(E, \mu)$ and \mathcal{H}

Let us now focus on the links between $L_{\mathbb{C}}^2(E, \mu)$ and \mathcal{H} . Let us recall that integral kernel operator \mathcal{L}_k defined eq.(1) is positive definite, compact and self-adjoint, with a range that is assumed to be dense in $L_{\mathbb{C}}^2(E, \nu)$. As a consequence, it admits a summable non-increasing sequence of strictly positive eigenvalues $(\mu_i)_{i \in \mathbb{N}}$ with corresponding eigenfunctions, $(\varphi_i)_{i \in \mathbb{N}}$ forming an orthonormal basis of $L_{\mathbb{C}}^2(E, \mu)$. Since $\mu_i > 0$ for all integers i , the integral operator \mathcal{L}_k admits a unique symmetric square root operator denoted by $\mathcal{L}_k^{1/2}$ and defined for all integers i as $\mathcal{L}_k^{1/2}(\varphi_i) = \mu_i^{1/2} \varphi_i$. The operator $\mathcal{L}_k^{1/2} : L_{\mathbb{C}}^2(E, \mu) \mapsto \mathcal{H}$ is a bijective isometry: for all $f, g \in L_{\mathbb{C}}^2(E, \mu)$

$$\langle \mathcal{L}_k^{1/2} f, \mathcal{L}_k^{1/2} g \rangle_{\mathcal{H}} = \langle f, g \rangle_{L_{\mathbb{C}}^2(E, \mu)}. \quad (57)$$

This operator must be understood as $\mathcal{L}_k^{1/2} \circ \mathcal{L}_k^{1/2} = \mathcal{L}_k$. The inverse of $\mathcal{L}_k^{1/2}$, noted $\mathcal{L}_k^{-1/2} : \mathcal{H} \mapsto L_{\mathbb{C}}^2(E, \mu)$, and defined for all integers $i \geq 0$ by $\mathcal{L}_k^{-1/2}(\varphi_i) = \mu_i^{-1/2} \varphi_i$ is an isometry as well: for all $f, g \in \mathcal{H}$

$$\langle \mathcal{L}_k^{-1/2} f, \mathcal{L}_k^{-1/2} g \rangle_{L_{\mathbb{C}}^2(E, \mu)} = \langle f, g \rangle_{\mathcal{H}}. \quad (58)$$

It can be checked easily that $\mathcal{L}_k^{1/2}$ and $\mathcal{L}_k^{-1/2}$ are Hermitian for the inner product of $L_{\mathbb{C}}^2(E, \mu)$: for all $f, g \in L_{\mathbb{C}}^2(E, \mu)$

$$\langle \mathcal{L}_k^{1/2}(f), g \rangle_{L_{\mathbb{C}}^2(E, \mu)} = \langle f, \mathcal{L}_k^{1/2}(g) \rangle_{L_{\mathbb{C}}^2(E, \mu)}, \quad (59)$$

and for all $f, g \in \mathcal{H}$

$$\langle \mathcal{L}_k^{-1/2}(f), g \rangle_{L_{\mathbb{C}}^2(E, \mu)} = \langle f, \mathcal{L}_k^{-1/2}(g) \rangle_{L_{\mathbb{C}}^2(E, \mu)}. \quad (60)$$

By construction, the family of Mercer eigenfunctions $(\varphi_i)_{i \in \mathbb{N}}$ is an orthonormal basis of $L_{\mathbb{C}}^2(E, \mu)$ and $\mathcal{L}_k^{1/2}$ enables to get an orthonormal basis of \mathcal{H} .

Proposition 10 (Orthonormal basis of \mathcal{H}) *The sequence of rescaled kernel eigenfunctions $(\mathcal{L}_k^{1/2}(\varphi_i))_{i \geq 0}$ is an orthonormal basis of \mathcal{H} .*

PROOF By (57), the family $(\mathcal{L}_k^{1/2}(\varphi_i))_i$ is orthonormal in \mathcal{H} . Furthermore, we have for all $f \in \mathcal{H}$

$$\sum_{i=0}^{\infty} \left| \langle f, \mathcal{L}_k^{1/2}(\varphi_i) \rangle_{\mathcal{H}} \right|^2 = \sum_{i=0}^{\infty} \left| \langle \mathcal{L}_k^{-1/2} f, \varphi_i \rangle_{L_{\mathbb{C}}^2(E, \mu)} \right|^2 = \|\mathcal{L}_k^{-1/2}(f)\|_{L_{\mathbb{C}}^2(E, \mu)}^2 = \|f\|_{\mathcal{H}}^2,$$

by (60) and (58). We obtain the result through Bessel characterization. \square

The next theorem and proposition provide us useful results for the following

Theorem 8 (Compact injection) *The injection $j : \mathcal{H} \mapsto L_{\mathbb{C}}^2(E, \mu)$ is continuous and compact.*

PROOF The comparison of the norms of \mathcal{H} and $L_{\mathbb{C}}^2(E, \mu)$ and (50) justifies the continuity of j (see remark 12). By proposition 10 and (50), we have

$$\sum_{i=0}^{\infty} \|j(\mathcal{L}_k^{1/2} \varphi_i)\|_{L_{\mathbb{C}}^2(E, \mu)}^2 = \sum_{i=0}^{\infty} \mu_i < \infty;$$

j is also an Hilbert-Schmidt operator and therefore a compact operator. \square

Proposition 11 (Compact injection range denseness) *The image of j , $j(\mathcal{H})$, is dense in $L_{\mathbb{C}}^2(E, \mu)$, more precisely the linear set spanned by $\{k(\cdot, X) : X \in E\}$ is dense in $L_{\mathbb{C}}^2(E, \mu)$.*

PROOF Let $f \in L_{\mathbb{C}}^2(E, \mu)$ such that $\langle f, g \rangle_{L_{\mathbb{C}}^2(E, \mu)} = 0$ for all $g \in \mathcal{H}$. In particular, for all $X \in E$ we have $\langle f, k(\cdot, X) \rangle_{L_{\mathbb{C}}^2(E, \mu)} = 0$, which means that $(\mathcal{L}_k f)(X) = 0$. Since $\mu_i > 0$ for all integers i , we obtain that $f = 0$. \square

From now on, we note $\mathcal{H}^0 := \text{Vect}\{k(\cdot, X) : X \in E\}$. The proposition 11 justifies that \mathcal{H}^0 is dense in $(L_{\mathbb{C}}^2(E, \mu), \|\cdot\|_{L_{\mathbb{C}}^2(E, \mu)})$. Furthermore, by the Moore-Aronszajn theorem [4], \mathcal{H}^0 is also dense in $(\mathcal{H}, \|\cdot\|_{\mathcal{H}})$.

Remark 14 (Higher-order RKHS differentiability) Let $p \geq 1$ be an integer. Assuming $k \in C^{p+1}(E \times E)$, the result of lemma 2 and theorem 6 can be extended to higher order differentiation. As a matter fact, for all $u \in E$, all $x \in E$ and all integer $0 \leq m \leq p$, we can prove that the m th order derivative $\partial_u^{(2), m} k(\cdot, x) \in \mathcal{H}$ and $\mathcal{H} \subset C^p(E, \mathbb{C})$. This corresponds, in slightly different forms, to results provided in [59, 60] for \mathbb{R}^d .

Remark 15 (Relaxation of \mathcal{L}_k range denseness) If some eigenvalues of \mathcal{L}_k are zero, the embedding space $L_{\mathbb{C}}^2(E, \mu)$ we are working with should be replaced by the linear subspace, $\mathcal{V} \subset L_{\mathbb{C}}^2(E, \mu)$ such that $\mathcal{V} = \text{Vect}\{\varphi_i : i \geq 0 \text{ with } \mu_i \neq 0\}$.

Remark 16 (Links between basis of $L_{\mathbb{C}}^2(E, \mu)$ and \mathcal{H}) Proposition 10 proves that if $(e_i)_i$ is an orthonormal basis of $L_{\mathbb{C}}^2(E, \mu)$ then $(\mathcal{L}_k^{1/2} e_i)_i$ is an orthonormal basis of \mathcal{H} . The converse statement is also true and can be shown with the same arguments: if $(f_i)_i$ is an orthonormal basis of \mathcal{H} then $(\mathcal{L}_k^{-1/2} f_i)_i$ is a Hilbert basis of $L_{\mathbb{C}}^2(E, \mu)$.

B Properties of the Extension and Restriction operators E_t and R_t

In this appendix we elaborate on the properties of the restriction and extension operators (namely that they are isometries; form an adjoint pairs in \mathcal{H}_t ; provide a representation of R_t in \mathcal{H} and state the continuity of R_t in $L^2(\Omega_t, \nu)$).

Proposition 12 (Extension/Restriction isometry) *For all $t \geq 0$, the mapping $E_t : \mathcal{H}_t \mapsto \mathcal{H}$ is an isometry :*

$$\forall f, g \in \mathcal{H} \quad \langle E_t(f), E_t(g) \rangle_{\mathcal{H}} = \langle f, g \rangle_{\mathcal{H}_t} \quad (61)$$

and $R_t : \mathcal{H} \mapsto \mathcal{H}_t$ is an isometry as well:

$$\forall f, g \in \mathcal{H}_t \quad \langle R_t(f), R_t(g) \rangle_{\mathcal{H}_t} = \langle f, g \rangle_{\mathcal{H}}. \quad (62)$$

PROOF (FOR THE EXTENSION) Let f and $g \in \text{Vect}\{k_t(\cdot, X_t) : X_t \in \Omega_t\} : f = \sum_{i=1}^{N_1} \alpha_i k_t(\cdot, X_t^{(i)})$ and $g = \sum_{j=1}^{N_2} \beta_j k_t(\cdot, Y_t^{(j)})$ with $X_t^{(i)}$ and $Y_t^{(j)} \in \Omega_t$. By definition, we have

$$\langle E_t(f), E_t(g) \rangle_{\mathcal{H}} = \sum_{i=1}^{N_1} \sum_{j=1}^{N_2} \alpha_i \bar{\beta}_j \langle k(\cdot, X_t^{(i)}), k(\cdot, Y_t^{(j)}) \rangle_{\mathcal{H}} = \sum_{i=1}^{N_1} \sum_{j=1}^{N_2} \alpha_i \bar{\beta}_j k(Y_t^{(j)}, X_t^{(i)}).$$

By construction of k , we have $k(Y_t^{(j)}, X_t^{(i)}) = k_t(Y_t^{(j)}, X_t^{(i)})$ for all i and j , hence

$$\begin{aligned} \langle E_t(f), E_t(g) \rangle_{\mathcal{H}} &= \sum_{i=1}^{N_1} \sum_{j=1}^{N_2} \alpha_i \bar{\beta}_j k_t(Y_t^{(j)}, X_t^{(i)}) \\ &= \sum_{i=1}^{N_1} \sum_{j=1}^{N_2} \alpha_i \bar{\beta}_j \langle k_t(\cdot, X_t^{(i)}), k_t(\cdot, Y_t^{(j)}) \rangle_{\mathcal{H}_t} = \langle f, g \rangle_{\mathcal{H}_t}. \end{aligned}$$

By density of $\text{Vect}\{k_t(\cdot, X_t) : X_t \in \Omega_t\}$ in $(\mathcal{H}_t, \|\cdot\|_{\mathcal{H}_t})$, this result extends to all f and $g \in \mathcal{H}_t$ and (61) is proved. \square

With similar arguments for R_t , (62) is also proved. By (10) and (11), the mappings E_t and R_t have inverse action on the mapping k and k_t . The next proposition justifies that the adjoint of E_t is R_t .

Proposition 13 *For all $t \geq 0$, the mappings E_t and R_t are such that for all $f \in \mathcal{H}_t$ and $g \in \mathcal{H}$*

$$\langle E_t(f), g \rangle_{\mathcal{H}} = \langle f, R_t(g) \rangle_{\mathcal{H}_t}. \quad (63)$$

PROOF For all X_t and $Y_t \in \Omega_t$, we have directly $\langle E_t[k_t(\cdot, X_t)], k(\cdot, Y_t) \rangle_{\mathcal{H}} = \langle k_t(\cdot, X_t), R_t[k(\cdot, Y_t)] \rangle_{\mathcal{H}_t}$. By linearity we obtain that $\langle E_t(f), g \rangle_{\mathcal{H}} = \langle f, R_t(g) \rangle_{\mathcal{H}_t}$ for all $f \in \text{Vect}\{k_t(\cdot, X_t) : X_t \in \Omega_t\}$ and $g \in \text{Vect}\{k(\cdot, X) : X \in \Omega\}$.

Let $f \in \mathcal{H}_t$ and $g \in \mathcal{H}$. There exists sequences $(f_n)_n$ of $\text{Vect}\{k_t(\cdot, X_t) : X_t \in \Omega_t\}$ and $(g_n)_n$ of $\text{Vect}\{k(\cdot, X) : X \in \Omega\}$ such that $\|f - f_n\|_{\mathcal{H}_t} \rightarrow 0$ and $\|g - g_n\|_{\mathcal{H}} \rightarrow 0$. We have for all n the equality

$$\langle E_t(f_n), g_n \rangle_{\mathcal{H}} = \langle f_n, R_t(g_n) \rangle_{\mathcal{H}_t}.$$

Furthermore, by Cauchy-Schwarz and (61) we have that

$$\begin{aligned} |\langle E_t(f_n), g_n \rangle_{\mathcal{H}} - \langle E_t(f), g \rangle_{\mathcal{H}}| &\leq |\langle E_t(f_n) - E_t(f), g \rangle_{\mathcal{H}}| + |\langle E_t(f), g_n - g \rangle_{\mathcal{H}}| \\ &\leq \|f_n - f\|_{\mathcal{H}_t} \|g\|_{\mathcal{H}} + \|f\|_{\mathcal{H}_t} \|g_n - g\|_{\mathcal{H}} \end{aligned}$$

and $\langle E_t(f_n), g_n \rangle_{\mathcal{H}} \rightarrow \langle E_t(f), g \rangle_{\mathcal{H}}$. With similar arguments, it can be readily proved that $\langle f_n, R_t(g_n) \rangle_{\mathcal{H}_t} \rightarrow \langle f, R_t(g) \rangle_{\mathcal{H}_t}$ by using (62). We obtain hence (63). \square

Remark 17 (Expression of $R_t(f)$ for $f \in \mathcal{H}$) The function $R_t(f)$ belongs to \mathcal{H}_t . By proposition 13, we have for all $X_t \in \Omega_t$

$$R_t(f)(X_t) = \langle R_t(f), k_t(\cdot, X_t) \rangle_{\mathcal{H}_t} = \langle f, k(\cdot, X_t) \rangle_{\mathcal{H}} = f(X_t).$$

More generally, for all $t \geq 0$ the mapping R_t can be extended on $L_{\mathbb{C}}^2(\Omega, \nu)$ (keeping the same notation) as:

$$R_t : \begin{array}{ccc} L_{\mathbb{C}}^2(\Omega, \nu) & \mapsto & L_{\mathbb{C}}^2(\Omega_t, \nu) \\ f & \mapsto & f \upharpoonright_{\Omega_t} \end{array}. \quad (64)$$

In particular, on $\mathcal{H} \subset L_{\mathbb{C}}^2(\Omega, \nu)$ this boils down to the definition given in (11). As $\Omega_t \subset \Omega$, we have directly the next proposition.

Proposition 14 (Continuity of the restriction) *The mapping $R_t : (L_{\mathbb{C}}^2(\Omega, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega, \nu)}) \mapsto (L_{\mathbb{C}}^2(\Omega_t, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega_t, \nu)})$ is continuous.*

C Proof of lemma 1 (Perron-Frobenius infinitesimal generator)

In this appendix we provide the proof on the Perron-Frobenius infinitesimal generator expression and of its domain. We recall that \mathcal{H}^0 is the linear span of the set $\{k(\cdot, x) : x \in \Omega\}$.

PROOF Let $f \in \mathcal{D}(A_p)$ and $\psi \in \mathcal{H}^0$, we have for all $t \geq 0$,

$$\left\langle \frac{P_t \circ f - f}{t}, \psi \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \left\langle f, \frac{U_t \circ \psi - \psi}{t} \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \quad (65)$$

For all $x \in \Omega$, we have

$$\frac{U_t(\psi)(x) - \psi(x)}{t} \xrightarrow{t \rightarrow 0^+} \frac{d}{dt} [U_t(\psi)(x)] \upharpoonright_{t=0} = \partial_{\mathcal{M}(x)} \psi(x).$$

D PROOF OF STEP1 AND STEP2 OF WONDERLAND SPECTRAL THEOREM46

We pass to the limit $[t \rightarrow 0^+]$ in (65), and recalling that \mathcal{M} is real valued, divergence free with $\mathcal{M} \upharpoonright_{\partial\Omega} = 0$, and measure invariant we obtain

$$\int_{\Omega} A_p f(x) \overline{\psi(x)} \nu(dx) = \int_{\Omega} -\partial_{\mathcal{M}(x)} f(x) \overline{\psi(x)} \nu(dx)$$

thus, $x \mapsto \partial_{\mathcal{M}(x)} f(x)$ belongs to $L^2_{\mathbb{C}}(\Omega, \nu)$ and

$$\mathcal{D}(A_p) \subset \{f \in L^2_{\mathbb{C}}(\Omega, \nu) : x \mapsto \partial_{\mathcal{M}(x)} f(x) \in L^2_{\mathbb{C}}(\Omega, \nu)\}$$

$$\text{and } A_p f(x) = -\partial_{\mathcal{M}(x)} f(x) \quad \text{for } f \in \mathcal{D}(A_p) \text{ and } x \in \Omega.$$

Conversely, considering now $f \in L^2_{\mathbb{C}}(\Omega, \nu)$ such that $x \mapsto \partial_{\mathcal{M}(x)} f(x) \in L^2_{\mathbb{C}}(\Omega, \nu)$, we show that $f \in \mathcal{D}(A_p)$. For all $\psi \in \mathcal{H}^0$, we have

$$\begin{aligned} \langle P_t(f) - f, \psi \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} &= \langle f, U_t(\psi) - \psi \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \\ &= \langle f, \int_0^t \frac{dU_s(\psi)}{ds} ds \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle f, \int_0^t \partial_{\mathcal{M}(\phi_s(\cdot))} \psi[\Phi_s(\cdot)] ds \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}. \end{aligned}$$

For all $\psi \in \mathcal{H}^0$, we have $\langle -\partial_{\mathcal{M}(\cdot)} f(\cdot), \psi \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle f, -\partial_{\mathcal{M}(\cdot)} \psi(\cdot) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}$ (since \mathcal{M} is divergence free) and we obtain also the following equality

$$\begin{aligned} \left\langle \frac{P_t(f) - f}{t} - \partial_{\mathcal{M}(\cdot)} f(\cdot), \psi \right\rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} &= \\ &= \left\langle f, \frac{1}{t} \int_0^t \partial_{\mathcal{M}(\phi_s(\cdot))} \psi[\Phi_s(\cdot)] ds - \partial_{\mathcal{M}(\cdot)} \psi(\cdot) \right\rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}. \end{aligned}$$

Since $t \mapsto \int_0^t \partial_{\mathcal{M}(\phi_s(\cdot))} \psi[\Phi_s(\cdot)] ds$ is differentiable in 0^+ , we obtain in particular

$$\left\langle \frac{P_t(f) - f}{t} - \partial_{\mathcal{M}(\cdot)} f(\cdot), \psi \right\rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \xrightarrow{t \rightarrow 0^+} 0,$$

for all function $\psi \in \mathcal{H}^0$. By density of \mathcal{H}^0 in $L^2_{\mathbb{C}}(\Omega, \nu)$ we get

$$\left\| \frac{P_t f - f}{t} - \partial_{\mathcal{M}(\cdot)} f(\cdot) \right\|_{L^2_{\mathbb{C}}(\Omega, \nu)} \xrightarrow{t \rightarrow 0^+} 0$$

and hence $f \in \mathcal{D}(A_p)$. □

D Proof of step1 and step2 of Wonderland spectral theorem

This appendix gathers step 1 and step 2 of the proof of the Wonderland representation theorem. Step 1 Uses Lions-Lax-Milgram theorem to show that an approximation \tilde{A}_U of the infinitesimal generator A_U is bijective. The inverse operator

D PROOF OF STEP1 AND STEP2 OF WONDERLAND SPECTRAL THEOREM47

\tilde{A}_U^{-1} seen as an operator of $L_{\mathbb{C}}^2(\Omega, \nu) \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ through the composition with the compact injection $j : \mathcal{H} \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ (Theorem 8, A.3) is shown to be compact and self-adjoint (Lemmas 5 and 6). The diagonalization of A_U is then obtained from the diagonalization of \tilde{A}_U^{-1} and \mathcal{L}_k (Prop.15). Step 2 finally constructs from the eigenfunctions of A_U an orthonormal basis of \mathcal{H}_t that diagonalizes the operator $A_{U,t}$ (Prop.16).

Step 1

Let \tilde{A}_U be the linear mapping defined by

$$\begin{aligned} \tilde{A}_U : (\mathcal{H}, \|\cdot\|_{\mathcal{H}}) &\mapsto (L_{\mathbb{C}}^2(\Omega, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega, \nu)}) \\ f &\mapsto [iA_U + \lambda \mathcal{L}_k^{-1/2}](f) \end{aligned} \quad (66)$$

with λ a fixed real such that

$$|\lambda| > \|A_U\| \left(\int_{\Omega} |k(x,x)|^2 \nu(dx) \right)^{1/2}.$$

By Theorem 7 (A.2), the norm of A_U verifies

$$\|A_U\| \leq \left(\nu(E) \sup_{x \in E} \|\partial^2 k(x,x)\| \sup_{x \in E} |\mathcal{M}(x)| \right)^{1/2},$$

and to lighten the notations in the following, we note $M := \|A_U\|$. We start analyzing the properties of \tilde{A}_U . First of all, the mapping \tilde{A}_U is continuous. We have for all $f \in \mathcal{H}$ the inequality

$$\begin{aligned} \|\tilde{A}_U f\|_{L_{\mathbb{C}}^2(\Omega, \nu)} &\leq \|iA_U f\|_{L_{\mathbb{C}}^2(\Omega, \nu)} + |\lambda| \|\mathcal{L}_k^{-1/2} f\|_{L_{\mathbb{C}}^2(\Omega, \nu)} = \\ &\|A_U f\|_{L_{\mathbb{C}}^2(\Omega, \nu)} + |\lambda| \|\mathcal{L}_k^{-1/2} f\|_{L_{\mathbb{C}}^2(\Omega, \nu)}. \end{aligned}$$

The identity of (58) and theorem 2 allow us to write

$$\|\tilde{A}_U f\|_{L_{\mathbb{C}}^2(\Omega, \nu)} \leq (M + |\lambda|) \|f\|_{\mathcal{H}}.$$

Lemma 4 (Bijectivity of \tilde{A}_U) *The operator $\tilde{A}_U : (\mathcal{H}, \|\cdot\|_{\mathcal{H}}) \mapsto (L_{\mathbb{C}}^2(\Omega, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega, \nu)})$ is bijective.*

PROOF Let a be the bilinear form defined by

$$\begin{aligned} a : \mathcal{H} \times \mathcal{H} &\mapsto \mathbb{C} \\ (f, g) &\mapsto \langle \tilde{A}_U(f), \mathcal{L}_k^{-1/2}(g) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}. \end{aligned}$$

The bilinear form a is continuous on \mathcal{H} . By (58), for all $f \in \mathcal{H}$ we have

$$a(f, f) = \langle iA_U(f), \mathcal{L}_k^{-1/2} f \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} + \lambda \|f\|_{\mathcal{H}}^2.$$

D PROOF OF STEP1 AND STEP2 OF WONDERLAND SPECTRAL THEOREM48

In particular, we get

$$|a(f, f)| \geq \left| |\lambda| \|f\|_{\mathcal{H}}^2 - |\langle A_U(f), \mathcal{L}_k^{-1/2} f \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}| \right| \geq \\ |\lambda| \|f\|_{\mathcal{H}}^2 - |\langle A_U(f), \mathcal{L}_k^{-1/2} f \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}|.$$

Using Cauchy-Schwarz inequality we have

$$-|\langle A_U(f), \mathcal{L}_k^{-1/2} f \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}| \geq -\|A_U(f)\|_{L_{\mathbb{C}}^2(\Omega, \nu)} \|\mathcal{L}_k^{-1/2} f\|_{L_{\mathbb{C}}^2(\Omega, \nu)}.$$

Through the continuity of $A_U : \mathcal{H} \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$, remark 12 and (58) we infer that

$$-|\langle A_U(f), \mathcal{L}_k^{-1/2} f \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}| \geq -M \left(\int_{\Omega} |k(x, x)|^2 \nu(dx) \right)^{1/2} \|f\|_{\mathcal{H}}^2.$$

We obtain hence that $|a(f, f)| \geq \gamma \|f\|_{\mathcal{H}}^2$ for all $f \in \mathcal{H}$ with the constant

$$\gamma := \left(|\lambda| - M \left(\int_{\Omega} |k(x, x)|^2 \nu(dx) \right)^{1/2} \right) > 0,$$

which is positive by definition of λ . The bilinear form a verifies thus a coercivity condition in \mathcal{H} .

To show that $\tilde{A}_U : \mathcal{H} \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ is bijective, let us consider $f \in L_{\mathbb{C}}^2(\Omega, \nu)$. The linear mapping $g \mapsto \langle f, \mathcal{L}_k^{-1/2}(g) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}$ is continuous on \mathcal{H} . By the Lions–Lax–Milgram theorem, there exists thus a unique $x_f \in \mathcal{H}$ such that

$$a(x_f, g) = \langle f, \mathcal{L}_k^{-1/2}(g) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \quad \text{for all } g \in \mathcal{H}.$$

As the operator $\mathcal{L}_k^{-1/2} : \mathcal{H} \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ is bijective, we obtain also

$$\langle \tilde{A}_U(x_f), h \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle f, h \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \quad \text{for all } h \in L_{\mathbb{C}}^2(\Omega, \nu). \quad (67)$$

In particular, we obtain hence $\tilde{A}_U(x_f) = f$. □

This lemma justifies the existence of the operator $\tilde{A}_U^{-1} : (L_{\mathbb{C}}^2(\Omega, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega, \nu)}) \mapsto (\mathcal{H}, \|\cdot\|_{\mathcal{H}})$. However, in order to diagonalize A_U , we will need to diagonalize the inverse of A_U in $L_{\mathbb{C}}^2(\Omega, \nu)$. To that end, we consider $j \circ \tilde{A}_U^{-1} : L_{\mathbb{C}}^2(\Omega, \nu) \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ with j the injection of \mathcal{H} in $L_{\mathbb{C}}^2(\Omega, \nu)$. As the injection is compact (Theorem 8), A.3), we obtain directly that the mapping

$$j \circ \tilde{A}_U^{-1} : (L_{\mathbb{C}}^2(\Omega, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega, \nu)}) \mapsto (L_{\mathbb{C}}^2(\Omega, \nu), \|\cdot\|_{L_{\mathbb{C}}^2(\Omega, \nu)})$$

is continuous and compact. By abuse of notation denoting the composition $j \circ \tilde{A}_U^{-1}$ as $\tilde{A}_U^{-1} : L_{\mathbb{C}}^2(\Omega, \nu) \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ we have the next lemma .

Lemma 5 (Compactness of \tilde{A}_U^{-1}) *The operator $\tilde{A}_U^{-1} : L_{\mathbb{C}}^2(\Omega, \nu) \mapsto L_{\mathbb{C}}^2(\Omega, \nu)$ is compact.*

Lemma 6 (Self-adjointness \tilde{A}_U^{-1}) *The operator \tilde{A}_U^{-1} is self-adjoint in $L^2_{\mathbb{C}}(\Omega, \nu)$.*

PROOF For all f and $g \in L^2_{\mathbb{C}}(\Omega, \nu)$, we have

$$\langle \tilde{A}_U^{-1}(f), g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle x_f, g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle x_f, \tilde{A}_U x_g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \quad (68)$$

where we used the notation of the proof of lemma 4 (and the elements x_f and x_g belong to \mathcal{H}). By definition, we have

$$\langle x_f, \tilde{A}_U x_g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = -i \langle x_f, A_U(x_g) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} + \lambda \langle x_f, \mathcal{L}_k^{-1/2}(x_g) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}.$$

Skew symmetry of the generator A_U (Proposition 4), symmetry of $\mathcal{L}_k^{-1/2}$ (60) and as λ is real allow us to write

$$\begin{aligned} \langle x_f, \tilde{A}_U x_g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} &= i \langle A_U(x_f), x_g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} + \lambda \langle \mathcal{L}_k^{-1/2}(x_f), x_g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} \\ &= \langle \tilde{A}_U x_f, x_g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}. \end{aligned}$$

With similar arguments as (68), we conclude that for all f and $g \in L^2_{\mathbb{C}}(\Omega, \nu)$ $\langle \tilde{A}_U^{-1}(f), g \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} = \langle f, \tilde{A}_U^{-1}(g) \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)}$. \square

With lemmas 5 and 6, we may apply the spectral theorem of compact self-adjoint operator and consequently there exists a orthonormal basis $(e_n)_{n \geq 0}$ of $(L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$ and a sequence of real $(\alpha_n)_n \in (\sigma_p(\tilde{A}_U^{-1}) - \{0\})^{\mathbb{N}}$ such that

$$\lim_{n \rightarrow \infty} \alpha_n = 0 \quad \text{and} \quad \tilde{A}_U^{-1}(e_n) = \alpha_n e_n \quad \text{for all } n \in \mathbb{N}. \quad (69)$$

Moreover, for all integers n , the real eigenvalue α_n has finite multiplicity. It can be noticed that $\alpha_n \neq 0$ for all n since \tilde{A}_U^{-1} is invertible and the eigenfunctions e_n belong to the RKHS \mathcal{H} . Finally, we have for all $n \in \mathbb{N}$

$$\tilde{A}_U(e_n) = [iA_U + \lambda \mathcal{L}_k^{-1/2}](e_n) = \alpha_n^{-1} e_n.$$

We now are in position to fully specify the eigenvalues of A_U . If all the eigenvalues of \tilde{A}_U^{-1} are simple, it is easier to relate them to the eigenvalues of A_U . This is explained in remark 18. In the other case, we need first to establish a commutativity property between \tilde{A}_U^{-1} and \mathcal{L}_k .

Lemma 7 (Commutation between A_U and \mathcal{L}_k) *The operators A_U and \mathcal{L}_k are commutative on \mathcal{H} . For all $f \in \mathcal{H}$ we have the identity: $A_U \circ \mathcal{L}_k(f) = \mathcal{L}_k \circ A_U(f)$.*

Before demonstrating this result, we first show a useful equality on \mathcal{H}^0 . Considering $f \in \mathcal{H}^0$ the element $\mathcal{L}_k(f)$ belongs to \mathcal{H} . We apply proposition 2 and we have for all $X \in \Omega$ and all $t \geq 0$

$$\mathcal{L}_k[U_t(f)](X) = \int_{\Omega} k(X, y) U_t(f)(y) \nu(dy) = \int_{\Omega} k(\Phi_t(X), \Phi_t(y)) f \circ \Phi_t(y) \nu(dy).$$

D PROOF OF STEP1 AND STEP2 OF WONDERLAND SPECTRAL THEOREM50

Since the dynamical system $(\Phi_t)_{t \geq 0}$ is measure preserving and $\mathcal{L}_k(f)$ belongs to \mathcal{H} , we obtain

$$\mathcal{L}_k[U_t(f)](X) = \int_{\Omega} k(\Phi_t(X), y) f(y) \nu(dy) = [\mathcal{L}_k f](\Phi_t(X)).$$

PROOF (LEMMA 7) First, we prove the result for $f \in \mathcal{H}^0$.

By the previous equality we have for all $g \in L_{\mathbb{C}}^2(\Omega, \nu)$ and $t \geq 0$

$$\left\langle \frac{[\mathcal{L}_k f](\Phi_t(\cdot)) - \mathcal{L}_k f}{t}, g \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \left\langle \frac{\mathcal{L}_k[U_t f] - \mathcal{L}_k f}{t}, g \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}.$$

The operator \mathcal{L}_k being self-adjoint on $L_{\mathbb{C}}^2(\Omega, \nu)$ we infer that

$$\left\langle \frac{[\mathcal{L}_k f](\Phi_t(\cdot)) - \mathcal{L}_k f}{t}, g \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \left\langle \frac{U_t f - f}{t}, \mathcal{L}_k(g) \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}.$$

Passing to the limit $[t \rightarrow 0^+]$, as $\mathcal{L}_k(f)$ belongs to $\mathcal{H} \subset \mathcal{D}(A_U)$, we have

$$\begin{aligned} \lim_{t \rightarrow 0^+} \left\langle \frac{[\mathcal{L}_k f](\Phi_t(\cdot)) - \mathcal{L}_k f}{t}, g \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} &= \left\langle \frac{d}{dt} [(\mathcal{L}_k f)(\Phi_t(\cdot))]_{t=0}, g \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \\ &= \langle \partial_{\mathcal{H}}(\cdot) \mathcal{L}_k(f)(\cdot), g \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \\ &= \langle A_U \mathcal{L}_k(f), g \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} \end{aligned}$$

and f belong to $\mathcal{H} \subset \mathcal{D}(A_U) = \mathcal{D}(A_P)$, we have too

$$\lim_{t \rightarrow 0^+} \left\langle \frac{U_t f - f}{t}, \mathcal{L}_k(g) \right\rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle A_U f, \mathcal{L}_k(g) \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle \mathcal{L}_k \circ A_U f, g \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}.$$

We deduce that $\langle A_U \circ \mathcal{L}_k(f), g \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)} = \langle \mathcal{L}_k \circ A_U(f), g \rangle_{L_{\mathbb{C}}^2(\Omega, \nu)}$ for all $g \in L_{\mathbb{C}}^2(\Omega, \nu)$.

The lemma is proved on \mathcal{H}^0 .

Then, we suppose that $f \in \mathcal{H}$. There exists a sequence $(f_n)_n$ of \mathcal{H}^0 which converges to f in \mathcal{H} .

The continuity of $A_U : \mathcal{H} \rightarrow L_{\mathbb{C}}^2(E, \mu)$ and remark 13 justifies for all integers n

$$\|A_U \circ \mathcal{L}_k(f_n) - A_U \circ \mathcal{L}_k(f)\|_{L_{\mathbb{C}}^2(E, \mu)} \leq C \|\mathcal{L}_k(f_n - f)\|_{\mathcal{H}} \leq C \|f_n - f\|_{\mathcal{H}}.$$

As $L_k : L_{\mathbb{C}}^2(E, \mu) \rightarrow L_{\mathbb{C}}^2(E, \mu)$ is continuous, we infer

$$\|\mathcal{L}_k \circ A_U(f_n) - \mathcal{L}_k \circ A_U(f)\|_{L_{\mathbb{C}}^2(E, \mu)} \leq C \|A_U(f_n - f)\|_{L_{\mathbb{C}}^2(E, \mu)} \leq C \|f_n - f\|_{\mathcal{H}}.$$

For all integers n , we have $A_U \circ \mathcal{L}_k(f_n) = \mathcal{L}_k \circ A_U(f_n)$. We pass to the limit to obtain $A_U \circ \mathcal{L}_k(f) = \mathcal{L}_k \circ A_U(f)$. \square

Lemma 8 (Commutation between \tilde{A}_U^{-1} and \mathcal{L}_k .) *The operators \tilde{A}_U^{-1} and \mathcal{L}_k are commutative on $L_{\mathbb{C}}^2(\Omega, \nu)$.*

PROOF We apply the lemma 7 and we infer for all $f \in \mathcal{H}$

$$\begin{aligned}\mathcal{L}_k \circ \tilde{A}_U(f) &= i\mathcal{L}_k \circ A_U(f) + \lambda \mathcal{L}_k \circ \mathcal{L}_k^{-1/2}(f) \\ &= iA_U \circ \mathcal{L}_k(f) + \lambda \mathcal{L}_k^{-1/2} \circ \mathcal{L}_k(f) = \tilde{A}_U \circ \mathcal{L}_k(f).\end{aligned}$$

As $\tilde{A}_U : \mathcal{H} \mapsto L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$ is bijective, we obtain that $\mathcal{L}_k \circ \tilde{A}_U^{-1}(g) = \tilde{A}_U^{-1} \circ \mathcal{L}_k(g)$ for all $g \in L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$.

Proposition 15 (Diagonalization of \tilde{A}_U^{-1} and \mathcal{L}_k) *The operators \tilde{A}_U^{-1} and \mathcal{L}_k are diagonalizable in the same orthonormal basis of $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$.*

PROOF For $n \in \mathbb{N}$, we note $E_{\alpha_n} \subset L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$ the eigenspace of \tilde{A}_U^{-1} associated with eigenvalue α_n . By lemma 8, we have for all $f \in E_{\alpha_n}$ that

$$[\tilde{A}_U^{-1} - \alpha_n I] \circ \mathcal{L}_k(f) = \mathcal{L}_k \circ [\tilde{A}_U^{-1} - \alpha_n I](f) = 0,$$

and E_{α_n} is thus an invariant subspace of \mathcal{L}_k . The operator $\mathcal{L}_k : E_{\alpha_n} \mapsto E_{\alpha_n}$ can be diagonalized. There exists an orthonormal basis $(f_{n,p})_p$ of E_{α_n} and reals $(\beta_{n,p})_p$ (with $1 \leq p \leq \dim E_{\alpha_n}$) such that

$$\mathcal{L}_k(f_{n,p}) = \beta_{n,p} f_{n,p} \quad \text{and} \quad \tilde{A}_U^{-1}(f_{n,p}) = \alpha_n f_{n,p}.$$

The family $(f_{n,p})_p$ diagonalizes both the operators \mathcal{L}_k and \tilde{A}_U^{-1} on E_{α_n} . It can be noticed that $\beta_{n,p} > 0$ since \mathcal{L}_k is positive definite. As $\alpha_n \neq 0$ for all n , the functions $f_{n,p}$ belong to \mathcal{H} . The concatenation of the vectors $(f_{n,p})_{n,p}$ for all $n \in \mathbb{N}$ and $1 \leq p \leq \dim E_{\alpha_n}$ forms a orthonormal basis of $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$. As a matter of fact, we have by construction that, $\text{Vect}\{f_{n,p} : n \in \mathbb{N} \text{ and } 1 \leq p \leq \dim E_{\alpha_n}\} = \text{Vect}\{e_n : n \in \mathbb{N}\}$ and the family $(f_{n,p})_{n,p}$ is also complete in $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$. For all n the family $(f_{n,p})_p$ is orthonormal in $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$ (by construction). The eigenspaces of \tilde{A}_U^{-1} are in addition orthogonal: the vectors $f_{n,p}$ and $f_{m,q}$ are orthogonal in $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$ for all $n \neq m$, $1 \leq p \leq \dim E_{\alpha_n}$ and $1 \leq q \leq \dim E_{\alpha_m}$. The family $(f_{n,p})_{n \in \mathbb{N}, 1 \leq p \leq \dim E_{\alpha_n}}$ is thus orthonormal in $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$ and constitutes a orthonormal basis of $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$. \square

The result of step 1 can now be obtained. Keeping the same notations for the eigenfunctions of \tilde{A}_U , for all $n \in \mathbb{N}$ and $1 \leq p \leq \dim E_{\alpha_n}$, we have $\tilde{A}_U(f_{n,p}) = \alpha_n^{-1} f_{n,p}$ and $\mathcal{L}_k^{-1/2}(f_{n,p}) = \beta_{n,p}^{-1/2} f_{n,p}$. With the expression of A_U in (66) we obtain

$$A_U(f_{n,p}) = i(\lambda \beta_{n,p}^{-1/2} - \alpha_n^{-1}) f_{n,p}. \quad (70)$$

Denoting for the sake of clarity, $(V_\ell)_\ell$ the orthonormal basis $(f_{n,p})_{n,p}$ of $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$ and $(\lambda_\ell)_\ell$ the associated eigenvalues of A_U (given (70) with $\lambda_\ell \in i\mathbb{R}$) and upon applying $\mathcal{L}_k^{1/2}$ on the orthonormal basis $(V_\ell)_{\ell \in \mathbb{N}}$ of $L_{\mathbb{C}}^2(\Omega, \mathfrak{v})$, we get also a orthonormal basis of \mathcal{H} (remark 16) which diagonalizes A_U :

$$A_U(\mathcal{L}_k^{1/2} V_\ell) = \lambda_\ell \mathcal{L}_k^{1/2} V_\ell. \quad (71)$$

This completes the proof of step 1.

It can be noticed that for all $g \in \mathcal{H}$ we have

$$A_U(g) = \sum_{\ell=0}^{\infty} \lambda_{\ell} \langle g, V_{\ell} \rangle_{L^2_{\mathbb{C}}(\Omega, \nu)} V_{\ell} \quad \text{and} \quad A_U(g) = \sum_{\ell=0}^{\infty} \lambda_{\ell} \langle g, \mathcal{L}_k^{1/2}(V_{\ell}) \rangle_{\mathcal{H}} \mathcal{L}_k^{1/2}(V_{\ell}).$$

Remark 18 (Particular case: \tilde{A}_U has simple eigenvalues) A simpler case arises when the eigenvalues of \tilde{A}_U are simple. This corresponds to the case of ergodic dynamical systems [30]. For all n , by (69) and lemma 7, the eigenspace E_{α_n} of \tilde{A}_U is an invariant space of $\mathcal{L}_k^{-1/2}$. As $\dim E_{\alpha_n} = 1$ and E_{α_n} is an invariant subspace of $\mathcal{L}_k^{-1/2}$, there exists reals β_n such that $\mathcal{L}_k^{-1/2} e_n = \beta_n e_n$ and we obtain

$$A_U(e_n) = i(\lambda \beta_n - \alpha_n^{-1}) e_n,$$

with the orthonormal basis $(e_n)_n$ of $L^2_{\mathbb{C}}(\Omega, \nu)$ obtained from the diagonalization of \tilde{A}_U^{-1} . We have as well for the associated orthonormal basis of \mathcal{H}

$$A_U(\mathcal{L}_k^{1/2} e_n) = i(\lambda \beta_n - \alpha_n^{-1}) (\mathcal{L}_k^{1/2} e_n).$$

Remark 19 (Exponential form of the Koopman semi-group) The range of \tilde{A}_U^{-1} is included in \mathcal{H} . It can be also noticed that V_{ℓ} belongs to \mathcal{H} and the following inclusions hold

$$\text{Vect}\{V_{\ell} : \ell \in \mathbb{N}\} \subset \mathcal{H} \subset L^2_{\mathbb{C}}(\Omega, \nu).$$

The set $\text{Vect}\{V_{\ell} : \ell \in \mathbb{N}\}$ is dense in $(L^2_{\mathbb{C}}(\Omega, \nu), \|\cdot\|_{L^2_{\mathbb{C}}(\Omega, \nu)})$ and is an invariant space of the Koopman infinitesimal generator A_U . For all f belonging to this set and all $t \geq 0$, as A_U is bounded, the Koopman operator is uniformly continuous and can be written in terms of an exponential series as

$$U_t(f) = \exp(tA_U)(f),$$

with convergence in the operator norm. Its adjoint, the Perron-Frobenius operator, can be written in the same way as

$$P_t(f) = \exp(-tA_U)(f).$$

Denoting for the sake of simplicity, $\psi_{\ell} = \mathcal{L}_k^{1/2}(V_{\ell})$ for all integers ℓ . The family $(\psi_{\ell})_{\ell}$ is a orthonormal basis of \mathcal{H} . We have $U_t(\psi_{\ell}) = \exp(t \lambda_{\ell}) \psi_{\ell}$ and we obtain hence

$$\psi_{\ell}(X_t) = \exp(t \lambda_{\ell}) \psi_{\ell}(X_0). \tag{72}$$

Step 2

From this orthonormal basis $(\psi_{\ell})_{\ell}$ of \mathcal{H} , we now build a orthonormal basis of \mathcal{H}_t that diagonalizes the operator $A_{U,t}$ for all time $t \geq 0$.

Proposition 16 (Hilbert basis of \mathcal{H}_t) For all $t \geq 0$, the family $(R_t(\psi_\ell))_\ell$ is an orthonormal basis of $(\mathcal{H}_t, \|\cdot\|_{\mathcal{H}_t})$.

PROOF For all ℓ and ℓ' , upon applying proposition 12 we have $\langle R_t(\psi_\ell), R_t(\psi_{\ell'}) \rangle_{\mathcal{H}_t} = \langle \psi_\ell, \psi_{\ell'} \rangle_{\mathcal{H}} = \delta_{\ell, \ell'}$, consequently the family $(R_t(\psi_\ell))_\ell$ is orthonormal in \mathcal{H}_t . Besides, the family $\text{Vect}\{R_t(\psi_\ell) : \ell \in \mathbb{N}\}$ is dense in \mathcal{H}_t . Let $h \in \mathcal{H}_t$ be such that $\langle h, R_t(\psi_\ell) \rangle_{\mathcal{H}_t} = 0$ for all integers ℓ . By proposition 13, we have $\langle h, R_t(\psi_\ell) \rangle_{\mathcal{H}_t} = \langle E_t(h), \psi_\ell \rangle_{\mathcal{H}} = 0$. As $(\psi_\ell)_\ell$ is an orthonormal basis of \mathcal{H} , we conclude that $E_t(h) = 0$ and thus $h = 0$ on \mathcal{H}_t . \square

The orthonormal basis $(R_t(\psi_\ell))_\ell$ of \mathcal{H}_t diagonalizes the operator $A_{U,t}$. As a matter of fact, for all ℓ ,

$$A_{U,t}[R_t\psi_\ell] = R_t \circ A_U \circ E_t \circ R_t(\psi_\ell) = \lambda_\ell [R_t\psi_\ell].$$

For conciseness purpose, we note in the following $\psi_\ell^t := R_t(\psi_\ell)$. Through remark 17 and (72), it can be noticed that for all $X_0 \in \Omega_0$

$$\psi_\ell^t(X_t) = \exp(t\lambda_\ell) \psi_\ell^0(X_0),$$

which ends the proof of step 2, and completes the proof of the spectral representation of Koopman operator in Wonderland.

Remark 20 The set $\text{Vect}\{\psi_\ell^t : \ell \in \mathbb{N}\}$ is an invariant space of $A_{U,t}$. As a matter of fact, considering remark 19 and proposition 12, for all $f \in \text{Vect}\{\psi_\ell^t : \ell \in \mathbb{N}\}$ we have $\exp(tA_{U,t})(f) = R_t \circ \exp(tA_U) \circ E_t(f)$.

E Numerical details of the QG implementation

The QG model is discretized on a staggered grid of size $N_x = 64 \times N_y = 128$ with order-2 finite differences. Advection term is computed using the Arakawa discretization [2] and centered differences are considered for the other terms. The Poisson equation is solved using spectral method by projection on a sine basis. Time integration is performed with a fourth-order Runge-Kutta scheme. An ensemble of $p = 100$ members is considered for the training, and another test ensemble of $p_2 = 100$ members is generated for predictions assessment.

Starting from an established state of a particular trajectory, a set of initial conditions is generated by perturbing this state by the potential vorticity of $N_{POD} = 50$ proper orthogonal decomposition (POD) modes [44] weighted by a Gaussian random perturbation $\mathcal{N}(0, 10\lambda_i/\lambda_1)$, where λ_i is the i^{th} POD eigenvalue. Then, a spin-up time integration of $t = 0.03$ is performed to evacuate eventual nonphysical flow features. The training and testing sets are generated independently using the same rule.

Following the dimensioning given in [56], $R_o = \frac{U_r}{\beta L_r^2} = 0.0036$ with U_r and L_r the reference velocity and length scale, respectively. We give as well the gradient of

the Coriolis parameter $\beta = 1.75 \cdot 10^{-11}$, typical of mid-latitudes at the center of the basin. Setting $L_r = 2000\text{km}$, leads to $U_r = 0.252\text{m}\cdot\text{s}^{-1}$, and then to a reference time $T_r = \frac{L_r}{U_r} = 92\text{days}$.

References

- [1] R. Alexander and D. Giannakis. Operator-theoretic framework for forecasting nonlinear time series with kernel analog techniques. *Physica D: Nonlinear Phenomena*, 409:132520, 2020.
- [2] A. Arakawa. Computational design for long-term numerical integration of the equations of fluid motion: Two-dimensional incompressible flow. Part I. *Journal of Computational Physics*, 1(1):119–143, 1966.
- [3] H. Arbabian and I. Mezic. Ergodic theory, dynamic mode decomposition and computation of spectral properties of the Koopman operator. *SIAM Journal on Applied Dynamical Systems*, 16(4):2096–2126, 2017.
- [4] N. Aronszajn. Theory of reproducing kernels. *Trans. of the American Math. Society*, 68(337-404), 1950.
- [5] AVISO. Ssalto/duacs user handbook: Msla and (m)adt near-real time and delayed time products. Technical Report CLS-DOS-NT-06-034, SALP-MU-P-EA-21065-CLS, 5rev0, CNES, 2016. available online at https://www.aviso.altimetry.fr/fileadmin/documents/data/tools/hdbk_duacs.pdf.
- [6] P. J. Baddoo, B. Herrmann, B. J. McKeon, and S. L. Brunton. Kernel learning for robust dynamic mode decomposition: linear and nonlinear disambiguation optimization. *Proceedings of the Royal Society A*, 478(2260):20210830, 2022.
- [7] R. Banisch and P. Koltai. Understanding the geometry of transport: Diffusion maps for lagrangian trajectory data unravel coherent sets. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 27(3):035804, 2017.
- [8] W. Bauer, P. Chandramouli, B. Chapron, L. Li, and E. Mémin. Deciphering the role of small-scale inhomogeneity on geophysical flow structuration: A stochastic approach. *Journal of Physical Oceanography*, 50(4):983 – 1003, 01 Apr. 2020.
- [9] T. Berlinet and C. Thomas-Agnan. *Reproducing kernel Hilbert spaces in Probability and Statistics*. Kluwer Academic Publishers, 2001.
- [10] T. Berry and T. Sauer. Local kernels and the geometric structure of data. *Applied and Computational Harmonic Analysis*, 40(3):439–469, 2016.

- [11] P. Bevanda, S. Sosnowski, and S. Hirche. Koopman operator dynamical models: Learning, analysis and control. *Annual Reviews in Control*, 52:197–212, 2021.
- [12] M. Bocquet, J. Brajard, A. Carrassi, and L. Bertino. Bayesian inference of chaotic dynamics by merging data assimilation, machine learning and expectation-maximization. *F. of Data Science*, 2(1):55–80, 2020.
- [13] S. Brunton, B. Brunton, J. Proctor, E. Kaiser, and J. Kutz. Chaos as an intermittently forced linear system. *Nat. Commun.*, 8(19), 2017.
- [14] S. Brunton, M. Budišić, E. Kaiser, and N. Kutz. Modern koopman theory for dynamical systems, 2021.
- [15] S. Brunton, J. Proctor, and N. Kutz. Discovering governing equations from data by sparse identification of nonlinear dynamical systems. *Proceedings of the National Academy of Sciences*, 113(15):3932–3937, 2016.
- [16] Steven L. Brunton, Bingni W. Brunton, Joshua L. Proctor, Eurika Kaiser, and J. Nathan Kutz. Chaos as an intermittently forced linear system. *Nature Communications*, 8, 2017.
- [17] Budišić, M. and Mohr, R. and Mezić, I. . Applied koopmanism. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 22(4):047510, 2012.
- [18] D. Burov, D. Giannakis, K. Manohar, and A. Stuart. Kernel analog forecasting: Multiscale test problems. *Multiscale Model. Simul.*, 19(2):1011–1040, jan 2021.
- [19] G. Buza, S. Jain, and G. Haller. Using spectral submanifolds for optimal mode selection in nonlinear model reduction. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 477(2246):20200725, 2021.
- [20] S. Chaturantabut and D. C. Sorensen. Nonlinear model reduction via discrete empirical interpolation. *SIAM Journal of Scientific Computing*, 32(5):2737–2764, September 2010.
- [21] R. R. Coifman and M. J. Hirn. Diffusion maps for changing data. *Applied and Computational Harmonic Analysis*, 36(1):79–107, 2014.
- [22] M. J. Colbrook and A. Townsend. Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems. *arXiv preprint <https://arxiv.org/abs/2111.14889>*, 2021.
- [23] F. Cucker and S. Smale. On the mathematical foundation of learning. *Bul. of the Amer. Math. Soc.*, 39(1):1–49, 2001.

- [24] S. Das and D. Giannakis. Koopman spectra in reproducing kernel Hilbert spaces. *Appl. Comput. Harmon. Anal.*, 49:573–607, 2020.
- [25] S. Das and D. Giannakis. Delay-coordinate maps and the spectra of Koopman operators. *J. Stat. Phys.*, 175:1107–1145, 2019.
- [26] S. Das, D. Giannakis, and J. Slawinska. Reproducing kernel Hilbert space compactification of unitary evolution groups. *Applied and Computational Harmonic Analysis*, 54:75–136, 2021.
- [27] A. M. DeGennaro and N. M. Urban. Scalable extended dynamic mode decomposition using random kernel approximation. *SIAM Journal on Scientific Computing*, 41(3):A1482–A1499, 2019.
- [28] M. Dellnitz and O. Junge. On the approximation of complicated dynamical behavior. *SIAM Journal on Numerical Analysis*, 36(2):491–515, 1999.
- [29] N. Ducet, P. Y. Le Traon, and G. Reverdin. Global high-resolution mapping of ocean circulation from topex/poseidon and ers-1 and -2. *Journal of Geophysical Research: Oceans*, 105(C8):19477–19498, 2000.
- [30] T. Eisner, B. Farkas, M. Haase, and R. Nagel. *Operator Theoretic Aspects of Ergodic Theory*. Graduate Texts in Mathematics. Springer, 2015.
- [31] D. Giannakis. Data-driven spectral decomposition and forecasting of ergodic dynamical systems. *Appl. Comput. Harmon. Anal.*, <https://doi.org/10.1016/j.acha.2017.09.001>, 2017.
- [32] D. Giannakis, J. Slawinska, and Z. Zhao. Spatiotemporal feature extraction with data-driven Koopman operators. In *Journ. of Mach. Learn. Res.: Workshop and Conference Proceedings*, volume 44, pages 103–115, 2015.
- [33] F. Gilani, D. Giannakis, and J. Harlim. Kernel-based prediction of non-Markovian time series. *Physica D: Nonlinear Phenomena*, 418:132829, 2021.
- [34] Georg A. Gottwald and Sebastian Reich. Supervised learning from noisy observations: Combining machine-learning techniques with data assimilation. *Physica D: Nonlinear Phenomena*, 423:132911, 2021.
- [35] R. J. Greatbatch and B. T. Nadiga. Four-gyre circulation in a barotropic model with double-gyre wind forcing. *Journal of Physical Oceanography*, 30(6):1461–1471, 2000.
- [36] F. Guillou, S. Metref, E. Cosme, C. Ubelmann, M. Ballarotta, J. Le Sommer, and J. Verron. Mapping altimetry in the forthcoming swot era by back-and-forth nudging a one-layer quasigeostrophic model. *Journal of Atmospheric and Oceanic Technology*, 38(4):697 – 710, 2021.

- [37] T. M. Hamill, J. S. Whitaker, and C. Snyder. Distance-dependent filtering of background error covariance estimates in an ensemble Kalman filter. *Monthly Weather Review*, 129(11):2776–2790, 2001.
- [38] S. Klus, I. Schuster, and K. Muandet. Eigendecompositions of transfer operators in reproducing kernel Hilbert spaces. *Journal of Nonlinear Science*, 30(1):283–315, 2020.
- [39] H. König. Eigenvalue distribution of compact operators with application to integral operators. *Linear Algebra and its application*, 84:111–122, 1986.
- [40] B. O. Koopman. Hamiltonian systems and transformation in Hilbert space. *Proceedings of the National Academy of Sciences*, 17(5):315–318, 1931.
- [41] N. Kutz, X. Fu, and S. Brunton. Multiresolution dynamic mode decomposition. *SIAM Journal on Applied Dynamical Systems*, 15(2):713–735, 2016.
- [42] P. Y. Le Traon, F. Nadal, and N. Ducet. An improved mapping method of multisatellite altimeter data. *Journal of Atmospheric and Oceanic Technology*, 15(2):522 – 534, 1998.
- [43] R. Lguensat, P. Tandeo, P. Ailliot, M. Pulido, and R. Fablet. The analog data assimilation. *Monthly Weather Review*, 145(10):4093 – 4107, 2017.
- [44] J. L. Lumley. The structure of inhomogeneous turbulent flows. *Atmospheric turbulence and radio wave propagation*, 1(1):166–178, 1967.
- [45] E. Mémin. Fluid flow dynamics under location uncertainty. *Geophys. & Astro. Fluid Dyn.*, 108(2):119–146, 2014.
- [46] I. Mezic. Spectral properties of dynamical systems, model reduction and decomposition. *Nonlinear Dynamics*, 41:309–325, 2005.
- [47] I. Mezić. Spectral properties of dynamical systems, model reduction and decompositions. *Nonlinear Dynamics*, 41(1-3):309–325, 2005.
- [48] H. Q. Minh. Some properties of Gaussian reproducing kernel Hilbert spaces and their implications for function approximation and learning theory. *Constructive Approximation*, 32(2):307–338, 2010.
- [49] S. Ouala, D. Nguyen, L. Drumetz, B. Chapron, A. Pascual, F. Collard, L. Gaultier, and R. Fablet. Learning latent dynamics for partially observed chaotic systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 30(10):103121, 2020.
- [50] S. Pan, N. Arnold-Medabalimi, and K. Duraisamy. Sparsity-promoting algorithms for the discovery of informative Koopman-invariant subspaces. *Journal of Fluid Mechanics*, 917:A18, 2021.

- [51] J. Pathak, B. Hunt, M. Girvan, Z. Lu, and E. Ott. Model-free prediction of large spatiotemporally chaotic systems from data: A reservoir computing approach. *Phys. Rev. Lett.*, 120:024102, Jan 2018.
- [52] J. Pathak, Z. Lu, B. Hunt, M. Girvan, and E. Ott. Using machine learning to replicate chaotic attractors and calculate Lyapunov exponents from data. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 27(12):121102, 2017.
- [53] M. Raissi, P. Perdikaris, and G.E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, 2019.
- [54] C. Rowley, I. Mezic, S. Bagheri, P. Schlatter, and D. Henningson. Spectral analysis of nonlinear flows. *J. Fluid Mech.*, 641:115–127, 2009.
- [55] C. Rowley S. Otto. Koopman operators for estimation and control of dynamical systems. *Annual Reviews of Control, Robotics, and Autonomous systems*, 4:59–87, 2021.
- [56] O. San, A. E. Staples, Z. Wang, and T. Iliescu. Approximate deconvolution large eddy simulation of a barotropic ocean circulation model. *Ocean Modelling*, 40(2):120–132, 2011.
- [57] Naratip Santitissadeekorn and Erik M. Bollt. Ensemble-based method for the inverse Frobenius–Perron operator problem: Data-driven global analysis from spatiotemporal “movie” data. *Physica D: Nonlinear Phenomena*, 411:132603, 2020.
- [58] P. Schmid. Dynamic mode decomposition of numerical and experimental data. *J. Fluid Mech.*, 656:5–28, 2010.
- [59] C.-J. Simon-Gabriel and B. Schölkopf. Kernel distribution embeddings: Universal kernels, characteristic kernels and kernel metrics on distributions. *Journal of Machine Learning Research*, 19(44):1–29, 2018.
- [60] I. Steinwart and A. Christmann. *Support Vector Machines. Information Science and Statistics*. Springer, 2008.
- [61] J. H. Tu, C. W. Rowley, D. M. Luchtenburg, S. L. Brunton, and J. N. Kutz. On dynamic mode decomposition: Theory and applications. *Journal of Computational Dynamics*, 1(2):391–421, 2014.
- [62] C. Ubelmann, B. Cornuelle, and L.-L. Fu. Dynamic mapping of along-track ocean altimetry: Method and performance from observing system simulation experiments. *Journal of Atmospheric and Oceanic Technology*, 33(8):1691–1699, 2016.

- [63] M. O. Williams, I. G. Kevrekidis, and C. W. Rowley. A data-driven approximation of the Koopman operator: Extending dynamic mode decomposition., *Journal of Nonlinear Science*, 25:1307–1346, 2015.
- [64] M. O. Williams, C. W. Rowley, and I. G. Kevrekidis. A kernel-based method for data-driven Koopman spectral analysis. *Journal of Computational Dynamics*, 2(2):247–265, 2015.
- [65] Zhizhen Zhao and Dimitrios Giannakis. Analog forecasting with dynamics-adapted kernels. *Nonlinearity*, 29(9):2888–2939, August 2016.
- [66] Y. Zhen, B. Chapron, E.Mémin, and L. Peng. Singular values of hankel matrix: A practical method to identify the koopman eigenfunctions. Technical report, arXiv:2107.01948, 2021.