



HAL
open science

Globally unsolvability of fractional evolution equations with exponential nonlinearity

Achouak Bekkai

► **To cite this version:**

Achouak Bekkai. Globally unsolvability of fractional evolution equations with exponential nonlinearity. CARI 2022, Oct 2022, Tunis, Tunisia. hal-03752999

HAL Id: hal-03752999

<https://inria.hal.science/hal-03752999>

Submitted on 17 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Globally unsolvability of fractional evolution equations with exponential nonlinearity

Achouak BEKKAI^{*1,1}

¹LAMIS Laboratory, Department of Mathematics and Informatic, University of Larbi Tebessi, Tebessa, Algeria

*E-mail : achouakbekkai@gmail.com

Abstract

In this paper the Cauchy problem for a time-space fractional evolution equation and a system with a time nonlocal nonlinearity of exponential growth are considered. The results on existence and uniqueness of local mild solution and nonexistence of global weak solution are presented, and under some suitable conditions, an estimate of the life span of blowing-up solutions is established.

Keywords

blow-up ; fractional integrals and derivatives ; life span ; local existence ; nonlinear fractional evolution equations

I INTRODUCTION

In this paper, we consider the following Cauchy problem:

$$\begin{cases} \mathbf{D}_{0|t}^\alpha u + (-\Delta)^{\beta/2} u = J_{0|t}^{1-\alpha} (e^u), & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^N, \end{cases} \quad (1.1)$$

where $N \geq 1, 0 < \alpha < 1, 0 < \beta \leq 2$, $\mathbf{D}_{0|t}^\alpha$ the Caputo fractional derivative operator of order α , $J_{0|t}^{1-\alpha} (e^u)$ is the Riemann-Liouville fractional integral of order $1 - \alpha$ for e^u , defined by

$$J_{0|t}^{1-\alpha} (e^u) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} e^{u(s)} ds,$$

where Γ is the gamma function, $(-\Delta)^{\beta/2}$ is the fractional Laplacian operator defined by

$$(-\Delta)^{\beta/2} u(x) = \mathcal{F}^{-1} \left(|\xi|^\beta \mathcal{F}(u)(\xi) \right) (x),$$

for $u \in D \left((-\Delta)^{\beta/2} \right) = H^\beta (\mathbb{R}^N)$, where $H^\beta (\mathbb{R}^N)$ is the homogeneous Sobolev space defined by

$$H^\beta (\mathbb{R}^N) = \left\{ u \in \mathcal{S}' ; (-\Delta)^{\beta/2} u \in L^2 (\mathbb{R}^N) \right\}, \text{ if } \beta \notin \mathbb{N},$$

$$H^\beta (\mathbb{R}^N) = \left\{ u \in L^2 (\mathbb{R}^N) ; (-\Delta)^{\beta/2} u \in L^2 (\mathbb{R}^N) \right\}, \text{ if } \beta \in \mathbb{N}$$

where \mathcal{S}' is the Schwartz space, \mathcal{F} is the Fourier transform, and \mathcal{F}^{-1} is its inverse, and $u_0 \in C_0(\mathbb{R}^N)$, where $C_0(\mathbb{R}^N)$ denotes the space of all continuous functions decaying to 0 at infinity.

We also consider local existence and global non-existence of the solution for the system:

$$\begin{cases} \mathbf{D}_{0|t}^{\alpha_1} u + (-\Delta)^{\beta/2} u = J_{0|t}^{1-\alpha_1}(e^v), & x \in \mathbb{R}^N, t > 0, \\ \mathbf{D}_{0|t}^{\alpha_2} v + (-\Delta)^{\beta/2} v = J_{0|t}^{1-\alpha_2}(e^u), & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \mathbb{R}^N. \end{cases} \quad (1.2)$$

Before we present and prove our results, let us dwell on existing literature related to our problems. The Cauchy problem for the semi-linear parabolic equation with a nonlinear memory

$$\begin{cases} u_t + (-\Delta)^{\beta/2} u = J_{0|t}^{1-\alpha}(|u|^{p-1}u), & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^N. \end{cases}$$

and system

$$\begin{cases} u_t + (-\Delta)^{\beta/2} u = J_{0|t}^{1-\alpha_1}(|v|^{p-1}v), & x \in \mathbb{R}^N, t > 0, \\ v_t + (-\Delta)^{\beta/2} v = J_{0|t}^{1-\alpha_2}(|u|^{q-1}u), & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \mathbb{R}^N, \end{cases}$$

has been considered by Fino and Kirane in [3] and [4].

In addition, Ahmad et al [2] considered the semi-linear nonlocal in time and space reaction diffusion equation:

$$\begin{cases} u_t + (-\Delta)^{\beta/2} u = J_{0|t}^{1-\alpha}(e^u), & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^N. \end{cases} \quad (1.3)$$

Moreover, the system of (1.3) was investigated by Ahmad et al [1] in the following form:

$$\begin{cases} u_t + (-\Delta)^{\beta/2} u = J_{0|t}^{1-\alpha_1}(e^v), & x \in \mathbb{R}^N, t > 0, \\ v_t + (-\Delta)^{\beta/2} v = J_{0|t}^{1-\alpha_2}(e^u), & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \mathbb{R}^N, \end{cases}$$

The here above cited papers treated the local existence, the blowing-up solutions and the blow-up profil of exploding solutions.

II FRACTIONAL DIFFUSION EQUATION WITH EXPONENTIAL NONLINEARITY

2.1 Local existence

In this section, we give the local existence and uniqueness of mild solution of the problem (1.1). First, we give the definition of a mild solution of (1.1).

Definition 1: Mild solution

Let $u_0 \in C_0(\mathbb{R}^N)$ and $T > 0$. We say that $u \in C([0, T]; C_0(\mathbb{R}^N))$ is a mild solution of (1.1) if

$$u(t) = P_{\alpha, \beta}(t) u_0 + \int_0^t (t-s)^{\alpha-1} S_{\alpha, \beta}(t-s) J_{0|s}^{1-\alpha}(e^{u(\tau)}) ds,$$

for $t \in [0, T]$.

Theorem 1: Local existence

Let $u_0 \in C_0(\mathbb{R}^N)$. Then, there exists a maximal time $T_{max} > 0$ such that problem (1.1) has a unique mild solution $u \in C([0, T_{max}); C_0(\mathbb{R}^N))$. Furthermore, either $T_{max} = +\infty$ or $T_{max} < +\infty$ and $\lim_{t \rightarrow T_{max}} \|u(t)\|_{L^\infty(\mathbb{R}^N)} = +\infty$. If, in addition, $u_0 \geq 0$, $u_0 \not\equiv 0$, then $u(t) > 0$ for all $0 < t < T_{max}$. Moreover, if $u_0 \in L^r(\mathbb{R}^N)$, for $1 \leq r < \infty$, then $u \in C([0, T_{max}); L^r(\mathbb{R}^N))$.

2.2 Blowing up solutions

Definition 2: Weak solution

Let $u_0 \in L_{loc}^\infty(\mathbb{R}^N)$ and $T > 0$. We say that u is a weak solution of (1.1) if $u \in L^p((0, T); L_{loc}^\infty(\mathbb{R}^N))$ and verify the equation

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^N} u_0 \mathbf{D}_{t|T}^\alpha \psi(x, t) + \int_0^T \int_{\mathbb{R}^N} J_{0|t}^{1-\alpha}(e^u) \psi(x, t) \\ &= \int_0^T \int_{\mathbb{R}^N} u(x, t) (-\Delta)^{\beta/2} \psi(x, t) + \int_0^T \int_{\mathbb{R}^N} u(x, t) \mathbf{D}_{t|T}^\alpha \psi(x, t), \end{aligned}$$

for all test function $\psi \in C^1([0, T]; H^\beta(\mathbb{R}^N))$ such that $\psi(x, T) = 0$.

Lemma 2:

Consider $u_0 \in C_0(\mathbb{R}^N)$ and let $u \in C([0, T]; C_0(\mathbb{R}^N))$ be a mild solution of (1.1), then u is also a weak solution of (1.1).

Theorem 3:

Let $u_0 \in C_0(\mathbb{R}^N)$ with $u_0 \geq 0$ and $u_0 \not\equiv 0$. Then, the solution of (1.1) blows up in a finite time.

2.3 Life span of blowing up solutions

In this section, we give an upper bound estimate of the life span of the blowing up solutions with some special initial datum. To this aim, we consider the following problem:

$$\begin{cases} \mathbf{D}_{0|t}^\alpha u_\epsilon + (-\Delta)^{\beta/2} u_\epsilon = J_{0|t}^{1-\alpha}(e^{u_\epsilon}), & x \in \mathbb{R}^N, t > 0, \\ u_\epsilon(x, 0) = \epsilon u_0(x), & x \in \mathbb{R}^N, \end{cases} \quad (1.4)$$

where $\epsilon > 0$, $0 < \alpha < 1$, $0 < \beta \leq 2$ and $u_0 \in C_0(\mathbb{R}^N)$ satisfies

$$u_0(x) \geq C_0 |x|^{-\delta}, \quad |x| \geq \epsilon_0, \quad N < \delta < \frac{\beta}{\alpha}, \quad (1.5)$$

for some positive constants C_0 and ϵ_0 .

Theorem 4:

Suppose that (1.5) holds. Let $[0, T_\epsilon]$ be the life span of the solution u_ϵ of the problem (1.4). Then, there exists a positive constant C such that

$$T_\epsilon \leq C\epsilon^{\frac{1}{\theta}}, \quad \theta = \frac{\alpha\delta}{\beta} - 1 < 0.$$

III FRACTIONAL DIFFUSION SYSTEM WITH NONLINEARITIES OF EXPONENTIAL GROWTH

In this section, we study the system (1.2). We start with the definition of the mild solution of (1.2).

3.1 Local existence

Definition 3: Mild solution

Let $u_0, v_0 \in C_0(\mathbb{R}^N)$. We say that $(u, v) \in C([0, T], C_0(\mathbb{R}^N)) \times C([0, T], C_0(\mathbb{R}^N))$ is a mild solution of (1.2) if (u, v) satisfies, for $t \in [0, T]$, the following equations

$$\begin{aligned} u(t) &= P_{\alpha_1, \beta}(t) u_0 + \int_0^t (t-s)^{\alpha_1-1} S_{\alpha_1, \beta}(t-s) J_{0|s}^{1-\alpha_1}(e^{v(\tau)}), \\ v(t) &= P_{\alpha_2, \beta}(t) v_0 + \int_0^t (t-s)^{\alpha_2-1} S_{\alpha_2, \beta}(t-s) J_{0|s}^{1-\alpha_2}(e^{u(\tau)}). \end{aligned}$$

Theorem 5: Local existence

Let $u_0, v_0 \in C_0(\mathbb{R}^N)$. Then, there exists a maximal time $T_{max} > 0$ such that the problem (1.2) has a unique mild solution $(u, v) \in C([0, T_{max}], C_0(\mathbb{R}^N)) \times C([0, T_{max}], C_0(\mathbb{R}^N))$. Furthermore, we have the alternative: - either $T_{max} = +\infty$; - or $T_{max} < +\infty$ and $\lim_{t \rightarrow T_{max}} (\|u(t)\|_{L^\infty(\mathbb{R}^N)} + \|v(t)\|_{L^\infty(\mathbb{R}^N)}) = +\infty$. If, in addition, $u_0, v_0 \geq 0, u_0 \not\equiv 0, v_0 \not\equiv 0$, then $u(t), v(t) > 0$ for all $0 < t < T_{max}$. Moreover, if $u_0, v_0 \in L^r(\mathbb{R}^N)$, for $1 \leq r < \infty$, then $u, v \in C([0, T_{max}]; L^r(\mathbb{R}^N))$.

3.2 Blowing up solutions

Definition 4: Weak solution

Let $u_0, v_0 \in L_{loc}^\infty(\mathbb{R}^N)$ and $T > 0$. We say that (u, v) is a weak solution of (1.2) if $(u, v) \in L^p((0, T); L_{loc}^\infty(\mathbb{R}^N)) \times L^p((0, T); L_{loc}^\infty(\mathbb{R}^N))$ and satisfies

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^N} u_0 \mathbf{D}_{t|T}^{\alpha_1} \psi_1(x, t) dx dt + \int_0^T \int_{\mathbb{R}^N} J_{0|t}^{1-\alpha_1}(e^v) \psi_1(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{R}^N} u(x, t) (-\Delta)^{\beta/2} \psi_1(x, t) + \int_0^T \int_{\mathbb{R}^N} u(x, t) \mathbf{D}_{t|T}^{\alpha_1} \psi_1(x, t), \\ & \int_0^T \int_{\mathbb{R}^N} v_0 \mathbf{D}_{t|T}^{\alpha_2} \psi_2(x, t) dx dt + \int_0^T \int_{\mathbb{R}^N} J_{0|t}^{1-\alpha_2}(e^u) \psi_2(x, t) dx dt \\ &= \int_0^T \int_{\mathbb{R}^N} v(x, t) (-\Delta)^{\beta/2} \psi_2(x, t) + \int_0^T \int_{\mathbb{R}^N} v(x, t) \mathbf{D}_{t|T}^{\alpha_2} \psi_2(x, t), \end{aligned}$$

for all test functions $\psi_1, \psi_2 \in C^1([0, T]; H^\beta(\mathbb{R}^N))$ such that $\psi_1(x, T) = \psi_2(x, T) = 0$.

Theorem 6:

Let $u_0, v_0 \in C_0(\mathbb{R}^N)$ with $u_0 \geq 0, u_0 \not\equiv 0, v_0 \geq 0$ and $v_0 \not\equiv 0$. blows up in a finite time.

3.3 Life span of blowing up solutions

We consider the problem:

$$\begin{cases} \mathbf{D}_{0|t}^{\alpha_1} u_\epsilon + (-\Delta)^{\beta/2} u_\epsilon = J_{0|t}^{1-\alpha_1} (e^{v_\epsilon}), & x \in \mathbb{R}^N, t > 0, \\ \mathbf{D}_{0|t}^{\alpha_2} v_\epsilon + (-\Delta)^{\beta/2} v_\epsilon = J_{0|t}^{1-\alpha_2} (e^{u_\epsilon}), & x \in \mathbb{R}^N, t > 0, \\ u_\epsilon(x, 0) = \epsilon u_0(x), \quad v_\epsilon(x, 0) = \epsilon v_0(x), & x \in \mathbb{R}^N, \end{cases} \quad (1.6)$$

where $\epsilon > 0, 0 < \alpha_1, \alpha_2 < 1, 0 < \beta \leq 2$ with $u_0, v_0 \in C_0(\mathbb{R}^N)$ satisfying

$$u_0(x) \geq m_0 |x|^{-\frac{\delta}{\alpha_1}}, \quad v_0(x) \geq n_0 |x|^{-\frac{\delta}{\alpha_2}}, \quad |x| \geq \epsilon_0, \quad \max\{\alpha_1, \alpha_2\} N < \delta < \beta, \quad (1.7)$$

for some positive constants m_0, n_0 and ϵ_0 .

Theorem 7:

Suppose that (1.7) holds. Let $[0, T_\epsilon)$ be the maximal time of existence of the solution (u_ϵ, v_ϵ) of problem (1.6). Then there exists a constant $C > 0$ such that

$$T_\epsilon \leq C \epsilon^{\frac{1}{\eta}}, \quad \eta = \frac{\delta}{\beta} - 1 < 0.$$

IV REFERENCES

REFERENCES

- B. Ahmad, A. Alsaedi, D. Hnaien, M. Kirane, "On a semi-linear system of nonlocal time and space reaction diffusion equations with exponential nonlinearities," J. Equations Appl. 30 (1) (2018), pp. 17–40.
- A. Alsaedi, B. Ahmad, M. Kirane, "On a reaction diffusion equation with nonlinear time-nonlocal source term," Math. Meth. Appl. Sci. 39 (2) (2016), pp. 236–244.
- A. Kirane M, "Qualitative properties of solutions to a nonlocal evolution system," Math Meth Appl Sci. 34 (9) (2011), pp. 1125-1143.
- A. Fino, M. Kirane, "Qualitative properties of solutions to a time-space fractional evolution equation," Quart. Appl. Math. 70 (1) (2012), pp. 133–157.
- G. H. Hardy, J. E. Littlewood, G. Pólya, "Inequalities," Cambridge: Cambridge University Press, 1952.
- S. G. Samko, A. A. Kilbas, O. I. Marichev, "Fractional integrals and derivatives. Theory and applications," Gordon and Breach Science Publishers, 1993.
- Q. G. Zhang, H. R. Sun, "The blow-up and global existence of solutions of Cauchy problems for a time fractional diffusion equation," Topol Methods Nonlinear Anal. 46 (1) (2015), 69–92.