



HAL
open science

The Topological Correctness of PL Approximations of Isomanifolds

Jean-Daniel Boissonnat, Mathijs Wintraecken

► **To cite this version:**

Jean-Daniel Boissonnat, Mathijs Wintraecken. The Topological Correctness of PL Approximations of Isomanifolds. Foundations of Computational Mathematics, 2021, 22, pp.967 - 1012. 10.1007/s10208-021-09520-0 . hal-03760378

HAL Id: hal-03760378

<https://inria.hal.science/hal-03760378>

Submitted on 25 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The Topological Correctness of PL Approximations of Isomanifolds

Jean-Daniel Boissonnat¹ · Mathijs Wintraecken²

Received: 18 June 2020 / Accepted: 7 April 2021 / Published online: 13 July 2021
© The Author(s) 2021

Abstract

Isomanifolds are the generalization of isosurfaces to arbitrary dimension and codimension, i.e. manifolds defined as the zero set of some multivariate multivalued smooth function $f : \mathbb{R}^d \rightarrow \mathbb{R}^{d-n}$. A natural (and efficient) way to approximate an isomanifold is to consider its piecewise-linear (PL) approximation based on a triangulation \mathcal{T} of the ambient space \mathbb{R}^d . In this paper, we give conditions under which the PL approximation of an isomanifold is topologically equivalent to the isomanifold. The conditions are easy to satisfy in the sense that they can always be met by taking a sufficiently fine and thick triangulation \mathcal{T} . This contrasts with previous results on the triangulation of manifolds where, in arbitrary dimensions, delicate perturbations are needed to guarantee topological correctness, which leads to strong limitations in practice. We further give a bound on the Fréchet distance between the original isomanifold and its PL approximation. Finally, we show analogous results for the PL approximation of an isomanifold with boundary.

Keywords Isomanifold · Solution manifolds · Piecewise-linear approximation · Isotopy · Fréchet distance

Communicated by Ulrike Tillmann.

This work was funded by the European Research Council under the European Union's ERC Grant Agreement number 339025 GUDHI (Algorithmic Foundations of Geometric Understanding in Higher Dimensions). This work was also supported by the French government, through the 3IA Côte d'Azur Investments in the Future project managed by the National Research Agency (ANR) with the reference number ANR-19-P3IA-0002. Mathijs Wintraecken also received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 754411.

✉ Mathijs Wintraecken
m.h.m.j.wintraecken@gmail.com

Jean-Daniel Boissonnat
jean-daniel.boissonnat@inria.fr

¹ Université Côte d'Azur, Inria, Sophia-Antipolis, France

² IST, Austria, Klosterneuburg, Austria

Mathematics Subject Classification 57Q55 · 65D18 · 68U05 · 51-08 · 32B25 · 53Z50

1 Introduction

Isosurfacing

Given a surface represented in \mathbb{R}^3 as the zero set of a function $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, the goal of isosurfacing is to find a piecewise-linear (PL) approximation of the surface. This question naturally extends to higher dimensions and codimensions, in which case the generalized surface is called an isomanifold. Isosurfaces play a crucial role in medical imaging, computer graphics and geometry processing [1]. Higher-dimensional isomanifolds are also of fundamental importance in many fields such as statistics [2], dynamical systems [3], econometrics or mechanics [1].

Marching Algorithms

The standard algorithmic solution to the isosurfacing problem is to use some marching algorithm. This approach was initiated by Lorensen and Cline, with their marching cube algorithm [4]. Many variants of the algorithm have been introduced, see, for example, [5–8], and the overview [1]. The approach, however, is always the same: one first subdivides the ambient space into cubes (in which case the algorithm is called a marching cube algorithm), or simplices [5,6,9] (in which case the algorithm is called a marching tetrahedra algorithm). One starts with a cube or a simplex (cell) in which a part of the zero set of the function is contained, and finds a piecewise-linear approximation of the zero set in that cell. One then propagates or marches to adjacent cells that also contain the zero set and approximates the zero set in that cell. This process can be continued until all cells that intersect the zero set have been visited.

For the marching cube algorithm [4], one also has to decide how to approximate the zero set inside a cube. As observed by Dürst, there is in general no canonical way how to do this due to *ambiguous configurations*, see [10] for an extensive discussion in the three-dimensional setting. For the marching tetrahedra algorithm, there is a canonical way to construct a piecewise-linear approximation of the zero set, as we will discuss below. However, the result of the algorithm is still not necessarily topologically correct.

Guarantees for Isosurfacing

For the marching simplex algorithm [5] in arbitrary dimensions, bounds have been given on the one-sided Hausdorff distance between the zero set of f and its PL approximation, and also on the difference between the gradient of f and the gradient of the PL approximation. It can be proven that the result of the algorithm is a manifold under appropriate assumptions [11,12].

An important requirement in the work of Allgower and Georg [12] is that the zero set avoids simplices that have dimension less than the codimension, see [12, Definition 12.2.2] and the text above [12, Theorem 15.4.1]. The idea to avoid these low-dimensional simplices originates with Whitney [13], with whom Allgower and

George [11,12] were apparently unfamiliar. Very heavy perturbation schemes for the vertices of the ambient triangulation \mathcal{T} are needed to ensure that the manifold stays sufficiently far from simplices in the ambient triangulation that have dimension less than the codimension of the manifold [13,14]. Various techniques have been developed to compute such perturbations with guarantees. They typically consist in perturbing the position of the sample points or in assigning weights to the points. Complexity bounds are then obtained using volume arguments. See, for example, [15–18]. However, these techniques suffer from several drawbacks. The constants in the complexity depend exponentially on the ambient dimension. Moreover, the analysis assumes that the probability of the simplices of dimension less than the codimension to intersect the manifold is zero, which is not true when dealing with finite precision. As a result, the actual implementations we are aware of fail to work well in practice except in very simple cases.

More complete correctness results have been achieved in three dimensions in the computational geometry community:

Boissonnat, Cohen-Steiner and Vegter [19] base their proof on a combination of Morse theory and simplicial collapses. Vegter and Plantinga's proof [20] is in its philosophy closely related to normal surface theory, see, for example, [21], but relies rather heavily on case analysis. The results of [19,20] seem not extendable to higher dimensions.

Triangulating General Manifolds (Without Boundary)

The approximation of a manifold that is the zero set of a function is an example of the more general question of how to triangulate a manifold. It is known that C^1 manifolds are triangulable, see, for example, [13], and algorithms have been proposed recently to triangulate smooth manifolds [14,17,22,23]. However, all known methods use intricate perturbation schemes to guarantee the correctness of the triangulation algorithms when the intrinsic dimension of the manifold exceeds 2. As for the case of isomanifolds, perturbation schemes work fine in theory but the constants are miserable and the methods do not work in practice in high dimensions.

Manifolds with Boundary

In this paper, we also consider the piecewise-linear approximation of manifolds with boundary (that are given as a zero set) and briefly mention the extension to stratifolds. Apart from some Delaunay-based work on triangulations of stratifolds in three dimensions [24–28], we are not aware of similar results on manifolds with boundary. Significant effort also went in the detection of strata, in this case in arbitrary dimension, see, for example, [29–31].

Contribution

This paper contains three main results, the first two (Theorem 25 and Corollary 27) concern manifolds without boundary and the third (Theorem 48) manifolds with boundary.

We state here simplified versions of the statements to be fully described later on.

Isomanifolds (without boundary)

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}^{d-n}$ be a smooth function and suppose that 0 is a regular value of f , meaning that at every point x such that $f(x) = 0$, the Jacobian of f is non-degenerate. Assume that \mathcal{T} is a triangulation of \mathbb{R}^d . Define the function f_{PL} as the linear interpolation of the values of f at the vertices if restricted to a single simplex $\sigma \in \mathcal{T}$. Then,

Theorem 25 (Ambient isotopy) *The zero set of f_{PL} is a manifold that is ambient isotopic to the zero set of f , provided that the triangulation \mathcal{T} is sufficiently fine and thick.*

We recall that the thickness of a simplex is the ratio of the height (smallest altitude) over the longest edge length, and it is a measure for the quality or how well shaped a simplex is.

Corollary 27 (Bound on the Fréchet distance) *The Fréchet distance between f_{PL} and f is of the order of D^2 , where D is the longest edge length of \mathcal{T} .*

We also give a variant of a result due to Allgower and George [11]:

Proposition 10 *The difference between the gradient of f and the gradient of its piecewise-linear approximation is of order dD inside each simplex of \mathcal{T} .*

Isomanifolds with boundary Suppose that apart from f we are also given another function $f_\partial : \mathbb{R}^d \rightarrow \mathbb{R}$ and $f_{\partial,PL}$ is defined similar to f_{PL} . Write f^i for the i th component of f . Let us further assume that the zero set is regular in the following sense: The gradients of f^i span a $(d-n)$ -dimensional space at each point of $f^{-1}(0)$ and the gradients of f^i and f_∂ span a $(d-n+1)$ -dimensional space at each point of $\partial M = f^{-1}(0) \cap f_\partial^{-1}(0)$. Then,

Theorem 48 (Manifolds with boundary) *The set $f^{-1}(0) \cap f_\partial^{-1}([0, \infty))$ is ambient isotopic to $f_{PL}^{-1}(0) \cap f_{\partial,PL}^{-1}([0, \infty))$, provided that the triangulation \mathcal{T} is sufficiently fine and thick.*

An important aspect of these results is that they hold under mild conditions: they simply ask for a sufficiently fine and thick triangulation \mathcal{T} . In contrast to previous results on the triangulation of manifolds, no perturbations are needed to guarantee topological correctness.

Our method provides guarantees on the piecewise-linear (PL) approximation of isomanifolds, regarding the topology, the Fréchet distance and the approximation of the gradients (the latter was already known to Allgower and Georg [11]).

However, we stress that it does not give lower bounds on the quality of the linear pieces in the PL approximation. This is a clear difference with previous methods [13, 14, 16, 23] whose output is a thick triangulation. Although this is an appealing property, it complicates the analysis further and requires unpractical perturbation schemes. Such perturbation techniques could be added to our method to improve the simplex quality (to some limited extent). However, they are not required to make the algorithm work and to obtain the guarantees mentioned above.

The techniques used in this paper are also different from many of the standard tools and do not rely on

Delaunay triangulations [32,33], the closed ball property [15,34,35], Whitney’s lemma [36] or collapses [37]. The current paper mainly relies on the non-smooth implicit function theorem [38] with some Morse theory.

Outline

The rest of this paper is subdivided as follows. In Sect. 2, we treat closed isomanifolds, i.e. compact manifolds without boundary. In Sect. 3, we treat isomanifolds with boundary. Extension to general isostratifolds is briefly discussed in Sect. 4. In the final section, we quantify the robustness of the method by studying how much the zero set of f changes if f is perturbed slightly in the C^1 -sense [39].

This paper is closely related to another paper where the data structure needed to efficiently propagate along the manifold is presented [40]. Altogether, these two papers show that one can construct PL approximations of isomanifolds in space and time polynomial in the resolution $1/D$ of the ambient triangulation and in the dimension d of the ambient space.

2 Isomanifolds (Without Boundary)

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}^{d-n}$ be a smooth (C^2 suffices) function and suppose that 0 is a regular value of f , meaning that at every point x such that $f(x) = 0$, the Jacobian of f is non-degenerate. Then, the zero set of f is an n -dimensional manifold as a direct consequence of the implicit function theorem, see, for example, [41, Section 3.5]. We further assume that $f^{-1}(0)$ is compact. As in [11] we consider a triangulation \mathcal{T} of \mathbb{R}^d . The function f_{PL} is the linear interpolation of the values of f at the vertices if restricted to a single simplex $\sigma \in \mathcal{T}$, i.e.

$$\forall x \in \sigma : f_{PL}(x) = \sum_{v \in \sigma} \lambda_v(x) f(v),$$

where the λ_v are the barycentric coordinates of x with respect to the vertices of σ . For any function $g : \mathbb{R}^d \rightarrow \mathbb{R}^{d-n}$, we write g^i , with $i = 1, \dots, d - n$, for the components of g .

We prove that under certain conditions there is an ambient isotopy from the zero set of f to the zero set of f_{PL} . The proof will be using the piecewise-smooth map

$$F_{PL}(x, \tau) = (1 - \tau)f(x) + \tau f_{PL}(x), \tag{1}$$

which interpolates between f and f_{PL} and is based on the generalized implicit function theorem.

We are, by definition, only interested in $f^{-1}(0)$ and so can ignore points that are sufficiently far from this zero set. More precisely, we observe the following: if $f^i(x)$ is positive for all x in a geometric simplex σ , then so is $f_{PL}^i(x)$ because $f_{PL}^i(x)$ is a convex combination of the (positive) values at the vertices. This in turn implies that

$F_{PL}^i(x, \tau)$ is positive on $\sigma \times [0, 1]$ as, for each τ , it is a convex combination of positive numbers. The same argument holds for negative values. So we see that

Remark 1 Write \mathcal{T}_0 for the set of all $\sigma \in \mathcal{T}$, such that $(f^i)^{-1}(0) \cap \sigma \neq \emptyset$ for all i . Then, for all τ , $\{x \mid F_{PL}(x, \tau) = 0\} \subset \mathcal{T}_0$.

The results will be expressed using constants defined in terms of f and the ambient triangulation \mathcal{T} .

Definition 2 We define

$$\gamma_{\max} = \max_{x \in \mathcal{T}_0} (\max_i |\nabla f^i(x)|) \tag{2}$$

$$\lambda_{\min} = \min_{x \in \mathcal{T}_0} \lambda_{\min}(x), \tag{3}$$

$$\alpha_{\max} = \max_{x \in \mathcal{T}_0} \max_i \|\text{Hes}(f^i)(x)\|_2 \tag{4}$$

$$D : \text{the longest edge length of a simplex in } \mathcal{T}_0 \tag{5}$$

$$T : \text{the smallest thickness of a simplex in } \mathcal{T}_0, \tag{6}$$

where

- $\nabla f^i = (\partial_j f_i)_j$ denotes the gradient of component f^i , for $i \in [1, d - n]$,
- $\text{Gram}(\nabla f)$ denotes the Gram matrix whose elements are $\nabla f^i \cdot \nabla f^j$ where \cdot stands for the dot product,
- $\lambda_{\min}(x)$ denotes the smallest absolute value of the eigenvalues of $\text{Gram}(\nabla f(x))$,¹
- $\text{Hes}(f) = (\partial_k \partial_l f_i)_{k,l}$ denotes the Hessian matrix of second order derivatives,
- $|\cdot|$ denotes the Euclidean norm of a vector and $\|\cdot\|_2$ the operator 2-norm of a matrix.²
- The thickness is the ratio of the height (smallest altitude) over the longest edge length.

We will assume that $\gamma_{\max}, \lambda_{\min}, \alpha_{\max}, D, T \in (0, \infty)$. The constant λ_{\min} quantifies how close 0 is to not being a regular value of f . D is a measure of the size of the simplices of \mathcal{T} . We will call $\delta = 1/D$ the resolution of \mathcal{T} . The thickness is a quality measure of a simplex. A good choice for \mathcal{T} is the Coxeter triangulation of type A_d , see [42,43], or the related Freudenthal triangulations, see [3,44–46], which can be defined for different values of D while keeping T constant (for a given dimension d).

Our results hold for any dimensions d and n . We are especially interested in the case where the ambient dimension d is large. We thus consider d and D as the two main parameters. For our bounds, we will give both exact and asymptotic expressions. The asymptotic expressions are given to emphasize the dependency on the two most important parameters d and D , and hold for any d, D and T such that $dD < T$ and for any fixed positive $\gamma_{\max}, \lambda_{\min}, \alpha_{\max}$. For Coxeter triangulations of type \tilde{A}_d (the only type of Coxeter triangulations mentioned in this paper), we have $T > \frac{2}{\sqrt{d+2}}$,

¹ Because a Gram matrix is a symmetric square matrix, its eigenvalues are well defined and real.

² The operator norm is defined as $\|A\|_p = \max_{x \in \mathbb{R}^n} \frac{|Ax|_p}{|x|_p}$, with $|\cdot|_p$ the p -norm on \mathbb{R}^n .




Fig. 1 A pictorial overview of the proof. The τ -direction goes upwards. Similar to Morse theory, we find that $f_{PL}^{-1}(0)$ (top) and $f^{-1}(0)$ (bottom) are ambient isotopic if the function τ restricted to $F_{PL}^{-1}(0)$ does not encounter a Morse critical point

see [43]. Therefore, the condition $dD < T$ is satisfied for these triangulations when $D < (2d\sqrt{d} + 2)^{-1}$. For convenience, exact expressions are gathered in Appendix A.

The Result

We are going to construct an ambient isotopy based on (1), see Fig. 1 for a pictorial overview. In fact, the map $\tau \mapsto \{x \mid F_{PL}(x, \tau) = 0\}$ gives an ambient isotopy between the zero set of $F_{PL}(x, 0)$, which is identical to the smooth isosurface $f^{-1}(0)$, and the zero set of $F_{PL}(x, 1)$, which is the PL approximation $f_{PL}^{-1}(0)$. The latter can be turned into a triangulation of the isosurface $f^{-1}(0)$ by triangulating the non-simplicial cells using barycentric subdivision. We will also bound the Fréchet distance between $f^{-1}(0)$ and $f_{PL}^{-1}(0)$.

Proving the ambient isotopy consists of three technical steps. The first two consume most of the space in the proof namely:

- *Local step.* Let $\sigma \in \mathcal{T}$. We first show that $\{(x, \tau) \mid F_{PL}(x, \tau) = 0\} \cap (\sigma \times [0, 1])$ is a smooth manifold, under certain conditions (Corollary 13).
- *Global step.* We prove that $F_{PL}^{-1}(0)$ is a manifold, under certain conditions, using techniques from non-smooth analysis (Corollary 24).

A crucial ingredient will be the implicit function theorem and its non-smooth extension. Along the way, we shall also see that $F_{PL}^{-1}(0)$ is never tangent to the $\tau = c$ planes, where c is a constant. The gradient of $(x, \tau), \mapsto \tau$ in

$\mathbb{R}^d \times \mathbb{R}$ is $(0, 1)$. Projecting this vector onto the tangent space of $F_{PL}^{-1}(0)$ gives the gradient of $(x, \tau), \mapsto \tau$ restricted to $F_{PL}^{-1}(0)$. Because of the non-tangency, this projection is nonzero. So the gradient field of the function $(x, \tau), \mapsto \tau$ restricted to $F_{PL}^{-1}(0)$, is piecewise-smooth (because $F_{PL}^{-1}(0)$ is piecewise-smooth) and never vanishes.

The third step is similar to a standard observation in Morse theory [47,48], with the exception that we now consider piecewise-smooth instead of smooth vector fields. We refer to Milnor [47] for an excellent introduction and to Lemma 2.4 and Theorem 3.1 in particular.

Lemma 3 (Gradient flow induced isotopies) *The flow of a non-vanishing piecewise-smooth gradient vector field of a function τ on a compact manifold generates a isotopy from $\tau = c_1$ to $\tau = c_2$, where c_1 and c_2 are constants.*

Proof This is a straightforward consequence of the existence and uniqueness of the solution to a differential equation. □

Bounds on the gradient of τ on the manifold give a bound on the Fréchet distance, which is defined in the following.

Definition 4 (*Fréchet distance for embedded manifolds*) Let \mathcal{M} and \mathcal{M}' be two homeomorphic, compact submanifolds of \mathbb{R}^d . Write \mathcal{H} for the set of all homeomorphisms from \mathcal{M} to \mathcal{M}' . The Fréchet distance between \mathcal{M} and \mathcal{M}' is

$$d_F(\mathcal{M}, \mathcal{M}') = \inf_{h \in \mathcal{H}} \sup_{x \in \mathcal{M}} d(x, h(x)).$$

2.1 Preliminaries

The following elementary lemma will be useful.

Lemma 5

$$|f^i(x_1) - f^i(x_2)| \leq \gamma_{\max} |x_1 - x_2|. \tag{7}$$

$$|\nabla f^i(x_1) - \nabla f^i(x_2)| \leq d\alpha_{\max} |x_1 - x_2|. \tag{8}$$

Proof The first statement follows from the fact that the supremum of the absolute value of a derivative of a function bounds the Lipschitz constant of the function. The second statement follows from standard bounds on matrix norm (see, for example, [49, Equation (2.3.11)]) together with (4). These bounds imply that $\sqrt{d}\alpha_{\max} \geq |\partial_k \partial_l f^i|$,

for all k, l and i . Arguing as before, we deduce a bound on the Lipschitz constant of $\partial_l f^i$:

$$|\partial_l f^i(x_1) - \partial_l f^i(x_2)| \leq \sqrt{d} \alpha_{\max} |x_1 - x_2|.$$

Bound (8) now follows. □

2.1.1 The Implicit Function Theorem

The main technical tool to prove the existence of the ambient isotopy from $f^{-1}(0)$ to $f_{PL}^{-1}(0)$ is the implicit function theorem which we recall now.

Theorem 6 (Smooth implicit function theorem) *Let $F : \mathbb{R}^{d+1} \rightarrow \mathbb{R}^{d-n}$ be a continuously differentiable function. Write $\mathbb{R}^{d+1} = \mathbb{R}^{n+1} \times \mathbb{R}^{d-n}$ and denote the coordinates of \mathbb{R}^{d+1} by (x, y) accordingly. Fix a point (a, b) , with $F(a, b) = 0 \in \mathbb{R}^{d-n}$. If the Jacobian $J_{F,y}(a, b) = (\frac{\partial F^i}{\partial y^j}(a, b))_{i,j}$ is of maximal rank (or equivalently is invertible), then there exists an open set $U \subset \mathbb{R}^{n+1}$ containing a such that there exists a unique continuously differentiable function $g : U \rightarrow \mathbb{R}^{d-n}$ such that $g(a) = b$ and $F(x, g(x)) = 0$ for all $x \in U$.*

To prove the existence of the isotopy from $f^{-1}(0)$ to $f_{PL}^{-1}(0)$, we will apply the implicit function theorem (Theorem 6) to several functions g that are close to f and we will therefore need to prove that their Jacobians are of maximal rank. A matrix has maximal rank if and only if the Gram matrix of its columns has a nonzero determinant or, equivalently, nonzero eigenvalues. In our context, we will need lower bounds on the absolute values of the eigenvalues of the Gram matrices $\text{Gram}(\nabla g)$, given the lower bound λ_{\min} on the absolute values of the eigenvalues of $\text{Gram}(\nabla f)$.

2.1.2 Eigenvalues and Perturbations

We will follow the convention that the eigenvalues of the matrices we consider are sorted by increasing order of their absolute values, i.e. $|\lambda_i| \leq |\lambda_j|$ if $i \leq j$. We first recall Weyl’s perturbation theorem that bounds the difference between the i th eigenvalues of two symmetric matrices:

Lemma 7 (Weyl’s bound, Corollary III.2.6 of [50]) *Let A and $\tilde{A} = A + E$ be two symmetric (or Hermitian) matrices and write λ_i and $\tilde{\lambda}_i$ for the eigenvalues of A and \tilde{A} , respectively. Then,*

$$\max_i |\lambda_i - \tilde{\lambda}_i| \leq \|E\|_2,$$

where $\|\cdot\|_p$ denotes the p -norm.

We further note that $\|E\|_2 \leq \|E\|_F$ where $\|\cdot\|_F$ denotes the Frobenius norm, see [49, (2.3.7)]. By definition of the Frobenius norm, we have that $|E_{ij}| \leq e_{\max}$, for all $i, j \in [1, d - n]$, implies that $\|E\|_F \leq (d - n)e_{\max}$ if $d - n$ is the dimension of E . Hence, we have

Corollary 8 *Under the conditions of Lemma 7, and assuming $\dim(E) = d - n$, and $|E_{ij}| \leq e_{\max}$, we have*

$$\max_i |\lambda_i - \tilde{\lambda}_i| \leq (d - n)e_{\max}.$$

2.2 Estimates for a Single Simplex

In this section, we concentrate on a single simplex σ and write f_L for the linear function whose values on the vertices of σ coincide with f . In other words, f_L is the linear extension of the interpolation of f . Note that f_L coincides with f_{PL} within the geometric simplex σ (but not necessarily outside).

2.2.1 Estimates on the Linear Approximation f_L and Its Gradient

We need a simple estimate similar to Proposition 2.1 of Allgower and George [11].

Lemma 9 *Let $\sigma \subset \mathcal{T}_0$ and let f_L be as described above. Then, for all $x \in \sigma$,*

$$|f_L^i(x) - f^i(x)| \leq 2D^2\alpha_{\max}.$$

We included a proof for completeness.

Proof Let v_k be a vertex of σ . Taylor’s theorem, see, for example, [41, Theorem 2.8.4], yields that

$$f^i(v_k) = f^i(x) + \sum_j \partial_j f^i(x)(v_k - x)^j + R(v_k), \tag{9}$$

with

$$\begin{aligned} R(v_k) &= 2 \sum_{j \neq l} (v_k - x)^j (v_k - x)^l \int_0^1 (1 - t)^2 \partial_j \partial_l f^i(v_k - t(v_k - x)) dt \\ &\quad + 2 \sum_j \frac{((v_k - x)^j)^2}{2} \int_0^1 (1 - t)^2 \partial_j^2 f^i(v_k - t(v_k - x)) dt \\ &\leq 2|v_k - x|^2 \alpha_{\max} && \text{by (4) and Cauchy–Schwarz} \\ &\leq 2D^2 \alpha_{\max} && \text{because } x \in \sigma. \end{aligned}$$

The function f_L at the point $x = \sum_k \lambda_k v_k$, where $\sum_k \lambda_k = 1$, is by construction

$$\begin{aligned} f_L^i(x) &= \sum_k \lambda_k f^i(v_k) \\ &= \sum_k \lambda_k \left(f^i(x) + \sum_j \partial_j f^i(x)(v_k - x)^j + R(v_k) \right) \tag{by (9)} \end{aligned}$$

$$\begin{aligned}
 &= \sum_k \lambda_k f^i(x) + \sum_j \partial_j f^i(x) \left(\sum_k \lambda_k v_k - x \right)^j + \sum_k \lambda_k R(v_k) \\
 &= f^i(x) + 0 + \sum_k \lambda_k R(v_k).
 \end{aligned}$$

Thanks to the bounds on $R(v_k)$ and Cauchy–Schwarz, one has

$$|f_L^i(x) - f^i(x)| \leq 2D^2\alpha_{\max}.$$

□

We will also be using an estimate similar to Proposition 2.2 of Allgower and George [11].

Proposition 10 *Let $\sigma \subset \mathcal{T}_0$ and let f_L be as described above. Then*

$$|\nabla f_L^i(x) - \nabla f^i(x)| = \sqrt{\sum_j (\partial_j f_L^i(x) - \partial_j f^i(x))^2} \leq \frac{4dD\alpha_{\max}}{T},$$

for all x in the simplex σ .

We provide a proof for completeness. In this proof, we use the following variant of a common refinement of two sets:

Claim 11 *Suppose we are given two finite sets $A = \{a_i \mid i = 0, \dots, i_{\max}\}$, $B = \{b_j \mid j = 0, \dots, j_{\max}\}$ of positive numbers such that $\sum a_i = \sum b_j$. Then, there exists a set $C = \{c_k \mid k = 0, \dots, k_{\max}\}$ of positive integers $0 = k_0 \leq k_1 \leq \dots \leq k_{i_{\max}}$, $0 = k'_0 \leq k'_1 \leq \dots \leq k'_{j_{\max}}$ such that*

$$\begin{aligned}
 \sum_{k=k_i}^{k_{i+1}-1} c_k &= a_i & \sum_{k=k'_j}^{k'_{j+1}-1} c_k &= b_j.
 \end{aligned}$$

Proof The proof goes via intersection as indicated in Fig. 2. In the figure, the sets A , B , C are represented as a union of intervals of lengths a_i , b_j , c_k , respectively. We observe the following:

- In the figure, the set C is obtained by the common refinement of the intervals. We thus have $k_{\max} \leq i_{\max} + j_{\max}$ with equality if there are no ι, ι' such that $\sum_{i=1}^{\iota} a_i = \sum_{j=1}^{\iota'} b_j$.
- The total length of C is the same as the length of A and B , i.e. $\sum_k c_k = \sum_i a_i = \sum_j b_j$.
- Any interval of A or B is the union of consecutive intervals of C , and the result follows.

□




Fig. 2 The construction of the c_k s from Claim 11

Proof of Proposition 10 We use the same notation as before and again use that

$$f^i(v_k) = f^i(x) + \sum_j \partial_j f^i(x)(v_k - x)^j + R(v_k), \tag{9}$$

with

$$|R(v_k)| \leq 2D^2\alpha_{\max}. \tag{10}$$

Subtracting $f^i(v_l)$ from $f^i(v_k)$ now yields

$$f^i(v_k) - f^i(v_l) = \sum_j \partial_j f^i(x)(v_k - v_l)^j + R(v_k) - R(v_l).$$

Because f_L is the linear interpolation of f , we have

$$f^i(v_k) - f^i(v_l) = \sum_j \partial_j f_L^i(x)(v_k - v_l)^j,$$

and thus

$$\left| \sum_j (\partial_j f_L^i(x) - \partial_j f^i(x))(v_k - v_l)^j \right| \leq |R(v_k) - R(v_l)| \leq 4D^2\alpha_{\max}. \tag{11}$$

Let now u and v be two points of σ . Writing u and v in terms of their barycentric coordinates with respect to the vertices v_k of σ , we have $u = \sum \mu_k v_k$ and $v = \sum v_k v_k$, with $\sum \mu_k = \sum v_k = 1$ and $\mu_k, v_k \geq 0$. We write K for the set of indices k , in particular $u - v = \sum_{k \in K} (\mu_k - v_k)$, K^+ for the set of $k \in K$ such that $\mu_k - v_k \geq 0$, and $K^- = K \setminus K^+$.

We can now apply Claim 11 to the two sets $\{(\mu_k - v_k), k \in K^+\}$ and $\{-(\mu_k - v_k), k \in K^-\}$ since $\sum_{k \in K^+} (\mu_k - v_k) = \sum_{k \in K^-} -(\mu_k - v_k)$. The refined set is denoted by $\{\pi_l | l = 0, \dots, L\}$, where $L \leq |K|$. The refinement associates with each l

a $k^+ \in K^+$ and a $k^- \in K^-$. We will write $l^+ = k^+(l)$ and $l^- = k^-(l)$ to emphasize this dependence. With this notation, we can write

$$u - v = \sum_{l \in L} \pi_l (v_{l^+} - v_{l^-}). \tag{12}$$

By Claim 11, we have

$$\sum_{l \in L} \pi_l = \sum_{k \in K^+} (\mu_k - \nu_k) \leq \sum_{k \in K} \mu_k = 1. \tag{13}$$

We now see that

$$\begin{aligned} & \left| \sum_j (\partial_j f_L^i(x) - \partial_j f^i(x))(u - v)^j \right| \\ &= \left| \sum_j \sum_{l \in L} \pi_l (\partial_j f_L^i(x) - \partial_j f^i(x))(v_{l^+}^j - v_{l^-}^j) \right| && \text{(by (12))} \\ &\leq \left| \left(\sum_{l \in L} \pi_l \right) |R(v_{l^+}) - R(v_{l^-})| \right| && \text{(by the triangle inequality)} \\ &\leq 4D^2\alpha && \text{(by (11) and (13)).} \end{aligned} \tag{14}$$

Because the simplex σ contains a ball of radius the smallest altitude over d centred at its barycentre, that is, TD/d with T the thickness, the vector $u - w$ can be chosen to be any vector of length less than TD/d . In particular, we can choose

$$(u - w)^j = \frac{TD}{d} \frac{(\partial_j f_L^i(x) - \partial_j f^i(x))}{\sqrt{\sum_j (\partial_j f_L^i(x) - \partial_j f^i(x))^2}}.$$

Plugging this choice into (14) gives

$$\frac{TD}{d} \sqrt{\sum_j (\partial_j f_L^i(x) - \partial_j f^i(x))^2} \leq 4D^2\alpha_{\max}$$

so that

$$\sqrt{\sum_j (\partial_j f_L^i(x) - \partial_j f^i(x))^2} \leq \frac{4dD\alpha_{\max}}{T}.$$

□

We stress that the bound in Proposition 10 depends on the quality of the simplices in the ambient triangulation \mathcal{T} but not on the shape of the cells of the PL approximation. This is fortunate since we know ambient triangulations of very good quality (e.g. Coxeter triangulations [43]), while we do not have control on the shapes of the cells of the PL approximation which depend on the way the isomanifold intersects \mathcal{T} .

2.2.2 Applying the Implicit Function Theorem

Let σ be a simplex of \mathcal{T} and let f_L be the linear approximation defined above. We now define a homotopy $F_L : \mathbb{R}^d \times [0, 1] \rightarrow \mathbb{R}^{d-n}$:

$$F_L(x, \tau) = (1 - \tau)f(x) + \tau f_L(x). \tag{15}$$

We intend to show that $F_L^{-1}(0)$ is a manifold in a neighbourhood of $\sigma \times [0, 1]$. This follows from the Implicit function theorem (Theorem 6) provided that, for any point such that $F_L(x, \tau) = 0$ in this neighbourhood, the Jacobian is of maximal rank or, equivalently as recalled above, if and only if the Gram matrix of its columns has nonzero eigenvalues. The following lemma will provide lower bounds on the eigenvalues of this Gram matrix.

We denote by $\nabla_x F_L$ or simply ∇F_L the gradient of the restriction of F_L to the x variable and $\nabla_{x,\tau} F_L$ the gradient of F_L . Note that

$$\nabla_{x,\tau} F_L^i(x, \tau) = \begin{pmatrix} \nabla(f^i(x) + \tau(f_L^i(x) - f^i(x))) \\ f_L^i(x) - f^i(x) \end{pmatrix}. \tag{16}$$

For convenience, we will write $\nabla_{x,\tau} f(x) = \begin{pmatrix} \nabla f^i(x) \\ 0 \end{pmatrix}$.

Lemma 12 *Let $G = \text{Gram}(\nabla f)$ and $\widehat{G} = \text{Gram}(\nabla_{x,\tau} F_L)$,³ and write λ_{\min} and $\widehat{\lambda}_{\min}$ for the smallest absolute values of the eigenvalues of G and \widehat{G} , respectively.*

$$|\widehat{\lambda}_{\min} - \lambda_{\min}| \leq e_L \tag{17}$$

where, assuming $dD < T$, $e_L = O(d^2D/T)$. If \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d ,

$e_L = O(d^{5/2}D)$. The precise expression of e_L is given in (19) and (18).

Proof Let, in addition to the notations of the lemma, $\widehat{G}' = \text{Gram}(\nabla_x F_L)$, λ'_{\min} be the smallest absolute value of the eigenvalues of \widehat{G}' , and write $G_{i,j}$, $\widehat{G}'_{i,j}$ and $\widehat{G}_{i,j}$ for the entries of G , \widehat{G}' and \widehat{G} , respectively. Proposition 10 yields that

$$\begin{aligned} & |\widehat{G}'_{i,j}(x) - G_{i,j}(x)| \\ &= |\nabla(f^i(x) + \tau(f_L(x)^i - f^i(x))) \cdot \nabla(f^j(x) + \tau(f_L(x)^j - f^j(x)))| \end{aligned}$$

³ As a general rule, we put a $\widehat{}$ over quantities that are related to PL functions.

$$\begin{aligned}
 & - \nabla(f^i(x)) \cdot \nabla(f^j(x))| \\
 = & |\nabla(\tau(f_L(x)^i - f^i(x))) \cdot \nabla(f^j(x)) \\
 & + \nabla(\tau(f_L(x)^j - f^j(x))) \cdot \nabla(f^i(x)) \\
 & + \nabla(\tau(f_L(x)^i - f^i(x))) \cdot \nabla(\tau(f_L(x)^j - f^j(x)))| \\
 \leq & 2\gamma_{\max} \frac{4dD\alpha_{\max}}{T} + \left(\frac{4dD\alpha_{\max}}{T}\right)^2 \stackrel{\text{def}}{=} e'_L. \tag{18}
 \end{aligned}$$

The addition of the τ component gives a small extra contribution.

$$\begin{aligned}
 |\widehat{G}_{i,j}(x) - G_{i,j}(x)| & = |\nabla_{x,\tau} F_L^i(x) \cdot \nabla_{x,\tau} F_L^j(x) - \nabla(f^i(x)) \cdot \nabla(f^j(x))| \\
 & = |\nabla(f^i(x) + \tau(f_L^i(x) - f^i(x))) \cdot \nabla(f^j(x) + \tau(f_L^j(x) - f^j(x))) \\
 & \quad + (f_L^i(x) - f^i(x))(f_L^j(x) - f^j(x)) - \nabla(f^i(x)) \cdot \nabla(f^j(x))| \\
 & \leq e'_L + (2D^2\alpha_{\max})^2. \tag{by Lemma 9}
 \end{aligned}$$

Applying Corollary 8, we obtain

$$|\widehat{\lambda}_{\min} - \lambda_{\min}| \geq (d - n)(e'_L + (2D^2\alpha_{\max})^2) \stackrel{\text{def}}{=} e_L. \tag{19}$$

From (18) and assuming $dD < T$, we see that $e'_L = O(dD/T)$. Hence, $e_L = O(d^2D/T)$. For Coxeter triangulations of type \tilde{A}_d , we have $T > \sqrt{\frac{2}{d+2}}$, see [43]. \square

The following corollary follows directly from the previous lemma and the discussion before the lemma.

Corollary 13 $(F_L^{-1}(0))$ is a manifold in a neighbourhood of $\sigma \times [0, 1]$ Under the regularity condition

$$\lambda_{\min} > e_L, \tag{20}$$

the implicit function theorem applies to $F_L(x, \tau)$ inside $\sigma \times [0, 1]$. (In fact, it applies to an open neighbourhood of this set.) It follows that $\{(x, \tau) \mid F_L(x, \tau) = 0\} \cap (\sigma \times [0, 1])$ is a smooth manifold.

2.2.3 Transversality with Regard to the τ -Direction

We now prove that inside each $\sigma \times [0, 1]$ the gradient of τ on $F_L = 0$ is smooth and does not vanish.

We need the following straightforward lemma. We include a proof for completeness.

Lemma 14 Now suppose that $A = (v_i)^t(v_i)$ is a Gram matrix, where (v_i) denotes the matrix whose column are the vectors v_i , that is, $A_{ij} = v_i \cdot v_j$. Similar to before, denote by $\lambda_{\min}(A)$ the smallest absolute value of an eigenvalue of the Gram matrix A . We have that $\sqrt{\lambda_{\min}(A)} \leq |v_k|$, for all k .

Proof We see that

$$\begin{aligned}
 \lambda_{\min} &= \min_{|u|=1} |u^t Au| \\
 &= \min_{|u|=1} |u^t (v_i)^t (v_i) u| \\
 &= \min_{|u|=1} |((v_i)u)^t ((v_i)u)| \\
 &= \min_{|u|=1} |((v_i)u)|^2 \\
 &\leq \min_{u=e_j} |((v_i)u)|^2 \\
 &= \min_j |v_j|^2 \\
 &\leq |v_k|^2.
 \end{aligned}$$

□

We also need to bound the angle of the vectors $\nabla_{x,\tau}(F_L^i)$ and the x plane, that is $\mathbb{R}^d \subset \mathbb{R}^{d+1}$. We recall the definition. If $v \in \mathbb{R}^{d+1}$ is a vector and $\mathcal{E} = \mathbb{R}^d \subset \mathbb{R}^{d+1}$ is the space spanned by the d basis vectors corresponding to the x -directions, the angle between v and \mathcal{E} is

$$\angle(v, \mathcal{E}) = \inf_{w \in \mathcal{E}} \angle(v, w).$$

Lemma 15 *Let \mathcal{E} be as above. We have*

$$\tan \angle(\nabla_{x,\tau} F_L^i, \mathcal{E}) \leq \theta \stackrel{\text{def}}{=} \frac{2D^2 \alpha_{\max}}{\sqrt{\lambda_{\min}} - \frac{4dD\alpha_{\max}}{T}}. \tag{21}$$

In particular, the manifold $F_L^{-1}(0)$ inside $\sigma \times [0, 1]$ is never tangent to the $\tau = c$ planes, where c is a constant, provided that the following transversality condition holds

$$\lambda_{\min} > \left(\frac{4dD\alpha_{\max}}{T} \right)^2. \tag{22}$$

Proof By (16),

the absolute value of the τ -component of $\nabla_{x,\tau} F_L^i$ is $|f_L(x)^i - f^i(x)|$, which is upper bounded by $2D^2 \alpha_{\max}$ (Lemma 9). On the other hand, the norm of the x -component of $\nabla_{x,\tau} F_L^i$ is lower bounded by

$$\sqrt{\lambda_{\min}} - \frac{4dD\alpha_{\max}}{T},$$

as a consequence of Proposition 10 and Lemma 14. The result now follows since $\tan \angle(\nabla_{x,\tau} F_L^i, \mathcal{E})$ is the ratio between the absolute value of the τ -component and the norm of the x -component of $\nabla_{x,\tau} F_L^i$. □

From Corollary 13 and Lemma 15, we immediately deduce

Corollary 16 *Under the regularity and transversality conditions (20) and (22), which both holds for $D/T = O(1/d^2)$, the gradient of τ on $F_L^{-1}(0)$ is smooth and does not vanish inside $\sigma \times [0, 1]$ for any $\sigma \in \mathcal{T}_0$. If \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d , the condition reduces to $D = O(d^{-3/2})$.*

2.3 Global Result

2.3.1 The Non-smooth Implicit Function Theorem

For the global result, we need to recall some definitions and results from non-smooth analysis. We refer to [38] for an extensive introduction.

Definition 17 (Generalized Jacobian, Definition 2.6.1 of [38]) Let $F : \mathbb{R}^{d+1} \rightarrow \mathbb{R}^{d-n}$, where F is assumed to be just Lipschitz. The generalized Jacobian of F at x_0 , denoted by $J_F(x_0)$, is the convex hull of all $(d - n) \times (d + 1)$ -matrices B obtained as the limit of a sequence of the form $J_F(x_i)$, where $x_i \rightarrow x_0$ and F is differentiable at x_i .

Following [38, page 253], we also define:

Definition 18 The generalized Jacobian $J_F(x_0)$ is said to be of maximal rank, provided every matrix in $J_F(x_0)$ is of maximal rank.

Write $\mathbb{R}^{d+1} = \mathbb{R}^{n+1} \times \mathbb{R}^{d-n}$ and denote the coordinates of \mathbb{R}^{d+1} by (x, y) accordingly. Fix a point (a, b) , with $F(a, b) = 0 \in \mathbb{R}^{d-n}$. We now write:

Notation 19 ([38, page 256]) $J_F(x_0, y_0)|_y$ is the set of all $(n + 1) \times (n + 1)$ -matrices M such that, for some $(n + 1) \times (d - n)$ -matrix N , the $(n + 1) \times (d + 1)$ -matrix $[N, M]$ belongs to $J_F(x_0, y_0)$.

With these definitions and notations, we now have:

Theorem 20 (Generalized implicit function theorem [38, page 256]) *Suppose that $J_F(a, b)|_y$ is of maximal rank. Then, there exists an open set $U \subset \mathbb{R}^{n+1}$ containing a such that there exists a Lipschitz function $g : U \rightarrow \mathbb{R}^{d-n}$, such that $g(a) = b$ and $F(x, g(x)) = 0$ for all $x \in U$.*

2.3.2 Applying the Non-smooth Implicit Function Theorem

We recall the definition of F_{PL} :

$$F_{PL}(x, \tau) = (1 - \tau)f(x) + \tau f_{PL}(x). \tag{1}$$

Further recall that the closed star of a vertex v in a simplicial complex is the closure of all simplices in the complex that contain v . We will also be using the following remark often.

Remark 21 Let v be a vertex in \mathcal{T} , $x_1, x_2 \in \text{star}(v)$, then

$$|x_1 - x_2| \leq 2D. \tag{23}$$

We now have

Lemma 22 Let v be a vertex in \mathcal{T} , $x_1, x_2 \in \text{star}(v)$, and $\tau_1, \tau_2 \in [0, 1]$, such that $\nabla_{x,\tau} F_{PL}^i(x_1, \tau_1)$ and $\nabla_{x,\tau} F_{PL}^i(x_2, \tau_2)$ are well defined, then

$$|\nabla_{x,\tau} F_{PL}^i(x_1, \tau_1) - \nabla_{x,\tau} F_{PL}^i(x_2, \tau_2)| \leq g_{PL} \stackrel{\text{def}}{=} 2dD\alpha_{\max} + \frac{8dD\alpha_{\max}}{T} + 4D^2\alpha_{\max}. \tag{24}$$

Proof Because

$$\nabla_{x,\tau} F_{PL}^i(x_1, \tau_1) = \begin{pmatrix} \nabla(f^i(x_1) + \tau_1(f_{PL}^i(x_1) - f^i(x_1))) \\ f_{PL}^i(x_1) - f^i(x_1) \end{pmatrix},$$

we get

$$\begin{aligned} & |\nabla_{x,\tau} F_{PL}^i(x_1, \tau_1) - \nabla_{x,\tau} F_{PL}^i(x_2, \tau_2)| \\ & \leq |\nabla f^i(x_1) - \nabla f^i(x_2)| \\ & \quad + |\nabla f_{PL}^i(x_1) - \nabla f^i(x_1)| + |\nabla f_{PL}^i(x_2) - \nabla f^i(x_2)| \quad \tau_i \in [0, 1] \\ & \quad + |f_{PL}^i(x_1) - f^i(x_1)| + |f_{PL}^i(x_2) - f^i(x_2)| \\ & \leq 2dD\alpha_{\max} + \frac{8dD\alpha_{\max}}{T} + 4D^2\alpha_{\max} \end{aligned}$$

(by (8) and (23), Proposition 10 and Lemma 9).

□

We generalize Lemma 12 as follows.

Lemma 23 Let v be a vertex in \mathcal{T} , $x_1, \dots, x_m \in \text{star}(v)$, and $\tau_1, \dots, \tau_m \in [0, 1]$. We assume that $\nabla_{x,\tau} F_{PL}^i(x_k, \tau_k)$ is well defined for $k = 1, \dots, m$. Define $\widehat{\mathbf{G}} = \text{Gram}(\sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL}^i(x_k, \tau_k))$, where μ_1, \dots, μ_m are positive weights such that $\mu_1 + \dots + \mu_m = 1$, and let $\widehat{\Lambda}_{\min}$ be the smallest modulus of the eigenvalues of $\widehat{\mathbf{G}}$. Then,

$$|\widehat{\Lambda}_{\min} - \widehat{\lambda}_{\min}| \leq e_{PL},$$

where $e_{PL} = O(d^2 D/T)$ (assuming $dD < T$) and is precisely defined in (26). When \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d , $e_{PL} = O(d^{5/2} D)$.

Proof Let $\widehat{\mathbf{G}} = \text{Gram}(\nabla_{x,\tau} F_L(x_0, y_0))$ and let $\widehat{\lambda}_{\min}$ be the smallest modulus of the eigenvalues of $\widehat{\mathbf{G}}$. We claim that the elements of the two matrices $\widehat{\mathbf{G}}$ and $\widehat{\mathbf{G}}$ are pairwise close. Specifically, using the identity $A \cdot B - C \cdot D = A \cdot (B - D) + (A - C) \cdot D$:

$$\begin{aligned} &|\widehat{\mathbf{G}}_{i,j} - \widehat{\mathbf{G}}_{i,j}| \\ &\leq \left| \sum_k \mu_k \nabla_{x,\tau} F_{PL}^i(x_k, \tau_k) \cdot \sum_k \mu_k \nabla_{x,\tau} F_{PL}^j(x_k, \tau_k) \right. \\ &\quad \left. - \sum_k \mu_k \nabla_{x,\tau} F_{PL}^i(x_0, \tau_0) \cdot \sum_k \mu_k \nabla_{x,\tau} F_{PL}^j(x_0, \tau_0) \right| \\ &= \left| \left(\sum_k \mu_k \nabla_{x,\tau} (F_{PL}^i(x_k, \tau_k)) \right) \cdot \left(\sum_k \mu_k (\nabla_{x,\tau} F_{PL}^j(x_k, \tau_k) - \nabla_{x,\tau} F_{PL}^j(x_0, \tau_0)) \right) \right. \\ &\quad \left. + \left(\sum_k \mu_k (\nabla_{x,\tau} F_{PL}^i(x_k, \tau_k) - \nabla_{x,\tau} F_{PL}^i(x_0, \tau_0)) \right) \cdot \left(\sum_k \mu_k \nabla_{x,\tau} (F_{PL}^j(x_0, \tau_0)) \right) \right| \\ &\leq g_{PL} \cdot \sum_k \mu_k \left(\left| \nabla_{x,\tau} F_{PL}^i(x_k, \tau_k) \right| + \left| \nabla_{x,\tau} F_{PL}^j(x_0, \tau_0) \right| \right), \end{aligned}$$

by Cauchy–Schwarz and the triangle inequality

where g_{PL} is given in Lemma 22. It remains to bound $|\nabla_{x,\tau} F_{PL}^i(x_k, \tau_k)|$:

$$\begin{aligned} |\nabla_{x,\tau} F_{PL}^i(x_k, \tau_k)| &\leq |\nabla_x (f^i(x_k) + \tau(f_{PL}^i(x_k) - f^i(x_k)))| + |f_{PL}^i(x_k) - f^i(x_k)| \\ &\leq \gamma_{\max} + \frac{4dD\alpha_{\max}}{T} + 2D^2\alpha_{\max}, \end{aligned} \tag{25}$$

where we used Lemma 9 and Proposition 10. We conclude that

$$|\widehat{\mathbf{G}}_{i,j} - \widehat{\mathbf{G}}_{i,j}| \leq g_{PL}(\gamma_{\max} + \frac{4dD\alpha_{\max}}{T} + 2D^2\alpha_{\max}).$$

Applying Corollary 8, we get

$$|\widehat{\lambda}_{\min} - \widehat{\lambda}_{\min}| \leq (d - n)g_{PL}(\gamma_{\max} + \frac{4dD\alpha_{\max}}{T} + 2D^2\alpha_{\max}) \stackrel{\text{def}}{=} e_{PL} \tag{26}$$

which completes the proof of the lemma. □

From the previous lemmas, we immediately have that,

Corollary 24 ($\{(x, \tau) \mid F_{PL}(x, \tau) = 0\}$ is a manifold) *Under the regularity condition*

$$\lambda_{\min} > e_{PL}, \tag{27}$$

where e_{PL} is defined in Lemma 23, the generalized implicit function theorem, Theorem 20, applies to $F_{PL}(x, \tau) = 0$. In particular, $\{(x, \tau) \mid F_{PL}(x, \tau) = 0\}$ is a manifold.

The second technical step of the proof is now completed.

The third step follows from an application of Lemma 3. The fact that $F_L(x, \tau) = 0$ is a piecewise-smooth manifold and transversality, as proven in Lemma 15, gives that the gradient of τ is a piecewise-smooth vector field whose flow we can integrate to give an ambient isotopy from the zero set of f to that of f_{PL} .

We summarize in a theorem:

Theorem 25 *If the regularity condition (27) and the transversality condition (22) hold, the zero set of f_{PL} is a manifold isotopic to the zero set of f . Note that both conditions hold when $D/T = O(d^{-2})$. For Coxeter triangulation of type \hat{A}_d , the condition reduces to $D = O(d^{-5/2})$.*

2.3.3 Fréchet Distance

To bound the Fréchet distance, denoted by d_F , between the zero sets of $f(x)$ and f_{PL} , it suffices to bound the angle that the gradient of τ (as restricted to $F_L(x, \tau) = 0$) makes with the τ -direction (in \mathbb{R}^{d+1}). We write e_τ for the unit vector in the τ direction (again in \mathbb{R}^{d+1}).

For this, we will use the angle bound of Lemma 15, together with some estimates that are similar in spirit to those in [51, Lemma C.13].

Lemma 26 *For any $w \in \text{span}(\nabla_{x,\tau} F_L^i)$, we have*

$$\cos \angle(w, e_\tau) \leq \frac{\sin(\theta)(\gamma_{\max} + \frac{4dD\alpha_{\max}}{T})}{\sqrt{\lambda_{\min} - e_{PL}}} = O(D^2)$$

where $e_{PL} = O(d^2 D/T)$ is defined in (26) and $\theta = O(D^2)$ is defined in (21).

Proof Write $v_i = \nabla_{x,\tau} F_L^i$ for $i \in [1, d - n]$, and $w = \mu_1 v^1 + \dots + \mu_{d-n} v^{d-n}$ with $\mu_1, \dots, \mu_{d-n} \in \mathbb{R}$. We have $|w|^2 = \sum_{i,j} \mu_i \mu_j v^i \cdot v^j$, and, by definition,

$$\cos \angle(w, e_\tau) = \frac{\sum_i \mu_i v^i \cdot e_\tau}{|w|} \quad \text{and} \quad |w|^2 = \sum_{i,j} \mu_i \mu_j v^i \cdot v^j \geq \hat{\Lambda}_{\min} |\mu|^2, \tag{28}$$

where $\hat{\Lambda}_{\min}$ is defined in Lemma 23. Proposition 10 and $|\nabla(f^i)| \leq \gamma_{\max}$ give

$$|v^i| \leq \gamma_{\max} + \frac{4dD\alpha_{\max}}{T}. \tag{29}$$

Lemma 15 states that

$$\angle(\nabla_{x,\tau} F_L^i, \mathcal{E}) \leq \theta = \arctan \frac{2D^2\alpha_{\max}}{\sqrt{\lambda_{\min} - \frac{4dD\alpha_{\max}}{T}}}.$$

By definition, \mathcal{E} is the space orthogonal to e_τ (with e_τ aligned with the τ direction), so that

$$\cos(\angle \nabla_{x,\tau} F_L^i, e_\tau) = \sin \angle(\nabla_{x,\tau} F_L^i, \mathcal{E}). \tag{30}$$

Hence, by definition of the cosine and (29), we see

$$|v^i \cdot e_\tau| \leq \sin(\theta) \left(\gamma_{\max} + \frac{4dD\alpha_{\max}}{T} \right).$$

Using (28) and Cauchy–Schwarz, we then obtain

$$\cos \angle(w, e_\tau) \leq \frac{|\mu| \sin(\theta) (\gamma_{\max} + \frac{4dD\alpha_{\max}}{T})}{\sqrt{\widehat{\Lambda}_{\min}} |\mu|^2} = \frac{\sin(\theta) (\gamma_{\max} + \frac{4dD\alpha_{\max}}{T})}{\sqrt{\widehat{\Lambda}_{\min}}}.$$

The result now follows thanks to Lemma 23. □

Let F_{PL_0} be the restriction of F_{PL} to $F_{PL}^{-1}(0, \tau)$.

Moreover, let $\nabla_\tau F_{PL_0}$ be the gradient of τ restricted to $F_{PL}^{-1}(0)$, whenever it exists. We want to bound the angle of $\nabla_\tau F_{PL_0}$ and the τ -direction. Because the isotopy is given by the gradient flow and we have a bound on the norm of the gradient, the Fréchet distance is bounded. Specifically, the bound is equal to the norm of the gradient since the time we follow the flow is 1.

There is one subtlety. Because the manifold is only piecewise-smooth, we need to take into account the points where $\nabla_\tau F_{PL_0}$ is not uniquely defined. Because, for each simplex σ , F_L extends to a neighbourhood of $\sigma \times [0, 1]$, there exists a limit of $\nabla_\tau F_{PL_0}(x_i, \tau_i)$ for any sequence (x_i, τ_i) that lies in $\text{int}(\sigma) \times [0, 1]$, where int denotes the interior. This means that, if we bound $\nabla_\tau F_{PL_0}$ for each simplex, we also bound its limits, where the limits are as just described.

Corollary 27 (Bound on the Fréchet distance) *Suppose that the conditions of Theorem 25 are satisfied. Then,*

$$d_F(f^{-1}(0), f_{PL}^{-1}(0)) \leq d_{PL}$$

where $d_{PL} = O(D^2)$ is defined in (31).

Proof Let, as before, $\mathcal{E} = \mathbb{R}^d \subset \mathbb{R}^{d+1}$ be the space spanned by the d basis vectors corresponding to the x -directions.

Lemma 26 gives, for $w \in \text{span}_i(\nabla_{x,\tau}(F^i))$,

$$\cos \angle(w, e_\tau) \leq \frac{\sin(\theta) (\gamma_{\max} + \frac{4dD\alpha_{\max}}{T})}{\sqrt{\lambda_{\min} - e_{PL}}}.$$

Since the tangent space to $F_L = 0$ is normal to $\text{span}_i(\nabla_{x,\tau}(F_L^i))$, the same bound holds for $\sin \angle(\nabla_\tau F_{PL_0}, e_\tau)$. This means that, as $\tau \in [0, 1]$, the distance between the

begin and the end points of the gradient flow, and thus the Fréchet distance, is bounded by $\tan \angle(\nabla_\tau F_{PL_0}, e_\tau)$, that is,

$$d_F(f^{-1}(0), f_{PL}^{-1}(0)) \leq \tan \arcsin \frac{\sin(\theta)(\gamma_{\max} + \frac{4dD\alpha_{\max}}{T})}{\sqrt{\lambda_{\min} - e_{PL}}} \stackrel{\text{def}}{=} d_{PL}. \tag{31}$$

The asymptotic dependence follows because $\sin(\arctan(x)) = \frac{x}{\sqrt{1+x^2}}$, and $\tan(\arcsin(x)) = \frac{x}{\sqrt{1-x^2}}$. □

3 Isomanifolds with Boundary

We will now consider isomanifolds with boundary. By this, we mean that on top of the function $f : \mathbb{R}^d \rightarrow \mathbb{R}^{d-n}$, we will have another function $f_\partial : \mathbb{R}^d \rightarrow \mathbb{R}$ and the set we consider is $M = f^{-1}(0) \cap f_\partial^{-1}([0, \infty))$. This is a manifold with boundary if the gradients of f^i span a $(d - n)$ -dimensional space at each point of $f^{-1}(0)$ and the gradients of f^i and f_∂ span a $(d - n + 1)$ -dimensional space at each point of $\partial M = f^{-1}(0) \cap f_\partial^{-1}(0)$, as a consequence of the submersion theorem.

We will again write f_{PL} for the PL interpolation of f . Similarly, we write $f_{\partial, PL}$ for the PL interpolation of f_∂ .

We prove that, under certain conditions, there is an isotopy from $f^{-1}(0) \cap f_\partial^{-1}([0, \infty))$ to $f_{PL}^{-1}(0) \cap f_{\partial, PL}^{-1}([0, \infty))$. The conditions are very similar to the conditions we have before but, of course, we need to include bounds on the gradient of $f_{\partial, PL}$.

Overview of the Proof

We will again construct an isotopy, but in this case it will consist of two steps, see Fig. 3 for a pictorial overview.

- In the **first step**, we isotope the part of $f^{-1}(0)$ that is far from $f_\partial^{-1}(0)$ to its piecewise-linear approximation, while leaving the part of $f^{-1}(0)$ that is close to $f_\partial^{-1}(0)$ smooth. We will denote the result by $M_1 = (F_{PL,1}(\cdot, 1))^{-1}(0)$.
- In the **second step**, we consider a (small) tubular neighbourhood around $f_\partial^{-1}(0)$ as restricted to M_1 by looking at all $f_\partial^{-1}(\epsilon)$ for $|\epsilon|$ sufficiently small.⁴ We then isotope $M_1 \cap f_\partial^{-1}(\epsilon)$ to its piecewise-linear approximation. Again, the isotopy is chosen in such a way that, for ϵ relatively large, it leaves $M_1 \cap f_\partial^{-1}(\epsilon)$ invariant (for the points such that M_1 is already piecewise-linear). This gives an isotopy of a tubular neighbourhood of $M_1 \cap f_\partial^{-1}(0)$ to its piecewise-linear approximation.

We will first partition the manifold in two parts using a smooth bump function $\phi : \mathbb{R} \rightarrow [0, 1]$, defined so that $\phi(y) = 0$ in a neighbourhood of zero and $\phi(y) = 1$ if $|y| > y_0$, for some $y_0 > 0$. Such bump functions can be easily constructed, see, for

⁴ We stress that ϵ may be negative.




Fig. 3 Top: we see the original isosurface with $f_{\partial}^{-1}(-1/10)$, $f_{\partial}^{-1}(0)$, $f_{\partial}^{-1}(1/10)$, and $f_{\partial}^{-1}(2/10)$ indicated in blue. Bottom left: we see that at the end of Step 1 the neighbourhood of the boundary is intact, while the rest has been isotoped to a piecewise-linear approximation. Bottom right: we have also isotoped the neighbourhood of the boundary to a piecewise-linear approximation by isotoping $f_{\partial}^{-1}(\epsilon)$, to its piecewise-linear approximation for all sufficiently small ϵ (Color figure online)

example, [39, Section 2.2]. We will be using the function $\phi(\sum_i (f^i)^2 + f_{\partial}^2)$ often. In fact, because it is used so often, it will be convenient to introduce the following shorthand

$$|f_B|^2 = \sum_i (f^i)^2 + f_{\partial}^2. \tag{32}$$

The first step will be using the zero set of the following function:

$$F_{PL,1}(x, \tau) = \left(1 - \tau\phi(|f_B|^2)\right) f(x) + \tau\phi(|f_B|^2) f_{PL}(x), \tag{33}$$

on which we will apply the same gradient flow argument as before.

The resulting set M_1 is the same zero set of f_{PL} as before if we stay sufficiently far away from ∂M and the isotopy leaves the manifold invariant close to ∂M . In particular, $\partial M_1 = \partial M$ (Fig. 3).

In the second step, we define an isotopy that will act only on a small neighbourhood of ∂M . Consider the sets $B_1(\epsilon) = M_1 \cap f_\partial^{-1}(\epsilon)$ and, for each of them, define the function

$$\begin{aligned}
 &F_{PL,2,\epsilon} : B_1(\epsilon) \times [0, 1] \rightarrow \mathbb{R}^{d-n+1} : \\
 &F_{PL,2,\epsilon}(x, \tau) = \left(1 - \tau \psi \left(|f_B|^2\right)\right) (F_{PL,1}(x, 1), f_\partial(x) - \epsilon) \\
 &\quad + \tau \psi \left(|f_B|^2\right) (f_{PL}(x), f_{\partial, PL}(x) - \epsilon), \tag{34}
 \end{aligned}$$

where $\psi : \mathbb{R} \rightarrow [0, 1]$ is now a smooth bump function that is 1 in a sufficiently large neighbourhood of zero (somewhat larger than y_0) and zero outside some compact set. Using the result for isomanifolds (with some modifications), we can prove that each individual set $B_1(\epsilon)$ is isotopic to $f_{PL}^{-1}(0) \cap f_{\partial, PL}^{-1}(\epsilon)$ for small ϵ while, for sufficiently large ϵ , it leaves the set invariant.

3.1 Step 1

The proof closely follows the proof for the case without boundary in Sect. 2. The main technical difficulty will be to provide bounds that serve as the counterparts to Lemma 22 for both steps in the proof. To be able to do so, we first need to discuss bounds on the bump functions ϕ and ψ .

3.1.1 Bump Functions

Following [39, Section 2.2], we write,

$$\zeta_1(x) = \begin{cases} 0 & \text{if } x \leq 0 \\ e^{-1/x} & \text{if } x > 0. \end{cases}$$

For $0 < y_1 < y_2$, we write $\zeta_2(x) = \zeta_1(x - y_1)\zeta_1(y_2 - x)$. Then, we define $\phi_l : \mathbb{R} \rightarrow [0, 1]$ by $\phi_l(x) = \int_x^{y_2} \zeta_2(x') dx' / \int_{y_1}^{y_2} \zeta_2(x') dx'$. Finally, define $\phi_b : \mathbb{R} \rightarrow [0, 1]$ by $\phi_b(x) = \phi_l(|x|)$, and let $\phi(x) = 1 - \phi_b(x)$.

Lemma 28 *We have $\phi_b(x) \in [0, 1]$ and, writing $2y_1 = y_2 = y_0$,*

$$\partial_x(\phi_l(x)) \leq 2 \frac{e^{\frac{4}{3(y_2-y_1)}}}{y_2 - y_1} = 4 \frac{e^{\frac{2}{3y_0}}}{y_0} = \gamma\phi. \tag{35}$$

Proof As mentioned, by construction $\phi_b(x) \in [0, 1]$. Because $\partial_x \beta \left(\frac{y_1+y_2}{2}\right) = 0$ and this is the only zero of the derivative in the open interval (y_1, y_2) , we see that

$$\beta(x) \leq \beta \left(\frac{y_1 + y_2}{2}\right) = e^{\frac{4}{y_1-y_2}}.$$

Hence, because $\beta(x) \geq 0$,

$$\int_{y_1}^{y_2} \beta(x) \leq (y_2 - y_1)e^{\frac{4}{y_1 - y_2}}.$$

Because $\beta(x)$ is monotone on $[y_1, \frac{y_1 + y_2}{2}]$, we also have

$$\int_{y_1}^{y_2} \beta(x) \geq \frac{y_2 - y_1}{2} \beta\left(\frac{3}{4}y_1 + \frac{1}{4}y_2\right) = \frac{y_2 - y_1}{2} e^{\frac{16}{3(y_1 - y_2)}}.$$

We now have

$$\begin{aligned} \partial_x(\phi_l(x)) &= \partial_x \left(\int_x^{y_2} \beta(x')dx' / \int_{y_1}^{y_2} \beta(x')dx' \right) \\ &= \beta(x) / \int_{y_1}^{y_2} \beta(x')dx' \\ &\leq \frac{e^{\frac{4}{y_1 - y_2}}}{\frac{y_2 - y_1}{2} e^{\frac{16}{3(y_1 - y_2)}}} \\ &= 2 \frac{e^{\frac{4}{y_1 - y_2} - \frac{16}{3(y_1 - y_2)}}}{y_2 - y_1} \\ &= 2 \frac{e^{\frac{4}{3(y_2 - y_1)}}}{y_2 - y_1}. \end{aligned}$$

□

3.1.2 Inside a Single Simplex

Similar to Corollary 13, we need a condition that ensures that the zero set of $F_{PL,1}^i(x, \tau)$ restricted to $\sigma \times [0, 1]$ is a smooth manifold. In fact, similar to (15), we define

$$\begin{aligned} F_{L,1}^i(x, \tau) &= \left(1 - \tau\phi(|f_B|^2)\right) f^i(x) + \tau\phi(|f_B|^2) f_L^i(x) \\ &= f^i(x) + \tau\phi(|f_B|^2) (f_L^i(x) - f^i(x)), \end{aligned}$$

where ϕ is as defined above. Observe that $F_{L,1}^i(x, \tau)$ can be extended to a neighbourhood of $\sigma \times [0, 1]$.

Remark 29 For the constants, it is better if y_0 can be chosen as large as possible, but we need y_1 to be quite a bit larger than y_0 . In turn, we cannot choose y_1 arbitrarily large because this would mean that the gradient field $\nabla f_\partial|_{f^{-1}(0)}$ (seen as restricted on $f^{-1}(0)$) would never vanish. The latter is in general impossible thanks to the hairy ball theorem [52].

We introduce the following definition that complements Definition 2:

Definition 30

$$\Gamma_{\max}^B = \max_{x \in \mathcal{T}_0} \left| \nabla \left(|f_B|^2 \right) \right| = 2 \max_{x \in \mathcal{T}_0} \left| \sum_l f^l \nabla f^l + f_\partial \nabla f_\partial \right| \tag{36}$$

$$\hat{\lambda}_{\min}^{B_1} = \min_{x \in \mathcal{T}_0} \lambda_{\min}(\widehat{G}^{B_1})(x), \tag{37}$$

where $\widehat{G}^{B_1} = \text{Gram}(\nabla_{x,\tau} F_{L,1})$ and $\lambda_{\min}(A)$ denotes, as before, the smallest absolute value of the eigenvalues of matrix A .

We have then the analogue of Lemma 12:

Lemma 31 *We have,*

$$|\hat{\lambda}_{\min}^{B_1} - \lambda_{\min}| \leq e_L^{B_1}$$

where $e_L^{B_1} = O(d^2 D/T)$ is precisely defined in (41) and (38). For Coxeter triangulations of type \tilde{A}_d , $e_L^{B_1} = O(d^{5/2} D)$.

Proof We start with an estimate on the individual $\nabla_{x,\tau} F_{L,1}^i(x, \tau)$. As noted before, we write for convenience $\nabla_{x,\tau} f(x) = \begin{pmatrix} \nabla f^i(x) \\ 0 \end{pmatrix}$.

$$\begin{aligned} & \left| \nabla_{x,\tau} f(x) - \nabla_{x,\tau} F_{L,1}^i(x, \tau) \right| \\ &= \left| \nabla_{x,\tau} f(x) - \nabla_{x,\tau} \left(f(x) + \tau \phi \left(|f_B|^2 \right) \left(f_L^i(x) - f^i(x) \right) \right) \right| \\ &= \left| -\nabla_{x,\tau} \left(\tau \phi \left(|f_B|^2 \right) \left(f_L^i(x) - f^i(x) \right) \right) \right| \\ &= \left| \begin{pmatrix} -\tau \nabla \left(\phi \left(|f_B|^2 \right) \right) \left(f_L^i(x) - f^i(x) \right) \\ \phi \left(|f_B|^2 \right) \left(f_L^i(x) - f^i(x) \right) \end{pmatrix} + \begin{pmatrix} -\tau \phi \left(|f_B|^2 \right) \nabla \left(f_L^i(x) - f^i(x) \right) \\ 0 \end{pmatrix} \right| \\ &\leq \left| \tau \nabla \left(\phi \left(|f_B|^2 \right) \right) \left(f_L^i(x) - f^i(x) \right) \right| + \left| \tau \phi \left(|f_B|^2 \right) \nabla \left(f_L^i(x) - f^i(x) \right) \right| \\ &\quad + \left| \phi \left(|f_B|^2 \right) \left(f_L^i(x) - f^i(x) \right) \right| \quad \text{(by the triangle inequality)} \\ &\leq \Gamma_{\max}^B \gamma \phi \left| f_L^i(x) - f^i(x) \right| + \left| \phi \left(|f_B|^2 \right) \nabla \left(f_L^i(x) - f^i(x) \right) \right| + \left| f_L^i(x) - f^i(x) \right| \quad \text{(because } \tau \leq 1, (36), (35), \phi \in [0, 1]) \\ &\leq \Gamma_{\max}^B \gamma \phi \left| f_L^i(x) - f^i(x) \right| + \left| \nabla \left(f_L^i(x) - f^i(x) \right) \right| \quad \text{(because } \phi \in [0, 1] + \left| f_L^i(x) - f^i(x) \right|) \\ &\leq 2D^2 \alpha_{\max} + 2\Gamma_{\max}^B \gamma \phi D^2 \alpha_{\max} + \frac{4dD\alpha_{\max}}{T} \stackrel{\text{def}}{=} 2D^2 \alpha_{\max} + e_i^{B_1} \\ & \tag{by Lemma 9 and Proposition 10.} \end{aligned} \tag{38}$$

We now write $G_{i,j}$ and $\widehat{G}_{i,j}^{B_1}$ for the elements (i, j) of G and \widehat{G}^{B_1} , respectively. Expanding yields

$$\begin{aligned} \widehat{G}_{i,j}^{B_1} &= \left(\nabla_{x,\tau} F_{L,1}^i(x, \tau) - \nabla_{x,\tau} f^i(x) + \nabla_{x,\tau} f^i(x) \right) \\ &\quad \cdot \left(\nabla_{x,\tau} F_{L,1}^j(x, \tau) - \nabla_{x,\tau} f^j(x) \right) \end{aligned}$$

$$\begin{aligned}
 &= G_{i,j} + \left(\nabla_{x,\tau} F_{L,1}^i(x, \tau) - \nabla_{x,\tau} f^j(x) \right) \cdot \nabla_{x,\tau} f^i(x) \\
 &\quad + \nabla_{x,\tau} f^j(x) \cdot \left(\nabla_{x,\tau} F_{L,1}^j(x, \tau) - \nabla_{x,\tau} f^j(x) \right) \\
 &\quad + \left(\nabla_{x,\tau} F_{L,1}^i(x, \tau) - \nabla_{x,\tau} f^j(x) \right) \cdot \left(\nabla_{x,\tau} F_{L,1}^j(x, \tau) - \nabla_{x,\tau} f^j(x) \right). \tag{39}
 \end{aligned}$$

We now see by Cauchy–Schwarz, the triangle inequality and Eq. (38) that

$$|\widehat{G}_{i,j}^{B_1} - G_{i,j}| \leq 2\gamma_{\max} \left(2D^2\alpha_{\max} + e_t^{B_1} \right) + \left(2D^2\alpha_{\max} + e_t^{B_1} \right)^2. \tag{40}$$

Corollary 8 now yields $\widehat{\lambda}_{\min}^{B_1} > \lambda_{\min} - e_L^{B_1}$ where

$$e_L^{B_1} \stackrel{\text{def}}{=} (d - n) \left(2\gamma_{\max} \left(2D^2\alpha_{\max} + e_t^{B_1} \right) + \left(2D^2\alpha_{\max} + e_t^{B_1} \right)^2 \right) \tag{41}$$

□

and $e_t^{B_1}$ is defined in (38). The following corollary is then the analogue of Corollary 13:

Corollary 32 ($F_{L,1}^{-1}(0)$ is a manifold) *If the regularity condition*

$$\lambda_{\min} > e_L^{B_1}, \tag{42}$$

holds, then $F_{L,1}^{-1}(0)$ is a smooth manifold inside an ϵ neighbourhood of $\sigma \times [0, 1]$.

3.1.3 Transversality with Regard to the τ -Direction

We note that, similar to Lemma 15, we have

Lemma 33 *Using the notation of Lemma 15, we have*

$$\tan \angle(\nabla_{x,\tau}(F_{L,1}), \Xi) \leq \frac{2D^2\alpha_{\max}}{\sqrt{\lambda_{\min}} - e_t^{B_1}}.$$

Proof The proof is identical to the proof of Lemma 15 with the replacement of $\frac{4dD\alpha_{\max}}{T}$ in the denominator by $e_t^{B_1}$

The latter constant is a consequence of (38). □

Now, similar to Corollary 16, we find that

Corollary 34 (Transversality with respect to τ for Step 1) *Assume that both the regularity condition (42) and the transversality condition*

$$\lambda_{\min} > e_t^{B_1} \tag{43}$$

hold.

Then, inside each $\sigma \times [0, 1]$, the gradient of τ on $F_{P,L,1}^{-1}(0)$ is smooth and does not vanish. Both conditions are satisfied if $D/T = O(d^{-2})$. When \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d , the condition reduces to $D = O(d^{-5/2})$.

3.1.4 Global Result

We now have to prove that $F_{P,L,1}^{-1}(0)$ is a manifold. For this, we again employ the generalized implicit function theorem. But first of all, we need the following bound, which is similar to Lemma 22.

Note that $\nabla_{x,\tau} F^i(x_0, \tau_0)$ is well defined as soon as x_0 lies in the interior of a d -simplex in \mathcal{T} .

Lemma 35 *Assuming that the gradients are well defined, we have*

$$|\nabla_{x,\tau} F_{P,L,1}^i(x_1, \tau_1) - \nabla_{x,\tau} F_{P,L,1}^i(x_2, \tau_2)| \leq g_{P,L}^{B_1},$$

where $g_{P,L}^{B_1} = O(dD/T)$ is precisely defined in (44). When \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d , $g_{P,L}^{B_1} = O(d^{3/2}D)$.

Proof By expansion, we see that

$$\begin{aligned} & |\nabla_{x,\tau} F_{P,L,1}^i(x_1, \tau_1) - \nabla_{x,\tau} F_{P,L,1}^i(x_2, \tau_2)| \\ &= \left| \nabla_{x,\tau} F_{P,L,1}^i(x_1, \tau_1) - \nabla_{x,\tau} f^i(x_1) + \nabla_{x,\tau} f^i(x_1) \right. \\ &\quad \left. - \nabla_{x,\tau} f^i(x_2) + \nabla_{x,\tau} f^i(x_2) - \nabla_{x,\tau} F_{P,L,1}^i(x_2, \tau_2) \right| \\ &\leq \left| \nabla_{x,\tau} F_{P,L,1}^i(x_1, \tau_1) - \nabla_{x,\tau} f^i(x_1) \right| \\ &\quad + \left| \nabla_{x,\tau} f^i(x_1) - \nabla_{x,\tau} f^i(x_2) \right| \\ &\quad + \left| \nabla_{x,\tau} f^i(x_2) - \nabla_{x,\tau} F_{P,L,1}^i(x_2, \tau_2) \right| \quad \text{(by the triangle inequality)} \\ &\leq 2d\alpha_{\max}D + 4D^2\alpha_{\max} + 2e_t^{B_1} \quad \text{(by (23), (8), and (38) twice)} \\ &\stackrel{\text{def}}{=} g_{P,L}^{B_1}. \end{aligned} \tag{44}$$

This completes the proof. □

Suppose $x_0, x_1, \dots, x_m \in \text{star}(v)$, $\tau_0, \dots, \tau_m \in [0, 1]$ and that $\nabla_{x,\tau} F_{P,L,1}^i(x_i, \tau_i)$ is well defined for all i . Further, assume that μ_1, \dots, μ_m are positive weights such that $\mu_1 + \dots + \mu_m = 1$. We write

$$\widehat{\mathbf{G}}^{B_1} = \text{Gram} \left(\sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{P,L,1}(x_k, \tau_k) \right),$$

and $\widehat{\mathbf{\Lambda}}_{\min}^{B_1}$ for the smallest modulus of the eigenvalues of $\widehat{\mathbf{G}}^{B_1}$.

Lemma 36 *We have*

$$|\widehat{\mathbf{A}}_{\min}^{B_1} - \lambda_{\min}| \leq e_{PL}^{B_1}$$

with $e_{PL}^{B_1} = O(d^2 D/T)$ being precisely defined in (47). For Coxeter triangulations of type \widetilde{A}_d , $e_{PL}^{B_1} = O(d^{5/2} D)$.

Proof The proof is more or less the same as the proof of Lemma 23, but with more complicated bounds. We assume that $x_0 \in \text{star}(v)$ and $\tau_0 \in [0, 1]$ are such that $\nabla_{x,\tau} F^i(x_0, \tau_0)$ is well defined (i.e. x_0 lies in the interior of a d -simplex of T). Lemma 31 gives that

$$\widehat{\mathbf{A}}_{\min}^{B_1} > \lambda_{\min} - e_L^{B_1}.$$

Using $\nabla(f^i) \leq \gamma_{\max}$ and (38), we note that

$$\left| \nabla_{x,\tau} F_{PL,1}^i(x_0, \tau_0) \right| \leq \gamma_{\max} + 2D^2\alpha_{\max} + e_t^{B_1}. \tag{45}$$

We want to use Weyl’s bound to determine a bound on the smallest absolute value of the eigenvalues of $\widehat{\mathbf{G}}^{B_1}$. Writing $\widehat{\mathbf{G}}_{i,j}^{B_1}$ and $\widehat{\mathbf{G}}_{i,j}^{B_1}$ for element (i, j) of matrices $\widehat{\mathbf{G}}^{B_1}(x_0, \tau_0)$ and $\widehat{\mathbf{G}}^{B_1}$, respectively, we show that $\widehat{\mathbf{G}}_{i,j}^{B_1}$ and $\widehat{\mathbf{G}}_{i,j}^{B_1}(x_0, \tau_0)$ are pairwise close (compare to (40)).

$$\begin{aligned} & \left| \widehat{\mathbf{G}}_{i,j}^{B_1} - \widehat{\mathbf{G}}_{i,j}^{B_1}(x_0, \tau_0) \right| \\ &= \left| \left(\sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,1}^i(x_k, \tau_k) \right) \cdot \left(\sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,1}^j(x_k, \tau_k) \right) \right. \\ &\quad \left. - \nabla_{x,\tau} F_{PL,1}^i(x_0, \tau_0) \cdot \nabla_{x,\tau} F_{PL,1}^j(x_0, \tau_0) \right| \\ &\leq \left| \nabla_{x,\tau} F_{PL,1}^i(x_0, \tau_0) - \sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,1}^i(x_k, \tau_k) \right| \\ &\quad \cdot \left| \nabla_{x,\tau} F_{PL,1}^j(x_0, \tau_0) \right| \\ &\quad + \left| \nabla_{x,\tau} F_{PL,1}^j(x_0, \tau_0) - \sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,1}^j(x_k, \tau_k) \right| \\ &\quad \cdot \left| \nabla_{x,\tau} F_{PL,1}^i(x_0, \tau_0) \right| \\ &\quad + \left| \nabla_{x,\tau} F_{PL,1}^i(x_0, \tau_0) - \sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,1}^i(x_k, \tau_k) \right| \\ &\quad \cdot \left| \nabla_{x,\tau} F_{PL,1}^j(x_0, \tau_0) - \sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,1}^j(x_k, \tau_k) \right| \end{aligned}$$

$$\leq 2g_{PL}^{B_1} \cdot \left(\gamma_{\max} + 2D^2\alpha_{\max} + e_t^{B_1} + \left(g_{PL}^{B_1} \right)^2 \right)$$

by Lemma 35 and (45). (46)

Using the result of Lemma 31 and invoking Corollary 8 once more gives

$$|\widehat{A}_{\min}^{B_1} - \lambda_{\min}| \leq e_{PL}^{B_1}$$

with

$$e_{PL}^{B_1} \stackrel{\text{def}}{=} e_L^{B_1} - (d - n) \left(2g_{PL}^{B_1} \cdot \left(\gamma_{\max} + 2D^2\alpha_{\max} + e_t^{B_1} \right) + \left(g_{PL}^{B_1} \right)^2 \right). \quad (47)$$

□

Lemma 36 immediately yields that

Corollary 37 ($F_{PL,1}^{-1}(0)$ is a manifold)
If the regularity condition

$$\lambda_{\min} > e_{PL}^{B_1} \quad (48)$$

holds, then the generalized implicit function theorem, Theorem 20, applies to $F_{PL,1}(x, \tau) = 0$. In particular, $F_{PL,1}^{-1}(0)$ is a manifold.

We stress again that, inside the set $\{x | \phi(\sum_i (f^i)^2(x) + f_\delta^2(x)) = 1\}$, the zero set of $F_{PL,1}(x, 1)$ coincides with the zero set of $f_{PL}(x)$.

3.2 Step 2

Before we can proceed, we have to specify the bump function ψ . We suppose that (the constants have not been optimized)

$$\psi(x) = \begin{cases} 1 & \text{if } |x| \leq \frac{101}{100}y_0 \\ 0 & \text{if } |x| \geq 2y_0. \end{cases}$$

In particular,

Definition 38 We pick $\psi(x) = \phi_b(x)$, with the choice $y_1 = \frac{101}{100}y_0$ and $y_2 = 2y_0$.

Remark 39 We stress that the choice of y_1 and y_2 for the function ψ is different from the choice we made for ϕ , in the first step of the proof.

First, we stress that the zero set of $F_{PL,2,\epsilon}(x, 1)$ coincides with the zero set of $(f_{PL}(x), f_{\partial,PL}(x) - \epsilon)$, provided that $\psi(\sum_i f_i(x)^2 + f_\delta(x)^2) = 1$.

Secondly, we now claim the following:

Lemma 40 *The zero set of $F_{PL,2,\epsilon}(x, 1)$ is a subset of the zero set of $f_{PL}(x)$, for each ϵ .*

Proof We focus on the first $d - n$ coordinates of (34). We see that

$$\begin{aligned} & (1 - \psi(|f_B|^2)) F_{PL,1}(x, 1) + \psi(|f_B|^2) f_{PL}(x) \\ &= (1 - \psi(|f_B|^2)) \left((1 - \phi(|f_B|^2)) f(x) + \phi(|f_B|^2) f_{PL}(x) \right) + \psi(|f_B|^2) f_{PL}(x) \\ &= (1 - \psi(|f_B|^2)) (1 - \phi(|f_B|^2)) f(x) + \left((1 - \psi(|f_B|^2)) \phi(|f_B|^2) + \psi(|f_B|^2) \right) f_{PL}(x) \\ &= \left((1 - \psi(|f_B|^2)) \phi(|f_B|^2) + \psi(|f_B|^2) \right) f_{PL}(x), \end{aligned} \tag{49}$$

where we used that

$$\begin{aligned} 1 - \psi(|f_B|^2) &= 0, & \text{if } |f_B|^2 \leq \frac{101}{100} y_0, \\ (1 - \phi(|f_B|^2)) &= 0, & \text{if } |f_B|^2 \geq y_0. \end{aligned}$$

We can further rewrite (49),

$$\begin{aligned} & \left((1 - \psi(|f_B|^2)) \cdot (\phi(|f_B|^2) - 1 + 1) + \psi(|f_B|^2) \right) f_{PL}(x) \\ &= \left((1 - \psi(|f_B|^2)) (\phi(|f_B|^2) - 1) + 1 \right) f_{PL}(x) \\ &= f_{PL}(x), \end{aligned}$$

where we used the same argument as before. □

The technical result that remains to be proven is the counterpart of Theorem 25 for $F_{PL,2,\epsilon}(x, \tau)$ and for each sufficiently small ϵ . To be precise, it suffices for $\epsilon \leq 2y_0$. We remark that it is likely that this bound on ϵ can be improved.

We again follow the same path to prove this result. That is, we first concentrate on a single simplex and prove that inside that simplex the zero set of $F_{PL,2,\epsilon}$ is a smooth manifold on which the gradient of τ as restricted to the manifold does not vanish. We then prove that the zero set of $F_{PL,2,\epsilon}$ is globally a manifold.

3.2.1 Assumptions and Notations

Because we are now faced with both $f(x)$ and $f_\partial(x)$, we need to introduce a bound on how far the gradients of all the entrees of these functions are from being collinear. We write

$$f_B(x) = (f(x), f_\partial(x)). \tag{50}$$

Before we were only interested in the set \mathcal{T}_0 . Similarly here, we sometimes concentrate on a neighbourhood of the zero set of both f_∂ and f . Therefore, we write \mathcal{T}_B for all $\sigma \in \mathcal{T}$ such that $(\sum_l (f^l)^2 + (f_\partial)^2)^{-1}([-2y_0, 2y_0]) \cap \sigma \neq \emptyset$.

We also write $G^B = \text{Gram}(\nabla f_B)$ and λ_{\min}^B for the minimal absolute value of the eigenvalues of G^B , where the minimization is over all simplices in the set $\mathcal{T}_B \cap \mathcal{T}_0$. The restriction to the set $\mathcal{T}_B \cap \mathcal{T}_0$ is important, because if the minimization would be just over \mathcal{T}_0 , G^B would generically be 0 as a consequence of the hairy ball theorem.

We note that by taking gradients the ϵ constant drops from the expression, so that the properties we now define are independent of ϵ . For the lengths of the gradients of f_B , we define,

$$\gamma_{\max}^B = \max_{x \in \mathcal{T}_0} \max_i |\nabla(f_B^i)|, \tag{51}$$

for all $1 \leq i \leq d - n + 1$. Similar to α_{\max} , we define α_{\max}^B as the bound on the operator 2-norm of all Hessians of f_B , that is,

$$\alpha_{\max}^B = \max_{x \in \mathcal{T}_0} \max_i \|\text{Hes}(f_B^i)\|_2 = \max_{x \in \mathcal{T}_0} \max_i \|(\partial_k \partial_l f_B^i)_{k,l}\|_2. \tag{52}$$

We stress that that $\alpha_{\max} \leq \alpha_{\max}^B$.

We use the same notation for the ambient triangulation \mathcal{T} , the lower bound on the thickness of the simplices T and upper bound on the longest edge length D . We also need to introduce a bound on the differential of the bump function ψ . Similar to (35), we define

$$\gamma_\psi = 2 \frac{e^{\frac{4}{3(y_2 - y_1)}}}{y_2 - y_1} = 2 \frac{e^{\frac{4}{3(2y_0 - \frac{101}{100}y_0)}}}{2y_0 - \frac{101}{100}y_0} = \frac{200}{99} \frac{e^{\frac{400}{297y_0}}}{y_0}, \tag{53}$$

because we picked $y_1 = \frac{101}{100}y_0$ and $y_2 = 2y_0$, for ψ , see Definition 38 and Remark 39.

3.2.2 Inside a Single Simplex

Similar to Lemma 31, we now give a condition that ensures that the zero set of $F_{PL,2,\epsilon}(x, \tau)$ is smooth inside $\sigma \times [0, 1]$. In fact, similar to (15), we define

$$\begin{aligned} F_{L,2,\epsilon}(x, \tau) = & \left(1 - \tau \psi(|f_B|^2)\right) (F_{L,1}(x, 1), f_\partial(x) - \epsilon) \\ & + \tau \psi(|f_B|^2) (f_L(x), f_{\partial,L}(x) - \epsilon), \end{aligned}$$

which can be extended to a neighbourhood of $\sigma \times [0, 1]$.

We also write $\widehat{G}^{B_2} = \text{Gram}(\nabla_{x,\tau} F_{PL,2,\epsilon})$ and $\hat{\lambda}_{\min}^{B_2}$ for the smallest absolute value of the eigenvalues of \widehat{G}^{B_2} .

Lemma 41 *For all ϵ*

$$|\hat{\lambda}_{\min}^{B_2} - \lambda_{\min}^B| \leq e_L^{B_2},$$

where $e_L^{B2} = O(d^2D/T)$ is precisely defined in (63). For Coxeter triangulations of type \tilde{A}_d , $e_L^{B2} = O(d^{5/2}D)$.

Proof We start with an estimate on the individual $\nabla_{x,\tau} F_{L,2,\epsilon}^i(x, \tau)$. We will write $(v, w)^i$ for the i th coordinate of the composed vector (v, w) . We now see that

$$\begin{aligned} & \left| \nabla_{x,\tau} f_B^i(x) - \nabla_{x,\tau} F_{L,2,\epsilon}^i(x, \tau) \right| \\ &= \left| \nabla_{x,\tau} f_B^i(x) \right. \\ &\quad - \nabla_{x,\tau} \left((1 - \tau \psi(|f_B|^2)) (F_{L,1}(x, 1), f_\partial(x) - \epsilon)^i \right. \\ &\quad \left. + \tau \psi(|f_B|^2) (f_L(x), f_{\partial,L}(x) - \epsilon)^i \right) \left. \right| \\ &= \left| \nabla_{x,\tau} f_B^i(x) - \nabla_{x,\tau} \left((1 - \tau \psi(|f_B|^2)) \right. \right. \\ &\quad \cdot \left((1 - \phi(|f_B|^2)) f(x) + \phi(|f_B|^2) f_L(x), f_\partial(x) - \epsilon \right)^i \left. \left. \right. \right| \quad (\text{by definition of } F_{L,1}) \\ &\quad + \tau \psi(|f_B|^2) (f_L(x), f_{\partial,L}(x) - \epsilon)^i \left. \right| \\ &= \left| \nabla_{x,\tau} f_B^i(x) - \nabla_{x,\tau} \left((1 - \tau \psi(|f_B|^2)) \right. \right. \\ &\quad \cdot \left(f(x) + \phi(|f_B|^2) (f_L(x) - f(x)), f_\partial(x) - \epsilon \right)^i \left. \left. \right. \right| \\ &\quad + \tau \psi(|f_B|^2) (f_L(x), f_{\partial,L}(x) - \epsilon)^i \left. \right| \\ &= \left| \begin{pmatrix} \nabla f_B^i(x) \\ 0 \end{pmatrix} \right. \\ &\quad - \nabla_{x,\tau} \left((f(x) + \phi(|f_B|^2) (f_L(x) - f(x)), f_\partial(x) - \epsilon)^i \right. \\ &\quad \left. + \tau \psi(|f_B|^2) \right. \\ &\quad \cdot \left. \left. (f_L(x) - f(x) - \phi(|f_B|^2) (f_L(x) - f(x)), f_{\partial,L}(x) - f_\partial(x))^i \right) \right. \left. \right| \\ &= \left| -\nabla_{x,\tau} \left((\phi(|f_B|^2) (f_L(x) - f(x)), 0)^i \right. \right. \quad (\text{by definition of } f_B^i(x)) \\ &\quad \left. \left. + \tau \psi(|f_B|^2) \right. \right. \\ &\quad \cdot \left. \left. ((1 - \phi(|f_B|^2)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_\partial(x))^i \right) \right. \left. \right| \\ &= \left| \begin{pmatrix} -\nabla (\phi(|f_B|^2) (f_L(x) - f(x)), 0)^i \\ 0 \end{pmatrix} \right. \\ &\quad \left. + \begin{pmatrix} \tau \nabla (\psi(|f_B|^2) ((1 - \phi(|f_B|^2)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_\partial(x))^i) \\ \psi(|f_B|^2) ((1 - \phi(|f_B|^2)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_\partial(x))^i \end{pmatrix} \right. \left. \right| \\ &\leq \max_j \left| \nabla (\phi(|f_B|^2) (f_L^j(x) - f^j(x))) \right| \quad (\text{by the triangle inequality}) \\ &\quad + \left| \nabla (\psi(|f_B|^2) \right. \\ &\quad \cdot \left. ((1 - \phi(|f_B|^2)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_\partial(x))^i \right) \left. \right| \quad (\text{because } \tau \in [0, 1]) \\ &\quad + \left| (f_L(x) - f(x), f_{\partial,L}(x) - f_\partial(x))^i \right| \quad (\text{because } \phi(y), \psi(y) \in [0, 1], \text{ for all } y) \\ &\leq \max_j \left| \nabla \phi(|f_B|^2) \right| |f_L^j(x) - f^j(x)| \end{aligned}$$

$$\begin{aligned}
 & + \max_j \left| \phi \left(|f_B|^2 \right) \right| \left| \nabla (f_L^j(x) - f^j(x)) \right| \\
 & \qquad \qquad \qquad \text{(by the Leibniz rule, Cauchy–Schwarz, and the triangle inequality)} \\
 & + \left| \nabla \left(\psi \left(|f_B|^2 \right) \right) \right| \\
 & \cdot \left| \left((1 - \phi \left(|f_B|^2 \right)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_{\partial}(x) \right)^i \right| \\
 & + \left| \psi \left(|f_B|^2 \right) \right| \\
 & \cdot \left| \nabla \left(\left((1 - \phi \left(|f_B|^2 \right)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_{\partial}(x) \right)^i \right) \right| \\
 & \qquad \qquad \qquad \text{(by the Leibniz rule, and the triangle inequality)} \\
 & + 2D^2 \alpha_{\max}^B \qquad \qquad \text{(by Lemma 9)}
 \end{aligned}$$

$$\begin{aligned}
 & \leq \max_j \gamma_\phi \Gamma_{\max}^B |f_L^j(x) - f^j(x)| + \max_j \left| \nabla (f_L^j(x) - f^j(x)) \right| \qquad \text{(by Lemmas 28, (35), (36), and since } \phi(y) \in [0, 1]) \\
 & + \gamma_\psi \Gamma_{\max}^B \left| \left((1 - \phi \left(|f_B|^2 \right)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_{\partial}(x) \right)^i \right| \qquad \text{(by Lemma 28, (36), (53))} \\
 & + \left| \nabla \left(\left((1 - \phi \left(|f_B|^2 \right)) (f_L(x) - f(x)), f_{\partial,L}(x) - f_{\partial}(x) \right)^i \right) \right| \qquad \text{(because } \psi(y) \in [0, 1]) \\
 & + 2D^2 \alpha_{\max}^B \\
 & \leq \gamma_\phi \Gamma_{\max}^B 2D^2 \alpha_{\max}^B + \frac{4dD\alpha_{\max}^B}{T} \qquad \text{(by Lemma 9 and Proposition 10)} \\
 & + \gamma_\psi \Gamma_{\max}^B 2D^2 \alpha_{\max}^B \qquad \text{(by Lemma 9 and since } \phi(y) \in [0, 1]) \\
 & + \left| \nabla (f_L(x) - f(x), f_{\partial,L}(x) - f_{\partial}(x))^i \right| \\
 & + \left| \nabla \left(\left(\phi \left(|f_B|^2 \right) (f_L(x) - f(x)), 0 \right)^i \right) \right| \qquad \text{(by the triangle inequality)} \\
 & + 2D^2 \alpha_{\max}^B \\
 & \leq \gamma_\phi \Gamma_{\max}^B 2D^2 \alpha_{\max}^B + \frac{4dD\alpha_{\max}^B}{T} \qquad \text{(because by definition } \alpha_{\max} \leq \alpha_{\max}^B) \\
 & + \gamma_\psi \Gamma_{\max}^B 2D^2 \alpha_{\max}^B \\
 & + \frac{4dD\alpha_{\max}^B}{T} \qquad \text{(by Proposition 10)} \\
 & + \max_j \left| \nabla \left(\phi \left(|f_B|^2 \right) (f_L^j(x) - f^j(x)) \right) \right| \\
 & + 2D^2 \alpha_{\max}^B \\
 & \leq (\Gamma_{\max}^B (\gamma_\phi + \gamma_\psi) + 1) 2D^2 \alpha_{\max}^B + \frac{8dD\alpha_{\max}^B}{T} \\
 & + \max_j \left| \nabla \phi \left(|f_B|^2 \right) \right| \left| (f_L^j(x) - f^j(x)) \right| \\
 & + \max_j \left| \phi \left(|f_B|^2 \right) \right| \left| \nabla (f_L^j(x) - f^j(x)) \right| \\
 & \qquad \qquad \qquad \text{(By the Leibniz rule and the triangle inequality)} \\
 & \leq (\Gamma_{\max}^B (\gamma_\phi + \gamma_\psi) + 1) 2D^2 \alpha_{\max}^B + \frac{8dD\alpha_{\max}^B}{T} \\
 & + \gamma_\phi \Gamma_{\max}^B 2D^2 \alpha_{\max}^B \\
 & \qquad \qquad \qquad \text{(by Lemma 28, (35), (36), and Lemma 9)} \\
 & + \frac{4dD\alpha_{\max}^B}{T} \qquad \text{because } \phi(y) \in [0, 1], \text{ (Proposition 10, and } \alpha_{\max} \leq \alpha_{\max}^B) \\
 & = (\Gamma_{\max}^B (2\gamma_\phi + \gamma_\psi) + 1) 2D^2 \alpha_{\max}^B + \frac{12dD\alpha_{\max}^B}{T} \stackrel{\text{def}}{=} e_t^{B2}. \qquad \qquad \qquad (54)
 \end{aligned}$$

We now write $G^B = \text{Gram}(\nabla f_B)$, $\widehat{G}^{B_2} = \text{Gram}(\nabla F_{L,2,\epsilon})$, and $G^{B_{i,j}}$ and $\widehat{G}_{i,j}^{B_2}$ for their (i, j) th elements, respectively. Similar to (39) (we simply need to replace $F_{L,1}$ by $F_{L,2,\epsilon}$ and f by f_B), we obtain

$$|\widehat{G}_{i,j}^{B_2} - G^{B_{i,j}}| \leq 2\gamma_{\max}^B e_t^{B_2} + (e_t^{B_2})^2. \tag{55}$$

By Corollary 8, we finally obtain

$$|\widehat{\lambda}_{\min}^{B_2} - \lambda_{\min}^B| \geq (d - n) \left(2\gamma_{\max}^B e_t^{B_2} + (e_t^{B_2})^2 \right) \stackrel{\text{def}}{=} e_L^{B_2} \tag{56}$$

□

We again have the following corollary.

Corollary 42 ($F_{L,2,\epsilon}^{-1}(0)$ is a manifold) *We have that $F_{L,2,\epsilon}^{-1}(0)$ is a smooth manifold inside an small neighbourhood of $\sigma \times [0, 1]$ provided that the following regularity condition holds*

$$\lambda_{\min}^B > e_L^{B_2}, \tag{57}$$

where $e_L^{B_2}$ is defined in Lemma 41.

3.2.3 Transversality with Regard to the τ -Direction

Once more, similar to Lemma 15 we have

Lemma 43 *Let \mathcal{E} be as in Lemma 15. We have*

$$\tan \angle(\nabla_{x,\tau}(F_{L,2,\epsilon}), \mathcal{E}) \leq \frac{2D^2 \alpha_{\max}^B}{\sqrt{\lambda_{\min}^B - e_t^{B_2}}}$$

where $e_t^{B_2} = O(dD/T)$ ($e_t^{B_2} = O(d^{3/2}D)$) if \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d is precisely defined in (54). In particular, if the following transversality condition holds:

$$\sqrt{\lambda_{\min}^B} > e_t^{B_2} \tag{58}$$

the manifold $F_{L,2,\epsilon}^{-1}(0)$ inside $\sigma \times [0, 1]$, if well defined, is never tangent to the $\tau = c$ planes, where c is a constant.

Proof The proof is identical to the proof of Lemma 15 with the replacement of α_{\max} by α_{\max}^B and of $\frac{4dD\alpha_{\max}}{T}$ in the denominator by $e_t^{B_2}$. The latter constant is a consequence of (54). □

Now, similar to Corollary 34, we have

Corollary 44 (Transversality with respect to τ for Step 2) *If both the regularity condition (57) and the transversality condition (58) hold, then, inside each $\sigma \times [0, 1]$, the gradient of τ on $F_{L,2,\epsilon}^{-1}(0)$ is smooth and does not vanish. Both conditions hold if $D = O(1/d^2)$.*

3.2.4 Global Result

We now have to prove that $F_{PL,2,\epsilon}^{-1}(0)$ is a manifold, for all sufficiently small ϵ . For this we first need the following bound, which is similar to the one in Lemma 35.

Lemma 45 *Let v be a vertex in \mathcal{T} , $x_1, x_2 \in \text{star}(v)$, and $\tau_1, \tau_2 \in [0, 1]$, such that $\nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_1, \tau_1)$ and $\nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_2, \tau_2)$ are well defined, then*

$$|\nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_1, \tau_1) - \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_2, \tau_2)| \leq g_{PL}^{B_2},$$

where $g_{PL}^{B_2} = \mathcal{O}(dD/T)$ is precisely defined in (59). If \mathcal{T} is a Coxeter triangulation of type \tilde{A}_d , $g_{PL}^{B_2} = \mathcal{O}(d^{3/2}D)$.

Proof The proof follows the same steps as the proof of Lemma 35. By expansion, we see that

$$\begin{aligned} & |\nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_1, \tau_1) - \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_2, \tau_2)| \\ &= \left| \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_1, \tau_1) - \nabla_{x,\tau} f_B^i(x_1) \right. \\ &\quad + \nabla_{x,\tau} f_B^i(x_1) - \nabla_{x,\tau} f_B^i(x_2) \\ &\quad \left. + \nabla_{x,\tau} f_B^i(x_2) - \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_2, \tau_2) \right| \\ &\leq \left| \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_1, \tau_1) - \nabla_{x,\tau} f_B^i(x_1) \right| \\ &\quad + \left| \nabla_{x,\tau} f_B^i(x_1) - \nabla_{x,\tau} f_B^i(x_2) \right| \\ &\quad + \left| \nabla_{x,\tau} f_B^i(x_2) - \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_2, \tau_2) \right| \quad \text{(by the triangle inequality)} \\ &\leq 2d\alpha_{\max}^B D + 2e_t^{B_2} \quad \text{(by (8), } |x_1 - x_2| \leq 2D, \text{ and (54) twice)} \\ &\stackrel{\text{def}}{=} g_{PL}^{B_2}. \end{aligned} \tag{59}$$

□

Suppose $x_0, x_1, \dots, x_m \in \text{star}(v)$, $\tau_0, \dots, \tau_m \in [0, 1]$, and that, for all i , $\nabla_{x,\tau} F_{PL,2}^i(x_i, \tau_i)$ is well defined. Further assume that μ_1, \dots, μ_m are positive weights such that $\mu_1 + \dots + \mu_m = 1$. We write

$$\widehat{\mathbf{G}}^{B_2} = \text{Gram} \left(\sum_{k=1}^m \mu_k \nabla_{x,\tau} F_{PL,2}(x_k, \tau_k) \right), \tag{60}$$

and $\widehat{\mathbf{A}}_{\min}^{B_2}$ for the smallest modulus of the eigenvalues of $\widehat{\mathbf{G}}^{B_2}$.

Lemma 46

$$|\widehat{\mathbf{A}}_{\min}^{B_2} - \lambda_{\min}^B| \leq e_{PL}^{B_2} \tag{61}$$

where $e_{PL}^{B_2} = O(d^2 D/T)$ is precisely defined in (63). If \mathcal{T} is a Coxeter triangulation of type \widehat{A}_d , $e_{PL}^{B_2} = O(d^{5/2} D)$.

Proof The proof is more or less the same as the proof of Lemma 36. Let $x_0 \in \text{star}(v)$ and $\tau_0 \in [0, 1]$, be such that $\nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_0, \tau_0)$ is well defined. Note that it is sufficient for x_0 to lie in the interior of a d -simplex in \mathcal{T} . Lemma 41 gives that

$$|\widehat{\mathbf{A}}_{\min}^{B_2} - \lambda_{\min}^B| \leq e_L^{B_2}.$$

Using $\nabla(f_B^i) \leq \gamma_{\max}^B$ and (54), we get

$$\left| \nabla_{x,\tau} F_{PL,2,\epsilon}^i(x_0, \tau_0) \right| \leq \gamma_{\max}^B + e_t^{B_2}. \tag{62}$$

We want to bound the smallest absolute value of the eigenvalues of $\widehat{\mathbf{G}}^{B_2} = \text{Gram}(\nabla_{x,\tau} F_{L,2,\epsilon})$. We proceed similarly to (46) (with $F_{PL,1}$ replaced by $F_{PL,2,\epsilon}^i$). Let $\widehat{\mathbf{G}}^{B_2}$ be as in (60), and denote by $\widehat{\mathbf{G}}_{i,j}^{B_2}$ and $\widehat{\mathbf{G}}_{i,j}^{B_2}$ the (i, j) elements of $\widehat{\mathbf{G}}^{B_2}$ and $\widehat{\mathbf{G}}^{B_2}$, respectively, and by $\widehat{\mathbf{A}}_{\min}^{B_2}$ the smallest absolute value of the eigenvalues of $\widehat{\mathbf{G}}^{B_2}$.

$$|\widehat{\mathbf{G}}_{i,j}^{B_2} - \widehat{\mathbf{G}}_{i,j}^{B_2}| \leq 2 \left(g_{PL}^{B_2} \right) \cdot (\gamma_{\max}^B + e_t^{B_2} + (g_{PL}^{B_2})^2) \tag{by (62)}.$$

Thanks to Corollary 8 and Lemma 41, we have that

$$\begin{aligned} |\widehat{\mathbf{A}}_{\min}^{B_2} - \lambda_{\min}^B| &\leq e_L^{B_2} + (d - n)(2g_{PL}^{B_2}) \cdot (\gamma_{\max}^B + e_t^{B_2} + (g_{PL}^{B_2})^2) \\ &\stackrel{\text{def}}{=} e_{PL}^{B_2}. \end{aligned} \tag{63}$$

□

Lemma 46 immediately yields that

Corollary 47 (The generalized implicit function theorem in Step 2) *If the regularity condition*

$$\lambda_{\min}^B > e_{PL}^{B_2}, \tag{64}$$

the generalized implicit function theorem, Theorem 20, applies to $F_{PL,1}(x, \tau) = 0$. In particular, $F_{PL,1}^{-1}(0)$ is a manifold.

Theorem 48 *If the regularity conditions (42) and (64) and the transversality conditions (43) and (58) hold, then $f^{-1}(0) \cap f_{\partial}^{-1}([0, \infty))$ is isotopic to $f_{PL}^{-1}(0) \cap f_{\partial, PL}^{-1}([0, \infty))$. All conditions hold when choosing $D = O(1/d^2)$.*

Proof The proof follows from Corollaries 32, 34, 37, 42, 44 and 47 . □

3.3 Fréchet Distance

The bounds on the Fréchet distance can be achieved in the same way as before.

Theorem 49 *Suppose that the conditions of Theorem 48 are satisfied. Then,*

$$d_F(f^{-1}(0), f_{PL}^{-1}(0)) \leq d_{PL}^B$$

where $d_{PL}^B = O(D^2)$ is defined in (65).

Proof We apply the same argument as in Lemma 26 and Corollary 27, for both steps of the proof. This yields the sum of two terms that are of the same form as (31). For the first step, we need the following substitutions:

- θ is replaced by $\theta_1 \stackrel{\text{def}}{=} \arctan \frac{2D^2\alpha_{\max}}{\sqrt{\lambda_{\min} - e_t^{B_1}}}$, as a consequence of Lemma 33.
- $\lambda_{\min} - e_{PL}$ is replaced by $\lambda_{\min} - e_{PL}^{B_1}$, as a consequence of Lemma 36.
- $\gamma_{\max} + \frac{4dD\alpha_{\max}}{T}$ is replaced by $\gamma_{\max} + 2D^2\alpha_{\max} + e_t^{B_1}$, as a consequence of (38).

For the second step, we need the following substitutions:


- θ is replaced by $\theta_2 \stackrel{\text{def}}{=} \arctan \frac{2D^2\alpha_{\max}^B}{\sqrt{\lambda_{\min}^B - e_t^{B_2}}}$, as a consequence of Lemma 43.
- $\sqrt{\lambda_{\min} - e_{PL}}$ is replaced by $\sqrt{\lambda_{\min} - e_{PL}^{B_2}}$, as a consequence of Lemma 46.
- $\gamma_{\max} + \frac{4dD\alpha_{\max}}{T}$ is replaced by $\gamma_{\max}^B + e_t^{B_2}$ as a consequence of (54).

This yields

$$\begin{aligned} & d_F(f^{-1}(0) \cap f_{\partial}^{-1}([0, \infty)), f_{PL}^{-1}(0) \cap f_{\partial, PL}^{-1}([0, \infty))) \\ & \leq \tan \arcsin \frac{\sin(\theta_1)(\gamma_{\max} + 2D^2\alpha_{\max} + e_t^{B_1})}{\sqrt{\lambda_{\min} - e_{PL}^{B_1}}} \\ & \quad + \tan \arcsin \frac{\sin(\theta_2)(\gamma_{\max}^B + e_t^{B_2})}{\sqrt{\lambda_{\min} - e_{PL}^{B_2}}} \\ & \stackrel{\text{def}}{=} d_{PL}^B. \end{aligned} \tag{65}$$

□

Fig. 4 An example of an isostratifold



4 Isostratifolds

There is no obstruction that prevents us from extending the approach above to isostratifolds. By isostratifolds, we mean stratifolds that are given by the zero sets of functions and inequalities. For example, suppose that we want to find a PL approximation of the unit sphere centred at 0 in \mathbb{R}^3 including the PL approximations of the intersections of the sphere with slightly deformed $x = 0$, $y = 0$, and $z = 0$ -planes, as depicted in Fig. 4. This would also give PL approximations of the respective ‘octants’ of the sphere.

We could follow the same procedure as for a manifold with boundary to give precise bounds on the longest edge length D of the ambient triangulation that ensure that the PL approximation is correct. However, this would mean that we have to introduce an extra bump function for each stratum as well as an extra isotopy. Even though this should be relatively straightforward, finding the precise constants involved would become prohibitively lengthy.

5 Robustness

Suppose that f and f_δ are smooth functions and moreover f_δ is small in terms of the C^1 -topology. Thanks to the implicit function theorem, we know that if 0 is a regular value of f , the zero set of f and the zero set of the slightly perturbed function $f + f_\delta$ are isotopic. We now give quantitative conditions that guarantee that $f^{-1}(0)$ and $(f + f_\delta)^{-1}(0)$ are ambient isotopic. Let $f, f_\delta : \mathbb{R}^d \rightarrow \mathbb{R}^{d-n}$, and write

$$\tilde{\alpha}_{\max} = \max_x \left(\max_i \{ |\nabla(f_\delta^i)(x)| \}, \max_i |f_\delta^i(x)| \right) \tag{66}$$

$$\tilde{\lambda}_{\min} = \min_{x \in f^{-1}([-\tilde{\alpha}_{\max}, \tilde{\alpha}_{\max}]^{d-n})} \lambda_{\min}(\text{Gram}(\nabla f)) \tag{67}$$

$$\tilde{\gamma}_{\max} = \max_{x \in \mathbb{R}^d} \max_i |\nabla(f^i)|. \tag{68}$$

Theorem 50 *If*

$$\tilde{\lambda}_{\min} > (d - n)(4\tilde{\gamma}_{\max}\tilde{\alpha}_{\max} + \tilde{\alpha}_{\max}^2).$$

and $\sqrt{\tilde{\lambda}_{\min}} > \tilde{\alpha}_{\max}$, then $f^{-1}(0)$ and $(f + f_{\delta})^{-1}(0)$ are ambient isotopic.

Proof We first note that if $|f^i(x)| > \tilde{\alpha}_{\max}$ then $f(x) + \tau f_{\delta}(x) \neq 0$ for all $\tau \in [0, 1]$, so we can restrict our attention to $f^{-1}([-\tilde{\alpha}_{\max}, \tilde{\alpha}_{\max}]^{d-n})$, conform (67). The proof is similar to the proof presented in the previous sections, but much simpler because here all functions are smooth. We start with the function $F(x, \tau) = f(x) + \tau f_{\delta}(x)$, where $\tau \in [0, 1]$. We, again, first establish that the zero set of this function is a $(n + 1)$ -dimensional manifold. Secondly, we will see that the gradient of τ restricted to this $(n + 1)$ -dimensional manifold never vanishes. As we have seen in the previous sections, this suffices to establish the isotopy $F^{-1}(0)$ from $f^{-1}(0)$ to $(f + f_{\delta})^{-1}(0)$, by Lemma 3.

As before, it suffices to prove that $\lambda_{\min}(\nabla_{x,\tau} F) > 0$ to establish that $F^{-1}(0)$ is a manifold. We write

$$\widehat{G} = \text{Gram}(\nabla_{x,\tau} F) \quad \text{and} \quad G = \text{Gram}(\nabla f).$$

We find that

$$\begin{aligned} |\widehat{G}_{i,j} - G_{i,j}| &= |(\nabla f^i + \tau \nabla f_{\delta}^i, f_{\delta}^i) \cdot (\nabla f^j + \tau \nabla f_{\delta}^j, f_{\delta}^j) - \nabla(f^i) \cdot \nabla(f^j)| \\ &= |\tau(\nabla f^i \cdot \nabla f_{\delta}^j + \nabla f_{\delta}^i \cdot \nabla f^j) + f_{\delta}^i f_{\delta}^j| \\ &\leq |\tau| |\nabla f^i| |\nabla f_{\delta}^j| + |\tau| |\nabla f_{\delta}^i| |\nabla f^j| + |f_{\delta}^i| |f_{\delta}^j| \\ &\hspace{10em} \text{(by the triangle inequality and Cauchy–Schwarz)} \\ &\leq 2|\tau| \tilde{\gamma}_{\max} \tilde{\alpha}_{\max} + \tilde{\alpha}_{\max}^2 \hspace{2em} \text{(by (66), (67) and (68))} \\ &\leq 4\tilde{\gamma}_{\max} \tilde{\alpha}_{\max} + \tilde{\alpha}_{\max}^2 \hspace{2em} \text{(because } |\tau| \leq 1). \end{aligned}$$

Corollary 8 implies that $F^{-1}(0)$ is a manifold if

$$\tilde{\lambda}_{\min} > (d - n)(4\tilde{\gamma}_{\max}\tilde{\alpha}_{\max} + \tilde{\alpha}_{\max}^2).$$

Lemma 14 further yields that $|\nabla f^i| \geq \sqrt{\tilde{\lambda}_{\min}}$ in $f^{-1}([-\tilde{\alpha}_{\max}, \tilde{\alpha}_{\max}]^{d-n})$. This means that the x component of $\nabla_{x,\tau}(F)$ does not vanish (again in the domain $f^{-1}([-\tilde{\alpha}_{\max}, \tilde{\alpha}_{\max}]^{d-n})$) as long as

$$\sqrt{\tilde{\lambda}_{\min}} > \tilde{\alpha}_{\max}.$$

□

Acknowledgements First and foremost, we acknowledge Siargey Kachanovich for discussions. We thank Herbert Edelsbrunner and all members of his group, all former and current members of the Datashape team (formerly known as Geometrica), and André Lieutier for encouragement. We further thank the reviewers of Foundations of Computational Mathematics and the reviewers and program committee of the Symposium on Computational Geometry for their feedback, which improved the exposition.

Funding Open access funding provided by Institute of Science and Technology (IST Austria).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

A Notations and Overview of Constants

For notations, we followed the following rules:

1. Greek letters, except τ and ϵ , are for constants related to functions.
2. We use $\hat{\cdot}$ for quantities related to PL functions.
3. Capital letters such as D, T are quantities related to the triangulation \mathcal{T} .
4. Bounds on gradients are denoted by g_x^y .
5. Bounds on eigenvalues are denoted by e_x^y .
6. For convenience, we write $\nabla_{x,\tau} f(x) = \begin{pmatrix} \nabla f^i(x) \\ 0 \end{pmatrix}$.

Overview of Constants

We give an overview. We write \mathcal{T}_0 for the set of all $\sigma \in \mathcal{T}$, such that $(f^i)^{-1}(0) \cap \sigma \neq \emptyset$ for all i . We write \mathcal{T}_B for all $\sigma \in \mathcal{T}$ such that $(\sum_l (f^l)^2 + (f_\partial)^2)^{-1}([-2y_0, 2y_0]) \cap \sigma \neq \emptyset$. We write

$$f_B(x) = (f(x), f_\partial(x)) \tag{50}$$

$$\lambda_{\min} = \min_{x \in \mathcal{T}_0} \min_k |E_k(\nabla(f^i) \cdot \nabla(f^j))_{i,j}|, \tag{3}$$

$$\gamma_{\max} = \max_{x \in \mathcal{T}_0} \max_i |\nabla(f^i)| \tag{2}$$

$$\gamma_{\max}^B = \max_{x \in \mathcal{T}_0} \max_i |\nabla(f_B^i)| \tag{51}$$

$$\Gamma_{\max}^B = \max_{x \in \mathcal{T}_0} \left| \nabla \left(|f_B|^2 \right) \right| = 2 \max_{x \in \mathcal{T}_0} \left| \sum_l f^l \nabla f^l + f_{\partial} \nabla f_{\partial} \right| \quad (36)$$

$$\alpha_{\max} = \max_{x \in \mathcal{T}_0} \max_i \|\text{Hes}(f^i)\|_2 = \max_x \max_i \|(\partial_k \partial_l f^i)_{k,l}\|_2 \quad (4)$$

$$\alpha_{\max}^B = \max_{x \in \mathcal{T}_0} \max_i \|\text{Hes}(f_B^i)\|_2 = \max_{x \in \mathcal{T}_0} \max_i \|(\partial_k \partial_l f_B^i)_{k,l}\|_2. \quad (52)$$

Parameters of the Triangulation

D : the longest edge length of a simplex in \mathcal{T}_0

T : the smallest thickness of a simplex in \mathcal{T}_0

$T > \frac{2}{\sqrt{d+2}}$ for Coxeter triangulations.

Bump Functions

$\mathcal{E} = \mathbb{R}^d \subset \mathbb{R}^{d+1}$ is the space spanned by the d basis vectors corresponding to the x -directions.

$$\gamma_{\phi} = 4 \frac{e^{\frac{2}{3y_0}}}{y_0} \quad (35)$$

$$\gamma_{\psi} = \frac{200}{99} \frac{e^{\frac{400}{297y_0}}}{y_0}. \quad (53)$$

Bounds on Gradients

$$g_{PL} = 2dD\alpha_{\max} + \frac{8dD\alpha_{\max}}{T} + 4D^2\alpha_{\max} = O\left(\frac{dD}{T}\right) \quad (24)$$

$$g_{PL}^{B_1} = 2d\alpha_{\max}D + 2\left((1 + \gamma_{\max}^B\gamma_\phi)2D^2\alpha_{\max} + \frac{4dD\alpha_{\max}}{T}\right) = O\left(\frac{dD}{T}\right) \tag{44}$$

$$g_{PL}^{B_2} = 2d\alpha_{\max}^B D + 2e_t^{B_2} = O\left(\frac{dD}{T}\right). \tag{59}$$

Bounds on Eigenvalues

$$e'_L = 2\gamma_{\max} \frac{4dD\alpha_{\max}}{T} + \left(\frac{4dD\alpha_{\max}}{T}\right)^2 = O\left(\frac{dD}{T}\right) \tag{18}$$

$$e_L = (d - n)(e'_L + (2D^2\alpha_{\max})^2) = O\left(\frac{d^2D}{T}\right) \tag{19}$$

$$e_{PL} = (d - n)g_{PL}(\gamma_1 + \frac{4dD\alpha_{\max}}{T} + 2D^2\alpha_{\max}) = O\left(\frac{d^2D}{T}\right) \tag{26}$$

$$e_t^{B_1} = 2D^2\alpha_{\max} + 2\gamma_{\max}^B\gamma_\phi D^2\alpha_{\max} + \frac{4dD\alpha_{\max}}{T} - 2D^2\alpha_{\max} = O\left(\frac{dD}{T}\right) \tag{38}$$

$$e_L^{B_1} = (d - n)2\gamma_{\max} \left(2D^2\alpha_{\max} + e_t^{B_1}\right) + \left(2D^2\alpha_{\max} + e_t^{B_1}\right)^2 = O\left(\frac{d^2D}{T}\right) \tag{41}$$

$$e_{PL}^{B_1} = e_L^{B_1} - (d - n) \left(2g_{PL}^{B_1} \cdot (\gamma_{\max} + 2D^2\alpha_{\max} + e_t^{B_1}) + (g_{PL}^{B_1})^2\right) = O\left(\frac{d^2D}{T}\right) \tag{47}$$

$$e_t^{B_2} = (\gamma_{\max}^B(2\gamma_\phi + \gamma_\psi) + 1)2D^2\alpha_{\max}^B + \frac{12dD\alpha_{\max}^B}{T} = O\left(\frac{dD}{T}\right) \tag{54}$$

$$e_L^{B_2} = (d - n) \left(2\gamma_{\max}^B e_t^{B_2} + (e_t^{B_2})^2 \right) = O\left(\frac{d^2 D}{T}\right) \quad (56)$$

$$e_{PL}^{B_2} = e_L^{B_2} + (d - n)(2g_{PL}^{B_2}) \cdot \left(\gamma_{\max}^B + e_t^{B_2} + (g_{PL}^{B_2})^2 \right) = O\left(\frac{d^2 D}{T}\right). \quad (63)$$

Bounds on Angles

$$\theta = \frac{2D^2 \alpha_{\max}}{\sqrt{\lambda_{\min}} - \frac{4dD\alpha_{\max}}{T}} = O(D^2). \quad (21)$$

References

1. Timothy S. Newman and Hong Yi. A survey of the marching cubes algorithm. *Computers & Graphics*, 30(5):854–879, 2006.
2. Yen-Chi Chen. Solution manifold and its statistical applications, 2020. 2002.05297.
3. Michael J Todd. *The computation of fixed points and applications*, volume 124 of *Lecture Notes in Economics and Mathematical Systems*. Springer-Verlag, Berlin Heidelberg, 1976.
4. William E. Lorensen and Harvey E. Cline. Marching cubes: A high resolution 3D surface construction algorithm. In *Proceedings of the 14th Annual Conference on Computer Graphics and Interactive Techniques*, SIGGRAPH '87, page 163–169, New York, NY, USA, 1987. Association for Computing Machinery.
5. Akio Doi and Akio Koide. An efficient method of triangulating equi-valued surfaces by using tetrahedral cells. *IEICE TRANSACTIONS on Information and Systems*, E74-D, 1991.
6. Chohong Min. Simplicial isosurfacing in arbitrary dimension and codimension. *Journal of Computational Physics*, 190(1):295–310, 2003.
7. Shek Ling Chan and Enrico O Purisima. A new tetrahedral tessellation scheme for isosurface generation. *Computers & Graphics*, 22(22):83–90, 1998.
8. Graham M Treece, Richard W Prager, and Andrew H Gee. Regularised marching tetrahedra: improved iso-surface extraction. *Computers & Graphics*, 23(4):583–598, 1999.
9. Eugene L. Allgower and Kurt Georg. Simplicial and continuation methods for approximating fixed points and solutions to systems of equations. *SIAM review*, 22(1):28–85, 1980.
10. Gregory M. Nielson and Bernd Hamann. The asymptotic decider: Resolving the ambiguity in marching cubes. In *Proceedings of the 2nd Conference on Visualization '91*, VIS '91, page 83–91, Washington, DC, USA, 1991. IEEE Computer Society Press.
11. Eugene L. Allgower and Kurt Georg. Estimates for piecewise linear approximations of implicitly defined manifolds. *Applied Mathematics Letters*, 2(2):111–115, 1989.
12. Eugene L. Allgower and Kurt Georg. *Numerical continuation methods: an introduction*, volume 13 of *Springer Series in Computational Mathematics*. Springer Science & Business Media, Berlin, Heidelberg, 1990.
13. Hassler Whitney. *Geometric Integration Theory*. Princeton University Press, Princeton, 1957.

14. Jean-Daniel Boissonnat, Siargey Kachanovich, and Mathijs Wintraecken. Triangulating submanifolds: An elementary and quantified version of Whitney's method. *Discrete & Computational Geometry* 66:386–434, 2021.
15. S-W. Cheng, T. K. Dey, and E. A. Ramos. Manifold reconstruction from point samples. In *Proc. 16th ACM-SIAM Symp. Discrete Algorithms*, pages 1018–1027, 2005.
16. Jean-Daniel Boissonnat and Arijit Ghosh. Manifold reconstruction using tangential Delaunay complexes. *Discrete & Computational Geometry*, 51(1):221–267, 2014.
17. Jean-Daniel Boissonnat, Ramsay Dyer, and Arijit Ghosh. Delaunay stability via perturbations. *International Journal of Computational Geometry & Applications*, 24(02):125–152, 2014.
18. Jean-Daniel Boissonnat, Frédéric Chazal, and Mariette Yvinec. *Geometric and Topological Inference*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2018.
19. Jean-Daniel Boissonnat, David Cohen-Steiner, and Gert Vegter. Isotopic implicit surface meshing. *Discrete & Computational Geometry*, 39(1):138–157, 2008.
20. Simon Plantinga and Gert Vegter. Isotopic approximation of implicit curves and surfaces. In *Proceedings of the 2004 Eurographics/ACM SIGGRAPH symposium on Geometry processing*, pages 245–254. ACM, 2004.
21. Jennifer Schultens. *Introduction to 3-manifolds*, volume 151 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, Rhode Island, 2014.
22. J.-D. Boissonnat, R. Dyer, and A. Ghosh. The Stability of Delaunay Triangulations. *International Journal of Computational Geometry & Applications*, 23(4-5):303–334, 2013.
23. Jean-Daniel Boissonnat, Ramsay Dyer, and Arijit Ghosh. Delaunay Triangulation of Manifolds. *Foundations of Computational Mathematics*, 45:38, 2017.
24. Steve Oudot, Laurent Rineau, and Mariette Yvinec. Meshing Volumes Bounded by Smooth Surfaces. *Computational Geometry, Theory and Applications*, 38:100–110, 2007.
25. Laurent Rineau. *Meshing Volumes bounded by Piecewise Smooth Surfaces*. Ph.D. thesis, Université Paris-Diderot - Paris VII, November 2007.
26. T.K. Dey and A. Slatton. Localized Delaunay refinement for piecewise-smooth complexes. *SoCG*, 2013.
27. Tamal K. Dey and Andrew G. Slatton. Localized Delaunay refinement for volumes. *Computer Graphics Forum*, 30(5):1417–1426, 2011.
28. Tamal K. Dey and Joshua A. Levine. Delaunay meshing of piecewise smooth complexes without expensive predicates. *Algorithms*, 2(4):1327–1349, 2009.
29. Paul Bendich, David Cohen-Steiner, Herbert Edelsbrunner, John Harer, and Dmitriy Morozov. Inferring local homology from sampled stratified spaces. In *Proceedings of the IEEE Symposium on Foundations of Computer Science (FoCS)*, pages 536–546, 2007.
30. Paul Bendich, Sayan Mukherjee, and Bei Wang. Stratification learning through homology inference. 2010.
31. Adam Brown and Bei Wang. Sheaf-Theoretic Stratification Learning. In Bettina Speckmann and Csaba D. Tóth, editors, *34th International Symposium on Computational Geometry (SoCG 2018)*, volume 99 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 14:1–14:14, Dagstuhl, Germany, 2018. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
32. T. K. Dey. *Curve and Surface Reconstruction; Algorithms with Mathematical Analysis*. Cambridge University Press, 2007.
33. Siu-Wing Cheng, Tamal K. Dey, and Jonathan R. Shewchuk. *Delaunay Mesh Generation*. CRC Press, 2013.
34. Herbert Edelsbrunner and Nimish R. Shah. Triangulating topological spaces. *International Journal of Computational Geometry & Applications*, 7(04):365–378, 1997.
35. N. Amenta and M. Bern. Surface reconstruction by Voronoi filtering. *Discrete and Computational Geometry*, 22(4):481–504, 1999.
36. Jean-Daniel Boissonnat, Ramsay Dyer, Arijit Ghosh, André Lieutier, and Mathijs Wintraecken. Local conditions for triangulating submanifolds of Euclidean space. 2020.
37. Dominique Attali and André Lieutier. Geometry-driven collapses for converting a Čech complex into a triangulation of a nicely triangulable shape. *Discrete Comput. Geom.*, 54(4):798–825, 2015.
38. Frank H. Clarke. *Optimization and Nonsmooth Analysis*, volume 5 of *Classics in applied mathematics*. SIAM, 1990.
39. M. W. Hirsch. *Differential Topology*. Springer-Verlag, 1976.

40. Jean-Daniel Boissonnat, Sargey Kachanovich, and Mathijs Wintraecken. Tracing isomanifolds in \mathbb{R}^d in time polynomial in d using Coxeter-Freudenthal-Kuhn triangulations. 2021.
41. J. J. Duistermaat and J. A. C. Kolk. *Multidimensional Real Analysis I: Differentiation*. Number 86 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2004.
42. Harold SM Coxeter. Discrete groups generated by reflections. *Annals of Mathematics*, pages 588–621, 1934.
43. Aruni Choudhary, Sargey Kachanovich, and Mathijs Wintraecken. Coxeter triangulations have good quality. *Mathematics in Computer Science*, 14:141–176, 2020.
44. Hans Freudenthal. Simplicialzerlegungen von beschränkter flachheit. *Annals of Mathematics*, pages 580–582, 1942.
45. Harold W Kuhn. Some combinatorial lemmas in topology. *IBM Journal of research and development*, 4(5):518–524, 1960.
46. B Curtis Eaves. *A course in triangulations for solving equations with deformations*, volume 234. Lecture Notes in Economics and Mathematical Systems, 1984.
47. John Milnor. *Morse Theory*. Princeton University Press, Princeton, 1969.
48. John Milnor. *Lectures on the H-Cobordism Theorem*. Princeton University Press, Princeton, 1965.
49. G. H. Golub and C. F. Van Loan. *Matrix computations*. Johns Hopkins Studies in the Mathematical Sciences. The Johns Hopkins University Press, Baltimore, 4 edition, 2013.
50. Rajendra Bhatia. *Matrix analysis*, volume 169 of *Graduate Texts in Mathematics*. Springer Science & Business Media, New York, 2013.
51. J-D. Boissonnat, M. Rouxel-Labbé, and M. Wintraecken. Anisotropic triangulations via discrete Riemannian Voronoi diagrams. *SIAM Journal on Computing*, 48(3):1046–1097, 2019.
52. L. E. J. Brouwer. Über Abbildung von Mannigfaltigkeiten. *Mathematische Annalen*, 71(4):598–598, 1912.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.