



HAL
open science

Duality analysis of interior penalty discontinuous Galerkin methods under minimal regularity and application to the a priori and a posteriori error analysis of Helmholtz problems

Théophile Chaumont-Frelet

► To cite this version:

Théophile Chaumont-Frelet. Duality analysis of interior penalty discontinuous Galerkin methods under minimal regularity and application to the a priori and a posteriori error analysis of Helmholtz problems. 2022. hal-03765207v1

HAL Id: hal-03765207

<https://inria.hal.science/hal-03765207v1>

Preprint submitted on 31 Aug 2022 (v1), last revised 20 Jan 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DUALITY ANALYSIS OF INTERIOR PENALTY DISCONTINUOUS GALERKIN METHODS UNDER MINIMAL REGULARITY AND APPLICATION TO THE A PRIORI AND A POSTERIORI ERROR ANALYSIS OF HELMHOLTZ PROBLEMS

T. CHAUMONT-FRELET*

ABSTRACT. We consider interior penalty discontinuous Galerkin discretizations of time-harmonic wave propagation problems modeled by the Helmholtz equation, and derive novel a priori and a posteriori estimates. Our analysis classically relies on duality arguments of Aubin–Nitsche type, and its originality is that it applies under minimal regularity assumptions. The estimates we obtain directly generalize known results for conforming discretizations, namely that the discrete solution is optimal in a suitable energy norm and that the error can be explicitly controlled by a posteriori estimators, provided the mesh is sufficiently fine.

KEYWORDS. a priori error estimates, a posteriori error estimates, Aubin–Nitsche trick, discontinuous Galerkin, Helmholtz problems, interior penalty, minimal regularity

1. INTRODUCTION

Discontinuous Galerkin methods are a popular approach to discretize PDE boundary value problems [32]. Similar to conforming finite element methods [12, 18], they can handle general meshes, which allows to account for complicated geometries. In addition, the use of discontinuous polynomial shape functions facilitates the presence of hanging nodes and varying polynomial degree. As a result, discontinuous Galerkin methods are especially suited for hp -adaptivity [14, 26]. Furthermore, since all their degrees of freedom are attached with mesh cells, discontinuous Galerkin methods allow linear memory access, which is crucial for efficient computer implementations, in particular on GPUs [6].

In the context of wave propagation problems, discontinuous Galerkin methods further display specific advantages. For time-dependent problems, the resulting mass-matrix is block-diagonal [2, 24], which enables explicit time-stepping schemes without mass-lumping [13]. For time-harmonic wave propagation too, discontinuous Galerkin formulations are attractive as they exhibit additional stability as compared to conforming alternatives [5, 20, 21]. These interesting properties have henceforth motivated a large number of works considering discontinuous Galerkin discretizations of wave propagation problems, and a non-exhaustive list includes [2, 14, 16, 20, 21, 24, 25, 33].

In this work, we consider the acoustic Helmholtz equation, which is probably the simplest model problem relevant to the difficulties of wave propagation. Specifically, given a domain

*University Côte d’Azur, Inria, CNRS, LJAD

$\Omega \subset \mathbb{R}^d$, $d = 2$ or 3 , and $f : \Omega \rightarrow \mathbb{C}$, the unknown $u : \Omega \rightarrow \mathbb{C}$ should satisfy

$$(1.1) \quad \begin{cases} -\omega^2 \mu u - \nabla \cdot (\mathbf{A} \nabla u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_D, \\ \mathbf{A} \nabla u \cdot \mathbf{n} = 0 & \text{on } \Gamma_N, \\ \mathbf{A} \nabla u \cdot \mathbf{n} - i\omega \gamma u = 0 & \text{on } \Gamma_R, \end{cases}$$

where $\overline{\Gamma_D} \cup \overline{\Gamma_N} \cup \overline{\Gamma_R} = \partial\Omega$ is a partition of the boundary and μ , \mathbf{A} and γ are given coefficients. Precise assumptions are listed in Section 2.1.

Our interest lies in interior penalty discontinuous Galerkin (IPDG) discretizations of (1.1). In particular, we focus on the ‘‘minimal regularity’’ case, where we do not assume any specific smoothness for the solution u . To the best of our knowledge, this problem has not been considered in the literature, and available works essentially assume that the solution belongs to $H^{3/2+\varepsilon}$, so that the traces of ∇u are well-defined on mesh faces, see e.g. [25, Eq. (4.5)] and [33, Lemma 2.6]. Unfortunately, such assumptions rule out important configurations of coefficients and boundary conditions, which may bring the regularity of the solution arbitrarily close to H^1 (see the appendix of [15] for instance).

When considering conforming finite element discretizations of (1.1), the so-called ‘‘Schatz argument’’ enables to show that the discrete solution u_h is quasi-optimal if the mesh is fine enough [34]. Assuming for simplicity that $\Gamma_R = \emptyset$ and introducing the energy norm

$$\|v\|_{\omega, \Omega}^2 := \omega^2 \|v\|_{\mu, \Omega}^2 + \|\nabla v\|_{\mathbf{A}, \Omega}^2 \quad v \in H_{\Gamma_D}^1(\Omega),$$

we have

$$(1.2) \quad \|u - u_h\|_{\omega, \Omega} \leq \frac{1}{1 - \gamma_{\text{ba}}^2} \min_{v_h \in V_h} \|u - v_h\|_{\omega, \Omega},$$

whenever the approximation factor

$$(1.3) \quad \gamma_{\text{ba}} := 2\omega \max_{\substack{\psi \in L^2(\Omega) \\ \|\psi\|_{\mu, \Omega} = 1}} \min_{v_h \in V_h} \|u_\psi - v_h\|_{\omega, \Omega}$$

is strictly less than one (all these notations are detailed in Section 2 below). Above, u_ψ solves (1.1) with right-hand side $\mu\psi$ instead of f . Similarly, when considering a posteriori error estimation [9], we have

$$(1.4) \quad \|u - u_h\|_{\omega, \Omega} \leq \sqrt{1 + \gamma_{\text{ba}}^2} \eta,$$

where η is a suitable a posteriori estimator.

Estimates similar to (1.2) and (1.4) are available for IPDG discretizations [28, 33], but with energy norms involving the normal trace of the gradient on faces, thus essentially requiring $H^{3/2+\varepsilon}$ regularity of u_ψ . Here, in contrast, we extend (1.2) and (1.4) to IPDG discretizations without additional regularity assumptions on the solutions u_ψ . As detailed below, our key finding is that it can be achieved by redefining the approximation factor as

$$(1.5) \quad \gamma_{\text{ba}}^2 := 4\omega^2 \max_{\substack{\psi \in L^2(\Omega) \\ \|\psi\|_{\mu, \Omega} = 1}} \left(\min_{v_h \in V_h} \|u_\psi - v_h\|_{1, \dagger, \mathcal{T}_h}^2 + \min_{\mathbf{w}_h \in \mathbf{W}_h} \|\mathbf{A} \nabla u_\psi - \mathbf{w}_h\|_{\text{div}, \dagger, \mathcal{T}_h}^2 \right),$$

where \mathbf{W}_h is the BDM finite element space built using the same mesh and polynomial degree than V_h , and $\|\cdot\|_{1, \dagger, \mathcal{T}_h}$ and $\|\cdot\|_{\text{div}, \dagger, \mathcal{T}_h}$ are $H_{\Gamma_D}^1(\Omega)$ and $\mathbf{H}_{\Gamma_N}(\text{div}, \Omega)$ norms appropriately scaled by the mesh size. The additional term in (1.5) as compared to (1.3) is necessary to account for the non-conformity of the scheme.

Actually, the interest of the subject exceeds time-harmonic wave propagation. In fact, the a priori and a posteriori error analysis of finite element discretizations to (1.1) rely on duality arguments of Aubin-Nitsche type [34]. Such techniques are crucial in time-harmonic wave propagation [29, 33], but there also useful in other contexts to establish convergence in weak norms, see e.g. [3, Section 5.1]. To the best of our knowledge, duality analysis for IPDG discretizations under minimal regularity has not been addressed in the literature, and it is our goal to do so here.

The remainder of this work is organized as follows. In Section 2, we precise the setting and introduce key notations. Section 3 presents the key argument that enables duality techniques for IPDG under minimal regularity. Sections 4 and 5 then employ the aforementioned reasoning to perform the a priori and a posteriori error analysis of IPDG discretization for Helmholtz problems.

2. SETTING AND PRELIMINARY RESULTS

2.1. Setting. Throughout this work, $\Omega \subset \mathbb{R}^d$, with $d = 2$ or 3 , is a Lipschitz polytopal domain. The boundary of Ω is partitioned into three open, Lipschitz and disjoint polytopal subsets Γ_D , Γ_N and Γ_R such that $\partial\Omega = \overline{\Gamma_D} \cup \overline{\Gamma_N} \cup \overline{\Gamma_R}$. We employ the notation \mathbf{n} for the unit vector normal to $\partial\Omega$ pointing outside Ω .

We consider coefficients $\mu : \Omega \rightarrow \mathbb{R}$, $\mathbf{A} : \Omega \rightarrow \mathbb{R}^{d \times d}$ and $\gamma : \Gamma_R \rightarrow \mathbb{R}$ satisfying the following properties. We assume that there exists a partition \mathcal{P} of Ω into a finite number disjoint open polytopal subdomains such that $\mu|_P = \mu_P \in \mathbb{R}$ and $\mathbf{A}|_P = \mathbf{A}_P \in \mathbb{R}^{d \times d}$ take constant values for all $P \in \mathcal{P}$. Similarly, there exists a finite partition \mathcal{Q} of Γ_R consisting of open polytopal subsets such that $\gamma|_Q = \gamma_Q \in \mathbb{R}$ is constant for each $Q \in \mathcal{Q}$. For each $P \in \mathcal{P}$, we assume that \mathbf{A}_P is symmetric, and let $\alpha_P := \min_{\boldsymbol{\xi} \in \mathbb{R}^d; |\boldsymbol{\xi}|=1} \mathbf{A}_P \boldsymbol{\xi} \cdot \boldsymbol{\xi}$. We then classically require that $\min_{P \in \mathcal{P}} \mu_P > 0$, $\min_{P \in \mathcal{P}} \alpha_P > 0$, and $\min_{Q \in \mathcal{Q}} \gamma_Q > 0$.

We also fix a real number $\omega > 0$ representing the (angular) frequency.

2.2. Functional spaces. If $D \subset \mathbb{R}^d$ is an open set, we denote by $L^2(D)$ the Lebesgue space of complex-valued square integrable functions defined over D , and we set $\mathbf{L}^2(D) := [L^2(D)]^d$ for vector-valued functions. The notations $(\cdot, \cdot)_D$ and $\|\cdot\|_D$ then stand for the usual inner-product and norm of $L^2(D)$ or $\mathbf{L}^2(D)$. In addition, if $w : D \rightarrow \mathbb{R}$ is a measurable function satisfying $0 < \text{ess inf}_D w$ and $\text{ess sup}_D w < +\infty$, then $\|\cdot\|_{w,D}^2 := (w \cdot, \cdot)_D$ defines a norm on $L^2(D)$ equivalent to the standard one. We use the same notation in $\mathbf{L}^2(D)$ with matrix-valued weights. Besides, we employ similar notations for $d - 1$ manifolds.

$H^1(D)$ is the Sobolev space of functions $v \in L^2(D)$ such that $\nabla v \in \mathbf{L}^2(D)$, where ∇ denotes the weak gradient defined in the sense of distributions. If $\Gamma \subset \partial D$ is a relatively open subset of the boundary of D , then $H_\Gamma^1(D)$ stands for the subset of functions $v \in H^1(D)$ such that $v|_\Gamma = 0$ in the sense of traces. We refer the reader to [1] for a detailed description of the above spaces. On $H_{\Gamma_D}^1(\Omega)$, we will often employ the following “energy” norm:

$$(2.1) \quad \|v\|_{\omega,\Omega}^2 := \omega^2 \|v\|_{\mu,\Omega}^2 + \omega \|v\|_{\gamma,\Gamma_R}^2 + \|\nabla v\|_{\mathbf{A},\Omega}^2 \quad \forall v \in H_{\Gamma_D}^1(\Omega).$$

It turns out that Sobolev spaces of vector-valued functions will be key in the duality analysis we are about to perform. Specifically, we will need the space $\mathbf{H}(\text{div}, D)$ of functions $\mathbf{v} \in \mathbf{L}^2(D)$ such that $\nabla \cdot \mathbf{v} \in L^2(D)$ where $\nabla \cdot$ is the weak divergence operator [23]. Following, e.g., [22], if $\Gamma \subset \partial D$ is a relatively open set, the normal trace $(\mathbf{w} \cdot \mathbf{n})|_\Gamma$ of $\mathbf{w} \in \mathbf{H}(\text{div}, D)$ can be defined in a weak sense, and $\mathbf{H}_\Gamma(\text{div}, D)$ will stand for the space of $\mathbf{w} \in \mathbf{H}(\text{div}, D)$ such that $(\mathbf{w} \cdot \mathbf{n})|_\Gamma = 0$.

We are studying ‘‘Robin-type’’ boundary conditions which are not naturally handled in the $\mathbf{H}(\operatorname{div})$ setting. We follow the standard remedy [8, 31] and introduce the space

$$\mathbf{X}(\operatorname{div}, \Omega) := \{ \mathbf{w} \in \mathbf{H}(\operatorname{div}, \Omega) \mid (\mathbf{w} \cdot \mathbf{n})|_{\Gamma_R} \in L^2(\Gamma_R) \},$$

where additional normal trace regularity is enforced on the Robin boundary. The notation $\mathbf{X}_{\Gamma_N}(\operatorname{div}, \Omega) := \mathbf{X}(\operatorname{div}, \Omega) \cap \mathbf{H}_{\Gamma_N}(\operatorname{div}, \Omega)$ will also be useful.

2.3. Helmholtz problem. Central to our considerations will be the sesquilinear form

$$(2.2) \quad b(\phi, v) := -\omega^2(\mu\phi, v)_\Omega - i\omega(\gamma\phi, v)_{\Gamma_R} + (\mathbf{A}\nabla\phi, \nabla v)_\Omega \quad \forall \phi, v \in H_{\Gamma_D}^1(\Omega),$$

corresponding to the weak formulation of (1.1). We will assume throughout this work that the considered Helmholtz problem is well-posed, meaning that there exists $\gamma_{\text{st}} > 0$ such that

$$(2.3) \quad \min_{\phi \in H_{\Gamma_D}^1(\Omega)} \max_{\substack{v \in H_{\Gamma_D}^1(\Omega) \\ \|\phi\|_{\omega, \Omega} = 1 \quad \|v\|_{\omega, \Omega} = 1}} \operatorname{Re} b(\phi, v) = \frac{1}{\gamma_{\text{st}}}.$$

In our setting, (2.3) always holds as soon as Γ_R has a positive measure due to the unique continuation principle. On the other hand, if $\Gamma_R = \emptyset$, then (2.3) fails to hold if and only if ω^2 is an eigenvalue of the resulting self-adjoint operator. In general, it is hard to quantitatively estimate γ_{st} , but a reasonable assumption is that it grows polynomially with the frequency [27]. It is also worth mentioning that the constant may be explicitly controlled in some specific configurations [4, 7, 11, 30].

In the remainder of this work, we fix a right-hand side $f \in L^2(\Omega)$, and let $u \in H_{\Gamma_D}^1(\Omega)$ be the unique element satisfying

$$(2.4) \quad b(u, v) = (f, v)_\Omega \quad \forall v \in H_{\Gamma_D}^1(\Omega).$$

2.4. Computational mesh. We consider a mesh \mathcal{T}_h of Ω consisting of non-overlapping (closed) simplicial elements K . We classically assume that \mathcal{T}_h is a matching mesh meaning that the intersection of two distinct elements either is empty, or it is a full face, edge or vertex of the two elements (see, e.g. [12, Section 2.2] or [18, Definition 6.11]). The set of faces of the mesh is denoted by \mathcal{F}_h . We also employ the standard notations h_K and ρ_K for the diameter of $K \in \mathcal{T}_h$ and the diameter of the largest ball contained in K (see [12, Theorem 3.1.3] or [18, Definition 6.4]). Then, $\kappa_K := h_K/\rho_K$ is the shape-regularity parameter of $K \in \mathcal{T}_h$, and $\kappa := \max_{K \in \mathcal{T}_h} \kappa_K$. We also introduce the global mesh size $h := \max_{K \in \mathcal{T}_h} h_K$. Similarly, h_F stands for the diameter of the face $F \in \mathcal{F}_h$.

We further assume that the mesh \mathcal{T}_h conforms with the partition of the boundary and coefficients in the sense that, for each $K \in \mathcal{T}_h$, there exists a (unique) $P \in \mathcal{P}$ such that $K \subset \overline{P}$ and for each $F \in \mathcal{F}_h$ with $F \subset \partial\Omega$, we have either $F \subset \overline{\Gamma_D}$, $F \subset \overline{\Gamma_N}$ or $F \subset \overline{\Gamma_R}$. If $F \subset \overline{\Gamma_R}$, we additionally require that $F \subset \overline{Q}$ for some (unique) $Q \in \mathcal{Q}$. We respectively denote by \mathcal{F}_h^D , \mathcal{F}_h^N and \mathcal{F}_h^R the set of faces $F \in \mathcal{F}_h$ such that $F \subset \overline{\Gamma_D}$, $\overline{\Gamma_N}$ or $\overline{\Gamma_R}$. We also set $\mathcal{F}_h^e := \mathcal{F}_h^D \cup \mathcal{F}_h^N \cup \mathcal{F}_h^R$, and $\mathcal{F}_h^i := \mathcal{F}_h \setminus \mathcal{F}_h^e$. For $K \in \mathcal{T}_h$, we can then set $\mu_K := \mu_P$, $\alpha_K := \alpha_P$, and $\vartheta_K := \sqrt{\alpha_K/\mu_K}$, where $P \in \mathcal{P}$ contains K . Similarly, if $F \in \mathcal{F}_h^e$, we set $\alpha_F := \alpha_K$ where $K \in \mathcal{T}_h$ is the only element having F as a face. If $F \in \mathcal{F}_h^R$, we also introduce $\gamma_F := \gamma_Q$ and $\vartheta_F := \alpha_F/\gamma_F$ where Q is the unique subset in \mathcal{Q} containing F .

Finally, we associate with each $F \in \mathcal{F}_h$ a unit normal vector \mathbf{n}_F . If $F \in \mathcal{F}_h^e$, we require that $\mathbf{n}_F = \mathbf{n}$. Otherwise, the orientation of \mathbf{n}_F is arbitrary, but fixed.

2.5. Polynomial spaces. In the remainder of this work, we fix a polynomial degree $p \geq 1$. For $K \in \mathcal{T}_h$, $\mathcal{P}_p(K)$ is the set of polynomial functions $K \rightarrow \mathbb{C}$ of total degree less than or equal to p . For vector-valued functions, we also set $\mathcal{P}_p(K) := [\mathcal{P}_p(K)]^3$. If $\mathcal{T} \subset \mathcal{T}_h$ is a collection of elements covering the (open) domain Θ , then $\mathcal{P}_p(\mathcal{T}) := \{v \in L^2(\Theta); v|_K \in \mathcal{P}_p(K) \forall K \in \mathcal{T}\}$ and $\mathcal{P}_p(\mathcal{T}) := [\mathcal{P}_p(\mathcal{T})]^3$.

2.6. Broken Sobolev spaces. We define $H^1(\mathcal{T}_h)$ as the subset of functions $v \in L^2(\Omega)$ such that $v|_K \in H^1(K)$ for all $K \in \mathcal{T}_h$. For functions in $H^1(\mathcal{T}_h)$, we still employ the notations ∇ for the element-wise weak gradient, so that $\nabla(H^1(\mathcal{T}_h)) \subset \mathbf{L}^2(\Omega)$. To avoid confusions, we will employ the alternative notations $(\cdot, \cdot)_{\mathcal{T}_h}$ and $\|\cdot\|_{\mathcal{T}_h}$ for the $L^2(\Omega)$ and $\mathbf{L}^2(\Omega)$ norms and inner-products when working with broken functions (we also employ the same notation for weighted norms). Similarly, we write

$$\|v\|_{\gamma, \mathcal{F}_h^R}^2 := \sum_{F \in \mathcal{F}_h^R} \|v\|_{\gamma, F}^2 \quad \forall v \in L^2(\Gamma_R).$$

The notation

$$(\phi, v)_{\mathcal{F}_h} := \sum_{F \in \mathcal{F}_h} (\phi, v)_F \quad \forall \phi, v \in \bigoplus_{F \in \mathcal{F}_h} L^2(F)$$

will also be useful.

2.7. Jumps, averages, lifting and discrete gradient. Considering $\phi \in H^1(\mathcal{T}_h)$, its jump through a face $F \in \mathcal{F}_h^i$ is defined by

$$[[\phi]]_F := \phi_+|_F \mathbf{n}_+ \cdot \mathbf{n}_F + \phi_-|_F \mathbf{n}_- \cdot \mathbf{n}_F$$

with $K_{\pm} \in \mathcal{T}_h$ the two elements such that $F = \partial K_+ \cap \partial K_-$, $\phi|_{\pm} := \phi|_{K_{\pm}}$ and $\mathbf{n}_{\pm} := \mathbf{n}_{K_{\pm}}$. For an exterior face $F \in \mathcal{F}_h^e$, we set instead

$$[[\phi]]_F := \phi|_F \text{ if } F \in \mathcal{F}_h^D \quad \text{and} \quad [[\phi]]_F := 0 \text{ otherwise.}$$

Similarly, if $\mathbf{w} \in \mathcal{P}_p(\mathcal{T}_h)$, its average on $F \in \mathcal{F}_h$ is given by

$$\{\mathbf{w}\}_F := \frac{1}{2}(\mathbf{w}_+|_F + \mathbf{w}_-|_F) \quad \text{if } F \in \mathcal{F}_h^i \quad \text{and} \quad \{\mathbf{w}\}_F := \mathbf{w}|_F \quad \text{if } F \in \mathcal{F}_h^e,$$

where $\mathbf{w}_{\pm} := \mathbf{w}|_{K_{\pm}}$ for the two elements $K_{\pm} \in \mathcal{T}_h$ such that $F = K_- \cap K_+$ in the case of an interior face.

A key part of our analysis will be to give a meaning to the terms $([[\phi]], \{\mathbf{A}\nabla v\} \cdot \mathbf{n}_F)_{\mathcal{F}_h}$ appearing in the IPDG form, for functions ϕ and v only belonging to $H^1(\mathcal{T}_h)$. This is subtle, since the normal trace of ∇v is actually not defined on faces. Following [32, Section 4.3], the solution is to introduce a lifting operator defined as follows.

For $\phi \in H^1(\mathcal{T}_h)$ we define the lifting $\ell_h([[\phi]])$ as the unique element of $\mathcal{P}_p(\mathcal{T}_h)$ such that

$$(\ell_h([[\phi]]), \mathbf{w}_h)_{\mathcal{T}_h} = ([[\phi]], \{\mathbf{w}_h\} \cdot \mathbf{n}_F)_{\mathcal{F}_h} \quad \forall \mathbf{w}_h \in \mathcal{P}_p(\mathcal{T}_h).$$

Notice that we can then write $(\mathbf{A}\ell_h([[\phi]]), \nabla v)_{\mathcal{T}_h}$ instead of $([[\phi]], \{\mathbf{A}\nabla v\} \cdot \mathbf{n}_F)_{\mathcal{F}_h}$ for functions $\phi, v \in \mathcal{P}_p(\mathcal{T}_h)$, with the advantage that the second expression is still well-defined for general $H^1(\mathcal{T}_h)$ arguments.

The notion of weak gradient will also be useful. Specifically, we set

$$\mathfrak{G}(\phi) := \nabla \phi - \ell_h([[\phi]]) \in \mathbf{L}^2(\mathcal{T}_h)$$

for all $\phi \in H^1(\mathcal{T}_h)$. Importantly, we have $\mathfrak{G}(\phi) = \nabla \phi$ whenever $\phi \in H_{\Gamma_D}^1(\Omega)$, and

$$(\mathfrak{G}(\phi), \mathbf{w}_h)_{\mathcal{T}_h} + (\phi, \nabla \cdot \mathbf{w}_h)_{\mathcal{T}_h} = (\phi, \mathbf{w}_h \cdot \mathbf{n})_{\Gamma_R}$$

for all $\phi \in H^1(\mathcal{T}_h)$ and $\mathbf{w}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{X}_{\Gamma_N}(\Omega, \text{div})$.

2.8. Broken norms. For $v \in H^1(\mathcal{T}_h)$, the broken counterpart of the energy norm introduced in (2.1) is defined by

$$\|v\|_{\omega, \mathcal{T}_h}^2 := \omega^2 \|v\|_{\mu, \mathcal{T}_h}^2 + \omega \|v\|_{\gamma, \mathcal{F}_h^R}^2 + \|\mathfrak{G}(v)\|_{\mathbf{A}, \mathcal{T}_h}^2.$$

Notice that it is equal to the $\|\cdot\|_{\omega, \Omega}$ norm if $v \in H_{\Gamma_D}^1(\Omega)$. We will also employ the mesh-dependent norms

$$\|v\|_{\dagger, 1, \mathcal{T}_h}^2 := \sum_{K \in \mathcal{T}_h} \left\{ \max \left(1, \frac{\omega^2 h_K^2}{\vartheta_K^2} \right) \frac{\alpha_K}{h_K^2} \|v\|_K^2 + \|\mathfrak{G}(v)\|_{\mathbf{A}, K}^2 \right\} + \sum_{F \in \mathcal{F}_h^R} \max \left(1, \frac{\omega h_F}{\vartheta_F} \right) \frac{\alpha_F}{h_F} \|v\|_F^2$$

and

$$\|\mathbf{w}\|_{\dagger, \text{div}, \mathcal{T}_h}^2 := \sum_{K \in \mathcal{T}_h} \left\{ \|\mathbf{w}\|_{\mathbf{A}^{-1}, K}^2 + \frac{h_K^2}{\alpha_K} \|\nabla \cdot \mathbf{w}\|_K^2 \right\} + \sum_{F \in \mathcal{F}_h^R} \frac{h_F}{\alpha_F} \|\mathbf{w} \cdot \mathbf{n}\|_F^2$$

for $\mathbf{w} \in \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$. These norms are ‘‘dual’’ to each other in the sense that

$$(2.5) \quad |(\mathfrak{G}(\phi), \mathbf{w})_{\mathcal{T}_h} + (\phi, \nabla \cdot \mathbf{w})_{\mathcal{T}_h} - (\phi, \mathbf{w})_{\mathcal{F}_h^R}| \leq \|\phi\|_{\dagger, 1, \mathcal{T}_h} \|\mathbf{w}\|_{\dagger, \text{div}, \mathcal{T}_h}$$

for all $\phi \in H^1(\mathcal{T}_h)$ and $\mathbf{w} \in \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$. For future references, we notice that the ‘‘max’’ in the definition of the $\|\cdot\|_{\dagger, 1, \mathcal{T}_h}$ are chosen so that

$$(2.6) \quad \|v\|_{\omega, \mathcal{T}_h} \leq \|v\|_{\dagger, 1, \mathcal{T}_h} \quad \forall v \in H^1(\mathcal{T}_h).$$

2.9. IPDG form. Following [32, Section 4.3.3], our DG approximation of the $(\mathbf{A}\nabla \cdot, \nabla \cdot)_{\Omega}$ form is given by

$$(2.7) \quad a_h(\phi, v) := (\mathbf{A}\mathfrak{G}(\phi), \mathfrak{G}(v))_{\mathcal{T}_h} + s_h(\phi, v) \quad \forall \phi, v \in H^1(\mathcal{T}_h)$$

where $s_h : H^1(\mathcal{T}_h) \times H^1(\mathcal{T}_h) \rightarrow \mathbb{C}$ is a sesquilinear form satisfying $s_h(\phi, v) = 0$ whenever $\phi \in H_{\Gamma_R}^1(\Omega)$ or $v \in H_{\Gamma_R}^1(\Omega)$. Importantly, we have

$$(2.8) \quad a_h(\phi, v) = (\mathbf{A}\nabla \phi, \nabla v)_{\Omega}$$

when $\phi, v \in H_{\Gamma_D}^1(\Omega)$.

It is worthy to note that the IPDG form is usually not presented (and not implemented) as it is presented in (2.7). Instead [2, 20, 24, 33], the method is usually written as

$$(2.9) \quad a_h(\phi, v) = (\mathbf{A}\nabla \phi, \nabla v)_{\mathcal{T}_h} - (\{\{\mathbf{A}\nabla \phi\}\} \cdot \mathbf{n}_F, \llbracket v \rrbracket)_{\mathcal{F}_h} - (\llbracket \phi \rrbracket, \{\{\mathbf{A}\nabla v\}\} \cdot \mathbf{n}_F)_{\mathcal{F}_h} + \sum_{F \in \mathcal{F}_h^i \cup \mathcal{F}_h^D} \frac{\beta_F}{h_F} (\llbracket \phi \rrbracket, \llbracket v \rrbracket)_F$$

where $\beta_F \sim p^2$ is a penalty parameter chosen to be large enough [2]. However, as shown in [32, Section 4.3.3], the formulation of (2.9) can be recast in the framework of (2.7) by setting

$$(2.10) \quad s_h(\phi, v) := \sum_{F \in \mathcal{F}_h^i \cup \mathcal{F}_h^D} \frac{\beta_F}{h_F} (\llbracket \phi \rrbracket, \llbracket v \rrbracket)_F - (\mathbf{A}\ell_h(\llbracket \phi \rrbracket), \ell_h(\llbracket v \rrbracket))_{\Omega}.$$

2.10. The discrete Helmholtz problem. Consider the sesquilinear form

$$b_h(\phi, v) := -\omega^2(\mu\phi, v)_{\mathcal{T}_h} - i\omega(\gamma\phi, v)_{\mathcal{F}_h^R} + a_h(\phi, v) \quad \forall \phi, v \in H^1(\mathcal{T}_h).$$

Then, the discrete problem consists in finding $u_h \in \mathcal{P}_p(\mathcal{T}_h)$ such that

$$(2.11) \quad b_h(u_h, v_h) = (f, v_h)_{\mathcal{T}_h} \quad \forall v_h \in \mathcal{P}_p(\mathcal{T}_h).$$

For any u_h satisfying (2.11), recalling (2.4) and due to (2.8), the Galerkin orthogonality property

$$(2.12) \quad b_h(u - u_h, w_h) = 0$$

holds true for all discrete conforming test functions $w_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_{\Gamma_D}^1(\Omega)$. Similarly, we have

$$(2.13) \quad |b_h(\phi, v)| \leq \|\phi\|_{\omega, \mathcal{T}_h} \|v\|_{\omega, \mathcal{T}_h}$$

for $\phi, v \in H^1(\mathcal{T}_h)$ whenever ϕ or v belongs to $H_{\Gamma_D}^1(\Omega)$.

2.11. Conforming subspaces and projections. The sets $\mathcal{P}_p(\mathcal{T}_h) \cap H_{\Gamma_D}^1(\Omega)$ and $\mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$ are the usual Lagrange and BDM finite element spaces (see, e.g., [18, Sections 8.5.1 and 11.5]). Although they are not needed to implement the IPDG discretization, they will be extremely useful for the analysis. The projections

$$\pi_h^g v := \arg \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_{\Gamma_D}^1(\Omega)} \|v - v_h\|_{1, \dagger, \mathcal{T}_h}$$

and

$$\pi_h^d \mathbf{w} := \arg \min_{\mathbf{w}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)} \|\mathbf{w} - \mathbf{w}_h\|_{\text{div}, \dagger, \mathcal{T}_h}$$

are well-defined for all $v \in H_{\Gamma_D}^1(\Omega)$ and $\mathbf{w} \in \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$, since the mesh-dependent norms appearing in the right-hand sides are naturally associated with inner-product. We will also need the conforming projector

$$\pi^g v := \arg \min_{s \in H_{\Gamma_D}^1(\Omega)} \|v - s\|_{1, \dagger, \mathcal{T}_h}$$

defined for all $v \in H^1(\mathcal{T}_h)$.

2.12. Approximation factors. Following [9, 28, 29, 33], our analysis will rely on duality arguments and approximation factors will be a central concept. For $\psi \in L^2(\Omega)$ and $\Psi \in L^2(\Gamma_R)$, let $u_\psi^*, U_\Psi^* \in H_{\Gamma_R}^1(\Omega)$ solve

$$b(w, u_\psi^*) = \omega(\mu w, \psi)_\Omega, \quad b(w, U_\Psi^*) = \omega^{1/2}(\gamma w, \Psi)_{\Gamma_R} \quad \forall w \in H_{\Gamma_R}^1(\Omega).$$

The approximation factors

$$(2.14a) \quad \check{\gamma}_{\text{ba}, g} := \max_{\substack{\psi \in L^2(\Omega) \\ \|\psi\|_{\mu, \Omega} = 1}} \|u_\psi^* - \pi_h^g u_\psi^*\|_{\dagger, 1, \mathcal{T}_h}, \quad \tilde{\gamma}_{\text{ba}, g} := \max_{\substack{\Psi \in L^2(\Gamma_R) \\ \|\Psi\|_{\gamma, \Gamma_R} = 1}} \|U_\Psi^* - \pi_h^g U_\Psi^*\|_{\dagger, 1, \mathcal{T}_h},$$

where previously employed in [9] for the a posteriori error analysis of conforming discretizations. Here, we will additionally need divergence-conforming approximation factors

$$(2.14b) \quad \check{\gamma}_{\text{ba}, d} := \max_{\substack{\psi \in L^2(\Omega) \\ \|\psi\|_{\mu, \Omega} = 1}} \|\mathbf{A}\nabla u_\psi^* - \pi_h^d(\mathbf{A}\nabla u_\psi^*)\|_{\dagger, \text{div}, \mathcal{T}_h}, \\ \tilde{\gamma}_{\text{ba}, d} := \max_{\substack{\Psi \in L^2(\Gamma_R) \\ \|\Psi\|_{\gamma, \Gamma_R} = 1}} \|\mathbf{A}\nabla U_\Psi^* - \pi_h^d(\mathbf{A}\nabla U_\Psi^*)\|_{\dagger, \text{div}, \mathcal{T}_h},$$

to deal with the non-conformity of the IPDG approximation under minimal regularity. We finally introduce the following short-hand notations

$$(2.14c) \quad \gamma_{\text{ba},g}^2 := 4\tilde{\gamma}_{\text{ba},g}^2 + 2\tilde{\gamma}_{\text{ba},g}^2, \quad \gamma_{\text{ba},d}^2 := 4\tilde{\gamma}_{\text{ba},d}^2 + 2\tilde{\gamma}_{\text{ba},d}^2,$$

$$(2.14d) \quad \tilde{\gamma}_{\text{ba}}^2 := \tilde{\gamma}_{\text{ba},g}^2 + \tilde{\gamma}_{\text{ba},d}^2, \quad \tilde{\gamma}_{\text{ba}}^2 := \tilde{\gamma}_{\text{ba},g}^2 + \tilde{\gamma}_{\text{ba},d}^2,$$

and

$$(2.14e) \quad \gamma_{\text{ba}}^2 := \gamma_{\text{ba},g}^2 + \gamma_{\text{ba},d}^2 = 4\tilde{\gamma}_{\text{ba}}^2 + 2\tilde{\gamma}_{\text{ba}}^2.$$

By combining elliptic regularity and approximation properties of finite element spaces, it is easy to see that $\gamma_{\text{ba}} \rightarrow 0$ as $h/p \rightarrow 0$. However, the dependency on the PDE coefficients and on the frequency may be hard to track. In particular, the stability constant γ_{st} typically plays a central role in estimating γ_{ba} . Qualitative estimates for γ_{ba} can be found in [10, 29], and explicit estimates are available in specific situations [9].

3. DUALITY UNDER MINIMAL REGULARITY

Here, we state in a separate section the key technical result that enables us to perform duality techniques under minimal regularity assumptions. We believe it will be important in other contexts, so that we make it easy to reference. Although the arguments are not complicated, we were not able to find a proof in the literature.

Lemma 3.1 (Lifting error). *For all $\phi \in H^1(\mathcal{T}_h)$ and $\boldsymbol{\sigma} \in \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$, we have*

$$(3.1) \quad (\mathfrak{G}(\phi), \boldsymbol{\sigma})_{\mathcal{T}_h} + (\phi, \nabla \cdot \boldsymbol{\sigma})_{\mathcal{T}_h} - (\phi, \boldsymbol{\sigma} \cdot \mathbf{n})_{\mathcal{F}_h^R} = \\ (\mathfrak{G}(\phi - \tilde{\phi}), \boldsymbol{\sigma} - \boldsymbol{\sigma}_h)_{\mathcal{T}_h} + (\phi - \tilde{\phi}, \nabla \cdot (\boldsymbol{\sigma} - \boldsymbol{\sigma}_h))_{\mathcal{T}_h} - (\phi - \tilde{\phi}, (\boldsymbol{\sigma} - \boldsymbol{\sigma}_h) \cdot \mathbf{n})_{\mathcal{F}_h^R}$$

and in particular

$$(3.2) \quad |(\mathfrak{G}(\phi), \boldsymbol{\sigma})_{\mathcal{T}_h} + (\phi, \nabla \cdot \boldsymbol{\sigma})_{\mathcal{T}_h} - (\phi, \boldsymbol{\sigma} \cdot \mathbf{n})_{\mathcal{F}_h^R}| \leq \|\phi - \tilde{\phi}\|_{\dagger, 1, \mathcal{T}_h} \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|_{\dagger, \text{div}, \mathcal{T}_h}$$

for all $\tilde{\phi} \in H_{\Gamma_D}^1(\Omega)$ and $\boldsymbol{\sigma}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$.

Proof. We first observe that if $\tilde{\phi} \in H_{\Gamma_D}^1(\Omega)$, integration by parts gives

$$(\mathfrak{G}(\tilde{\phi}), \boldsymbol{\tau})_{\mathcal{T}_h} + (\tilde{\phi}, \nabla \cdot \boldsymbol{\tau}) - (\tilde{\phi}, \boldsymbol{\tau} \cdot \mathbf{n})_{\Gamma_R} = (\nabla \tilde{\phi}, \boldsymbol{\tau})_{\mathcal{T}_h} + (\tilde{\phi}, \nabla \cdot \boldsymbol{\tau}) - (\tilde{\phi}, \boldsymbol{\tau} \cdot \mathbf{n})_{\Gamma_R} = 0$$

for all $\boldsymbol{\tau} \in \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$ due to the essential boundary conditions on Γ_D and Γ_N . Thus,

$$(\mathfrak{G}(\phi), \boldsymbol{\sigma})_{\mathcal{T}_h} + (\phi, \nabla \cdot \boldsymbol{\sigma})_{\mathcal{T}_h} - (\phi, \boldsymbol{\sigma} \cdot \mathbf{n})_{\mathcal{F}_h^R} = (\mathfrak{G}(\phi - \tilde{\phi}), \boldsymbol{\sigma})_{\mathcal{T}_h} + (\phi - \tilde{\phi}, \nabla \cdot \boldsymbol{\sigma})_{\mathcal{T}_h} - (\phi - \tilde{\phi}, \boldsymbol{\sigma} \cdot \mathbf{n})_{\mathcal{F}_h^R}.$$

If $\boldsymbol{\sigma}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{H}(\text{div}, \Omega)$, we also have

$$0 = (\mathfrak{G}(\phi), \boldsymbol{\sigma}_h)_{\mathcal{T}_h} + (\phi, \nabla \cdot \boldsymbol{\sigma}_h)_{\mathcal{T}_h} - (\phi, \boldsymbol{\sigma}_h \cdot \mathbf{n})_{\mathcal{F}_h^R} \\ = (\mathfrak{G}(\phi) - \nabla \tilde{\phi}, \boldsymbol{\sigma}_h)_{\mathcal{T}_h} + (\nabla \tilde{\phi}, \boldsymbol{\sigma}_h)_{\mathcal{T}_h} + (\phi, \nabla \cdot \boldsymbol{\sigma}_h)_{\mathcal{T}_h} - (\phi, \boldsymbol{\sigma}_h \cdot \mathbf{n})_{\mathcal{F}_h^R} \\ = (\mathfrak{G}(\phi - \tilde{\phi}), \boldsymbol{\sigma}_h)_{\mathcal{T}_h} + (\phi - \tilde{\phi}, \nabla \cdot \boldsymbol{\sigma}_h)_{\mathcal{T}_h} - (\phi - \tilde{\phi}, \boldsymbol{\sigma}_h \cdot \mathbf{n})_{\mathcal{F}_h^R},$$

and (3.1) follows after summation. Then, (3.2) is a direct consequence of (2.5). \square

A simple consequence of Lemma 3.1 is the following result, which is crucial to estimate the non-conformity error in the context of duality arguments.

Corollary 3.2 (Control of the non-conformity). *For all $\phi \in H^1(\mathcal{T}_h)$ and $\xi \in H^1(\Omega)$ with $\mathbf{A}\nabla\xi \in \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$, we have*

$$(3.3) \quad \left| a_h(\phi, \xi) + (\phi, \nabla \cdot (\mathbf{A}\nabla\xi))_{\mathcal{T}_h} - (\phi, \mathbf{A}\nabla\xi \cdot \mathbf{n})_{\mathcal{F}_h^R} \right| \leq \|\phi - \tilde{\phi}\|_{\dagger, 1, \mathcal{T}_h} \|\mathbf{A}\nabla\xi - \boldsymbol{\sigma}_h\|_{\dagger, \text{div}, \mathcal{T}_h}$$

for all $\tilde{\phi} \in H_{\Gamma_D}^1(\Omega)$ and $\boldsymbol{\sigma}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$.

4. A PRIORI ANALYSIS

The purpose of this section is to establish the existence and uniqueness of a discrete solution $u_h \in \mathcal{P}_p(\mathcal{T}_h)$ to (2.11) and to derive a priori error estimates controlling $u - u_h$ in suitable norms. The proof follows the line of the Schatz argument [34], with suitable modifications to take into account the non-conforming nature of the discrete scheme.

In this section, we require that there exists $\rho > 0$ such that

$$(4.1) \quad \text{Re } s_h(v_h, v_h) \geq \rho^2 \|v_h - \pi^g v_h\|_{\dagger, 1, \mathcal{T}_h}^2 \quad \forall v_h \in \mathcal{P}_p(\mathcal{T}_h).$$

This assumption always holds true when $s_h(\cdot, \cdot) = \beta/h(\llbracket \cdot \rrbracket, \llbracket \cdot \rrbracket)_{\mathcal{F}_h}$ with $\beta > 0$. It is also the case when s_h takes the form (2.10) if the penalization parameter is sufficiently large [2]. We note that this assumption rules out some choices of stabilization including complex-value stabilization parameters, as proposed e.g. in [20, 33]. For the sake of simplicity, we only consider stabilizations satisfying (4.1), but slight modifications of our proofs could handle more general situations.

We start with the main duality argument, which is a generalization of the Aubin-Nitsche trick.

Lemma 4.1 (Aubin-Nitsche). *Assume that $u_h \in \mathcal{P}_p(\mathcal{T}_h)$ satisfies (2.11). Then, we have*

$$(4.2) \quad \omega \|u - u_h\|_{\mu, \mathcal{T}_h} \leq \check{\gamma}_{\text{ba}, g} \|u - u_h\|_{\omega, \mathcal{T}_h} + \check{\gamma}_{\text{ba}, d} \|u_h - \pi^g u_h\|_{\dagger, 1, \mathcal{T}_h}$$

and

$$(4.3) \quad \omega^{1/2} \|u - u_h\|_{\gamma, \mathcal{F}_h^R} \leq \tilde{\gamma}_{\text{ba}, g} \|u - u_h\|_{\omega, \mathcal{T}_h} + \tilde{\gamma}_{\text{ba}, d} \|u_h - \pi^g u_h\|_{\dagger, 1, \mathcal{T}_h}.$$

Proof. For the sake of shortness, we set $e_h := u - u_h$. We first establish (4.2). To do so, we define ξ as the unique element of $H_{\Gamma_D}^1(\Omega)$ such that $b(\phi, \xi) = \omega(\mu\phi, e_h)_\Omega$ for all $\phi \in H_{\Gamma_D}^1(\Omega)$. Integrating by parts, we see that the identities

$$-\omega^2 \mu \xi - \nabla \cdot (\mathbf{A}\nabla\xi) = e_h \quad \mathbf{A}\nabla\xi \cdot \mathbf{n} + i\omega\xi = 0$$

respectively hold in $L^2(\Omega)$ and $L^2(\Gamma_R)$. As a result,

$$\omega \|e_h\|_{\mu, \mathcal{T}_h}^2 = -\omega^2 (\mu e_h, \xi) - i\omega (e_h, \phi)_{\mathcal{F}_h^R} - \{(e_h, \nabla \cdot (\mathbf{A}\nabla\xi))_{\mathcal{T}_h} - (e_h, \mathbf{A}\nabla\xi \cdot \mathbf{n})_{\mathcal{F}_h^R}\}$$

and we have from Corollary 3.2 that

$$\omega \|e_h\|_{\mu, \mathcal{T}_h}^2 \leq |b_h(e_h, \xi)| + \|e_h - \tilde{\phi}\|_{\dagger, 1, \mathcal{T}_h} \|\mathbf{A}\nabla\xi - \boldsymbol{\sigma}_h\|_{\dagger, \text{div}, \mathcal{T}_h}$$

for all $\tilde{\phi} \in H_{\Gamma_D}^1(\Omega)$ and $\boldsymbol{\sigma}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{X}_{\Gamma_N}(\text{div}, \Omega)$. On the one hand, we select $\tilde{\phi} = u - \pi^g u_h$ and $\boldsymbol{\sigma}_h = \pi_h^d(\mathbf{A}\nabla\xi)$, so that using (2.14b), we have

$$\omega \|e_h\|_{\mu, \mathcal{T}_h}^2 \leq |b_h(e_h, \xi)| + \check{\gamma}_{\text{ba}, d} \|u_h - \pi^g u_h\|_{\dagger, 1, \mathcal{T}_h} \|e_h\|_{\mu, \mathcal{T}_h}.$$

On the other hand, using (2.12) and (2.13), we have

$$|b_h(e_h, \xi)| = |b_h(e_h, \xi - \pi_h^g \xi)| \leq \|e_h\|_{\omega, \mathcal{T}_h} \|\xi - \pi_h^g \xi\|_{\omega, \Omega} \leq \check{\gamma}_{\text{ba}, g} \|e_h\|_{\omega, \mathcal{T}_h} \|e_h\|_{\mu, \mathcal{T}_h}.$$

We then prove (4.3). Instead of ξ , we define χ as the unique element of $H_{\Gamma_D}^1(\Omega)$ such that $b(\phi, \chi) = \omega^{1/2}(\gamma\chi, e_h)_{\Gamma_R}$ for all $\phi \in H_{\Gamma_D}^1(\Omega)$. We then have

$$-\omega^2\mu\chi - \nabla \cdot (\mathbf{A}\nabla\chi) = 0 \quad \mathbf{A}\nabla\chi \cdot \mathbf{n} + i\omega\chi = e_h$$

in $L^2(\Omega)$ and $L^2(\Gamma_R)$, respectively. We thus have

$$\omega^{1/2}\|e_h\|_{\gamma, \mathcal{F}_h^R}^2 = -\omega^2(\mu e_h, \chi) - i\omega(e_h, \phi)_{\mathcal{F}_h^R} - \{(e_h, \nabla \cdot (\mathbf{A}\nabla\chi))_{\mathcal{T}_h} - (e_h, \mathbf{A}\nabla\chi \cdot \mathbf{n})_{\mathcal{F}_h^R}\},$$

and applying again Corollary 3.2 we arrive at

$$\omega^{1/2}\|e_h\|_{\gamma, \mathcal{F}_h^R}^2 \leq |b_h(e_h, \chi - \pi_h^g\chi)| + \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h} \|\mathbf{A}\nabla\chi - \pi_h^d(\mathbf{A}\nabla\chi)\|_{\dagger, \text{div}, \mathcal{T}_h},$$

and the result follows from the definitions of $\tilde{\gamma}_{\text{ba},g}$ and $\tilde{\gamma}_{\text{ba},d}$ given in (2.14c). \square

We can now complete the Schatz argument, leading to quasi-optimality of the discrete solution for sufficiently refined meshes.

Theorem 4.2 (A priori estimate). *If $\gamma_{\text{ba},g} < 1$ and $\gamma_{\text{ba},d} \leq \rho$, then there exists a unique solution $u_h \in \mathcal{P}_p(\mathcal{T}_h)$ to (2.11) and the estimate*

$$(4.4) \quad \|u - u_h\|_{\omega, \mathcal{T}_h} \leq \frac{1}{1 - \gamma_{\text{ba},g}^2} \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_{\Gamma_D}^1(\Omega)} \|u - v_h\|_{\omega, \Omega}$$

holds true. In addition, if $\gamma_{\text{ba},d} < \rho$, we also have

$$(4.5) \quad \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h} \leq \left(\frac{1}{(1 - \gamma_{\text{ba},g}^2)(\rho^2 - \gamma_{\text{ba},d}^2)} \right)^{1/2} \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_{\Gamma_D}^1(\Omega)} \|u - v_h\|_{\omega, \Omega}.$$

Proof. We first observe that

$$\begin{aligned} \text{Re } b_h(e_h, e_h) + \omega \|e_h\|_{\gamma, \mathcal{F}_h^R}^2 + 2\omega^2 \|e_h\|_{\mu, \mathcal{T}_h}^2 &= \|e_h\|_{\omega, \mathcal{T}_h}^2 + \text{Re } s_h(u_h, u_h) \\ &\geq \|e_h\|_{\omega, \Omega}^2 + \rho^2 \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h}^2. \end{aligned}$$

As a result, we have

$$\begin{aligned} \|e_h\|_{\omega, \mathcal{T}_h}^2 + \rho^2 \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h}^2 &\leq |b_h(e_h, u - v_h)| + 2\omega^2 \|e_h\|_{\mu, \mathcal{T}_h}^2 + \omega \|e_h\|_{\gamma, \mathcal{F}_h^R}^2 \\ &\leq \|e_h\|_{\omega, \mathcal{T}_h} \|u - v_h\|_{\omega, \Omega} \\ &\quad + 2(\tilde{\gamma}_{\text{ba},g} \|e_h\|_{\omega, \mathcal{T}_h} + \tilde{\gamma}_{\text{ba},d} \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h})^2 \\ &\quad + (\tilde{\gamma}_{\text{ba},g} \|e_h\|_{\omega, \mathcal{T}_h} + \tilde{\gamma}_{\text{ba},d} \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h})^2 \\ &\leq \|e_h\|_{\omega, \mathcal{T}_h} \|u - v_h\|_{\omega, \Omega} \\ &\quad + (4\tilde{\gamma}_{\text{ba},g} + 2\tilde{\gamma}_{\text{ba},g}) \|e_h\|_{\omega, \mathcal{T}_h}^2 + (4\tilde{\gamma}_{\text{ba},d} + 2\tilde{\gamma}_{\text{ba},d}) \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h}^2 \end{aligned}$$

so that

$$(1 - 4\tilde{\gamma}_{\text{ba},g}^2 - 2\tilde{\gamma}_{\text{ba},g}^2) \|e_h\|_{\omega, \mathcal{T}_h}^2 + (\rho^2 - 4\tilde{\gamma}_{\text{ba},d}^2 - 2\tilde{\gamma}_{\text{ba},d}^2) \|u_h - \pi_h^g u_h\|_{\dagger, 1, \mathcal{T}_h}^2 \leq \|e_h\|_{\omega, \mathcal{T}_h} \|u - v_h\|_{\omega, \Omega},$$

from which (4.4) and (4.5) follows, recalling the definitions of $\gamma_{\text{ba},g}$ and $\gamma_{\text{ba},d}$ in (2.14). \square

5. A POSTERIORI ANALYSIS

We now provide a posteriori error estimates under minimal regularity assumptions. We first present an abstract framework, and then show how it can be applied to the particular of residual-based estimators.

Throughout this section, we assume that u_h is a fixed function in $\mathcal{P}_p(\mathcal{T}_h)$ satisfying (2.11). Notice that unique solvability of (2.11) is not required. In particular, the proposed analysis applies without any restriction on the mesh size or polynomial degree. Also, in contrast to the a priori analysis, we do not need any specific positivity assumption on the stabilisation form s_h .

5.1. Abstract reliability analysis. Following [19, Theorem 3.3], the discretisation error may be bounded using two different terms that respectively control the equation residual, and the non-conformity of the discrete solution. For the equation residual, we introduce the residual functional and its norm

$$(5.1a) \quad \langle \mathcal{R}_r, v \rangle := b_h(e_h, v) \quad \forall v \in H_{\Gamma_D}^1(\Omega), \quad R_r := \sup_{\substack{v \in H_{\Gamma_D}^1(\Omega) \\ \|\nabla v\|_{\mathbf{A}, \Omega} = 1}} |\langle \mathcal{R}_r, v \rangle|.$$

Notice that this first term only involves conforming test functions. To account for the non-conformity of the discrete solution, we additionally introduce

$$(5.1b) \quad R_c := \|u_h - \pi^g u_h\|_{\dagger, 1, \mathcal{T}_h}.$$

The total abstract estimator is then the Hilbertian sum of the two above components:

$$(5.1c) \quad R^2 := R_r^2 + R_c^2.$$

Following [9, 33], our a posteriori analysis follows the lines of the Schatz argument [34]. Specifically, we start by controlling weak norms of the errors with the estimator and the approximation factor using a duality argument.

Lemma 5.1 (Aubin-Nitsche). *The following estimates hold true:*

$$(5.2) \quad \omega \|u - u_h\|_{\mu, \mathcal{T}_h} \leq \check{\gamma}_{\text{ba}} R, \quad \omega^{1/2} \|u - u_h\|_{\gamma, \mathcal{F}_h^R} \leq \check{\gamma}_{\text{ba}} R.$$

Proof. Let us set $e_h := u - u_h$. We start with the first estimate in (5.2). We define ξ as the unique element of $H_{\Gamma_D}^1(\Omega)$ such that $b(w, \xi) = \omega(\mu w, e_h)_{\mathcal{T}_h}$ for all $w \in H_{\Gamma_D}^1(\Omega)$. Following the lines the proof of Lemma 4.1, we write

$$\begin{aligned} b_h(e_h, \xi) &= -\omega^2(\mu e_h, \xi)_{\mathcal{T}_h} - i\omega(\gamma e_h, \xi)_{\mathcal{F}_h^R} + (\mathfrak{G}(e_h), \mathbf{A}\nabla\xi)_{\mathcal{T}_h} \\ &= (e_h, -\omega^2\mu\xi - \nabla \cdot (\mathbf{A}\nabla\xi))_{\mathcal{T}_h} - (\gamma e_h, \mathbf{A}\nabla\xi \cdot \mathbf{n})_{\mathcal{F}_h^R} + (e_h, \nabla \cdot (\mathbf{A}\nabla\xi)) + (\mathfrak{G}(e_h), \mathbf{A}\nabla\xi)_{\mathcal{T}_h} \\ &= \omega \|e_h\|_{\mu, \mathcal{T}_h}^2 + (\mathfrak{G}(\pi^g u_h - u_h), \boldsymbol{\sigma} - \boldsymbol{\sigma}_h)_{\mathcal{T}_h} + (\pi^g u_h - u_h, \nabla \cdot (\boldsymbol{\sigma} - \boldsymbol{\sigma}_h))_{\mathcal{T}_h}, \end{aligned}$$

for all $\boldsymbol{\sigma}_h \in \mathcal{P}_p(\mathcal{T}_h) \cap \mathbf{H}(\text{div}, \Omega)$, where we employed (3.1) in the last identity. It follows that

$$\omega \|e_h\|_{\mu, \mathcal{T}_h}^2 \leq |b_h(e_h, \xi)| + \|u_h - \pi^g u_h\|_{\dagger, 1, \mathcal{T}_h} \|\mathbf{A}\nabla\xi - \boldsymbol{\sigma}_h\|_{\dagger, \text{div}, \mathcal{T}_h}.$$

On the one hand, recalling the Galerkin orthogonality property stated in (2.12) and the definition of R_r at (5.1a), we have

$$|b_h(e_h, \xi)| = |b_h(e_h, \xi - \pi_h^g \xi)| \leq R_r \|\nabla(\xi - \pi_h^g \xi)\|_{\mathbf{A}, \Omega} \leq \check{\gamma}_{\text{ba}}^g R_r \|e_h\|_{\mu, \mathcal{T}_h}.$$

On the other hand, recalling (2.14b) and (5.1b), we have

$$\|u_h - \pi^g u_h\|_{\dagger, 1, \mathcal{T}_h} \|\mathbf{A}\nabla\xi - \pi_h^d(\mathbf{A}\nabla\xi)\|_{\dagger, \text{div}, \mathcal{T}_h} \leq \check{\gamma}_{\text{ba}}^d R_c \|e_h\|_{\mu, \mathcal{T}_h}.$$

Then, the first estimate of (5.2) follows from (5.1c) since

$$\check{\gamma}_{\text{ba}}^{\text{g}} \mathbf{R}_{\text{r}} + \check{\gamma}_{\text{ba}}^{\text{d}} \mathbf{R}_{\text{c}} \leq \left((\check{\gamma}_{\text{ba}}^{\text{g}})^2 + (\check{\gamma}_{\text{ba}}^{\text{d}})^2 \right)^{1/2} (\mathbf{R}_{\text{r}}^2 + \mathbf{R}_{\text{c}}^2)^{1/2} = \check{\gamma}_{\text{ba}} \mathbf{R}.$$

For the sake of shortness, we do not detail the proof of the second estimate in (5.2) as it follows the lines of the first estimate but using the function χ from the proof of Lemma 4.1 instead of ξ . \square

We can now conclude the abstract reliability analysis. Following [19], our proof uses a Pythagorean identity to separate the conforming and non-conforming parts of the error.

Theorem 5.2 (Abstract reliability). *We have*

$$(5.3) \quad \|u - u_h\|_{\omega, \mathcal{T}_h} \leq \sqrt{1 + \gamma_{\text{ba}}^2} \mathbf{R}.$$

Proof. For the sake of simplicity, we introduce

$$b_h^+(\phi, v) := \omega^2(\mu\phi, v)_{\mathcal{T}_h} + \omega(\gamma\phi, v)_{\mathcal{F}_h^{\text{R}}} + (\mathbf{A}\mathfrak{G}(\phi), \mathfrak{G}(v))_{\mathcal{T}_h} \quad \forall \phi, v \in H^1(\mathcal{T}_h),$$

and we define \tilde{u} as the only element of $H_{\Gamma_{\text{D}}}^1(\Omega)$ such that

$$b_h^+(\tilde{u} - u_h, v) = 0 \quad \forall v \in H_{\Gamma_{\text{D}}}^1(\Omega).$$

Notice that this indeed a well-formed definition, since b_h^+ is equivalent to the usual inner-product of $H_{\Gamma_{\text{D}}}^1(\Omega)$. Classically, we have the Pythagorean identity

$$(5.4) \quad \begin{aligned} \|u - u_h\|_{\omega, \mathcal{T}_h}^2 &= \|\tilde{u} - u_h\|_{\omega, \mathcal{T}_h}^2 + 2 \operatorname{Re} b^+(\tilde{u} - u_h, u - \tilde{u}) + \|u - \tilde{u}\|_{\omega, \mathcal{T}_h}^2 \\ &= \|u_h - \tilde{u}\|_{\omega, \mathcal{T}_h}^2 + \|u - \tilde{u}\|_{\omega, \mathcal{T}_h}^2. \end{aligned}$$

Then, on the one hand, it is clear using (2.6) that

$$(5.5) \quad \|u_h - \tilde{u}\|_{\omega, \mathcal{T}_h} \leq \|u_h - \pi^{\text{g}} u_h\|_{\omega, \mathcal{T}_h} \leq \|u_h - \pi^{\text{g}} u_h\|_{\dagger, 1, \mathcal{T}_h} = \mathbf{R}_{\text{c}},$$

and on the other hand, we have

$$\begin{aligned} \|u - \tilde{u}\|_{\omega, \Omega}^2 &= b_h^+(u - \tilde{u}, u - \tilde{u}) = b_h^+(e_h, u - \tilde{u}) \\ &= b_h(e_h, u - \tilde{u}) + 2\omega^2(\mu e_h, u - \tilde{u})_{\mathcal{T}_h} + (1 - i)\omega(\gamma e_h, u - \tilde{u})_{\mathcal{F}_h^{\text{R}}} \\ &\leq \mathbf{R}_{\text{r}} \|\nabla(u - \tilde{u})\|_{\mathbf{A}, \Omega} + 2\check{\gamma}_{\text{ba}} \mathbf{R} \omega \|u - \tilde{u}\|_{\mu, \mathcal{T}_h} + \sqrt{2}\check{\gamma}_{\text{ba}} \mathbf{R} \omega \|u - \tilde{u}\|_{\gamma, \mathcal{F}_h^{\text{R}}} \\ &\leq (\mathbf{R}_{\text{r}}^2 + (4\check{\gamma}_{\text{ba}}^2 + 2\check{\gamma}_{\text{ba}}^2) \mathbf{R}^2)^{1/2} \|u - \tilde{u}\|_{\omega, \Omega}, \end{aligned}$$

and recalling (2.14c), we arrive at

$$(5.6) \quad \|u - \tilde{u}\|_{\omega, \Omega}^2 \leq \mathbf{R}_{\text{r}}^2 + (4\check{\gamma}_{\text{ba}}^2 + 2\check{\gamma}_{\text{ba}}^2) \mathbf{R}^2 = \mathbf{R}_{\text{r}}^2 + \gamma_{\text{ba}}^2 \mathbf{R}^2.$$

The estimate in (5.3) then follows from (5.4), (5.5) and (5.6). \square

5.2. Residual-based a posteriori estimator. The residuals \mathbf{R}_{r} and \mathbf{R}_{c} can be straightforwardly controlled using flux and potential reconstructions. We refer the reader to [14, 19] for details about these constructions. Here, we focus instead on residual-based estimators, for which the link may be less clear. In this section, the letter C refers to a generic constant that may change from one occurrence to the other, and that only depends on the contrasts of the

coefficients μ , \mathbf{A} and γ as well as the polynomial degree p and the shape-regularity parameter κ . The stability result for the lifting operator

$$(5.7) \quad \|\ell_h(\llbracket v_h \rrbracket)\|_{\mathbf{A}, \mathcal{T}_h}^2 \leq C \sum_{K \in \mathcal{T}_h} \frac{\alpha_K}{h_K} \|\llbracket v_h \rrbracket\|_{\partial K}^2 \quad \forall v_h \in \mathcal{P}_p(\mathcal{T}_h)$$

can be found, e.g., in [32, Section 4.3] or [26, Lemma 4.1].

We first establish that R_c can be controlled by the jumps of u_h . The notations

$$\frac{\omega h_F^*}{\vartheta_F^*} := \max_{F \in \mathcal{F}_h^R} \frac{\omega h_F}{\vartheta_F}, \quad \frac{\omega h_K^*}{\vartheta_K^*} := \max_{K \in \mathcal{T}_h} \frac{\omega h_K}{\vartheta_K},$$

will be useful.

Lemma 5.3 (Control of the non-conformity). *We have*

$$(5.8) \quad R_c^2 \leq C \max \left(1, \frac{\omega h_F^*}{\vartheta_F^*}, \left(\frac{\omega h_K^*}{\vartheta_K^*} \right)^2 \right) \sum_{K \in \mathcal{T}_h} \frac{\alpha_K}{h_K} \|\llbracket u_h \rrbracket\|_{\partial K \setminus (\Gamma_N \cup \Gamma_R)}^2.$$

Proof. We start with two estimates that are easily inferred from [17, Lemma 4.3]. For all $v_h \in \mathcal{P}_p(\mathcal{T}_h)$, there exists a conforming approximation $\mathcal{J}v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_{\Gamma_D}^1(\Omega)$ such that and all $K \in \mathcal{T}_h$, we have

$$\|v_h - \mathcal{J}v_h\|_K^2 \leq Ch_K \sum_{K' \in \tilde{\mathcal{T}}_K} \|\llbracket v_h \rrbracket\|_{\partial K'}^2, \quad \|\nabla(v_h - \mathcal{J}v_h)\|_K^2 \leq C \frac{1}{h_K} \sum_{K' \in \tilde{\mathcal{T}}_K} \|\llbracket v_h \rrbracket\|_{\partial K'}^2,$$

where $\tilde{\mathcal{T}}_K$ collects those elements $K' \in \mathcal{T}_h$ sharing at least one vertex with K . Employing the multiplicative trace inequality

$$\|\theta\|_{\partial K}^2 \leq C \left\{ \frac{1}{h_K} \|\theta\|_K^2 + \|\theta\|_K \|\nabla \theta\|_K \right\} \leq C \left\{ \frac{1}{h_K} \|\theta\|_K^2 + h_K \|\nabla \theta\|_K^2 \right\} \quad \forall \theta \in H^1(K),$$

we also have

$$\|v_h - \mathcal{J}v_h\|_{\partial K}^2 \leq C \sum_{K' \in \tilde{\mathcal{T}}_K} \|\llbracket v_h \rrbracket\|_{\partial K'}^2.$$

It is therefore clear that

$$\|u_h - \mathcal{J}u_h\|_{\dagger, 1, \mathcal{T}_h}^2 \leq C \max \left(1, \frac{\omega h_F^*}{\vartheta_F^*}, \left(\frac{\omega h_K^*}{\vartheta_K^*} \right)^2 \right) \sum_{K \in \mathcal{T}_h} \frac{\alpha_K}{h_K} \|\llbracket u_h \rrbracket\|_{\partial K}^2,$$

and (5.8) follows from the definition of R_c in (5.1b). \square

We then control the conforming residual R_r . This is classically done by combining element-wise integration by parts and a quasi-interpolation operator.

Lemma 5.4 (Control of the residual). *We have*

$$(5.9) \quad R_r^2 \leq C \left\{ \sum_{K \in \mathcal{T}_h} \left(\frac{h_K^2}{\alpha_K} \|f + \omega^2 \mu u_h + \nabla \cdot (\mathbf{A} \nabla u_h)\|_{\mathcal{T}_h}^2 + \frac{h_K}{\alpha_K} \|\llbracket \mathbf{A} \nabla u_h \rrbracket \cdot \mathbf{n}_K\|_{\partial K \setminus \partial \Omega}^2 + \frac{\alpha_K}{h_K} \|\llbracket u_h \rrbracket\|_{\partial K \setminus \partial \Omega}^2 \right) \right. \\ \left. + \sum_{F \in \mathcal{F}_h^D} \frac{\alpha_F}{h_F} \|u_h\|_F^2 + \sum_{F \in \mathcal{F}_h^N} \frac{h_F}{\alpha_F} \|\mathbf{A} \nabla u_h \cdot \mathbf{n}\|_F^2 + \sum_{F \in \mathcal{F}_h^R} \frac{h_F}{\alpha_F} \|\mathbf{A} \nabla u_h \cdot \mathbf{n} - i\omega \gamma u_h\|_F^2 \right\}.$$

Proof. We consider an arbitrary $v \in H_{\Gamma_D}^1(\Omega)$, and let $\tilde{v} := v - \mathcal{Q}_h v \in H_{\Gamma_D}^1(\Omega)$, where \mathcal{Q}_h is a quasi-interpolation operator satisfying

$$\frac{1}{h_K^2} \|v - \mathcal{Q}_h v\|_K^2 + \frac{1}{h_K} \|v - \mathcal{Q}_h v\|_{\partial K}^2 + \|\nabla(v - \mathcal{Q}_h v)\|_K^2 \leq C \frac{1}{\alpha_K} \|\nabla v\|_{\mathbf{A}, \tilde{K}},$$

with \tilde{K} the domain corresponding to all the elements $K' \in \mathcal{T}_h$ that share at least one vertex with K . We refer the reader to, e.g., [17, Theorem 5.2], for the construction of \mathcal{Q}_h . Thanks to Galerkin orthogonality (2.12), we can write that

$$\begin{aligned} (5.10) \quad & \langle \mathcal{R}_r, v \rangle \\ &= b_h(e_h, v) = b_h(e_h, \tilde{v}) \\ &= (f, \tilde{v})_\Omega + \omega^2 (\mu u_h, \tilde{v})_{\mathcal{T}_h} + i\omega (\gamma u_h, \tilde{v})_{\mathcal{F}_h^R} + (\mathbf{A}\mathcal{G}(u_h), \nabla \tilde{v})_{\mathcal{T}_h} \\ &= \underbrace{(f, \tilde{v})_\Omega + \omega^2 (\mu u_h, \tilde{v})_{\mathcal{T}_h} + i\omega (\gamma u_h, \tilde{v})_{\mathcal{F}_h^R} - (\mathbf{A}\nabla u_h, \nabla \tilde{v})_{\mathcal{T}_h}}_{r_1} - \underbrace{(\mathbf{A}\ell_h(\llbracket u_h \rrbracket), \nabla \tilde{v})_{\mathcal{T}_h}}_{r_2}. \end{aligned}$$

Then, on the one hand, we see that

$$\begin{aligned} r_1 &= (f + \omega^2 \mu u_h + \nabla \cdot (\mathbf{A}\nabla u_h), \tilde{v})_{\mathcal{T}_h} + i\omega (\gamma u_h, \tilde{v})_{\mathcal{F}_h^R} - \sum_{K \in \mathcal{T}_h} (\mathbf{A}\nabla u_h \cdot \mathbf{n}_K, \tilde{v})_{\partial K} \\ &= (f + \omega^2 \mu u_h - \nabla \cdot (\mathbf{A}\nabla u_h), \tilde{v})_{\mathcal{T}_h} + \sum_{F \in \mathcal{F}_h^i} (\llbracket \mathbf{A}\nabla u_h \rrbracket \cdot \mathbf{n}_F, \tilde{v})_F \\ &\quad - (\mathbf{A}\nabla u_h \cdot \mathbf{n}, \tilde{v})_{\mathcal{F}_h^N} - (\mathbf{A}\nabla u_h \cdot \mathbf{n} - i\omega \gamma u_h, \tilde{v})_{\mathcal{F}_h^R}, \end{aligned}$$

and therefore,

$$\begin{aligned} (5.11) \quad |r_1| &\leq \sum_{K \in \mathcal{T}_h} (\|f + \omega^2 \mu u_h + \nabla \cdot (\mathbf{A}\nabla u_h)\|_K \|\tilde{v}\|_K + \|\llbracket \nabla u_h \rrbracket \cdot \mathbf{n}_K\|_{\partial K} \|\tilde{v}\|_{\partial K}) \\ &\quad + \sum_{F \in \mathcal{F}_h^N} \|\mathbf{A}\nabla u_h \cdot \mathbf{n}\|_F \|\tilde{v}\|_F + \sum_{F \in \mathcal{F}_h^R} \|\mathbf{A}\nabla u_h \cdot \mathbf{n} - i\omega \gamma u_h\|_F \|\tilde{v}\|_F \\ &\leq C \left\{ \sum_{K \in \mathcal{T}_h} \left(\frac{h_K^2}{\alpha_K} \|f + \omega^2 \mu u_h + \nabla \cdot (\mathbf{A}\nabla u_h)\|_K + \frac{h_K}{\alpha_K} \|\llbracket \nabla u_h \rrbracket \cdot \mathbf{n}_K\|_{\partial K} \right) \right. \\ &\quad \left. + \sum_{F \in \mathcal{F}_h^N} \frac{h_F}{\alpha_F} \|\mathbf{A}\nabla u_h \cdot \mathbf{n}\|_F + \sum_{F \in \mathcal{F}_h^R} \frac{h_F}{\alpha_F} \|\mathbf{A}\nabla u_h \cdot \mathbf{n} - i\omega \gamma u_h\|_F \right\} \|\nabla v\|_{\mathbf{A}, \Omega}. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} (5.12) \quad |r_2| &\leq \|\ell_h(\llbracket u_h \rrbracket)\|_{\mathbf{A}, \Omega} \|\nabla \tilde{v}\|_{\mathbf{A}, \Omega} \\ &\leq C \left\{ \sum_{K \in \mathcal{T}_h} \frac{\alpha_K}{h_K} \|\llbracket u_h \rrbracket\|_{\partial K \setminus \partial \Omega}^2 + \sum_{F \in \mathcal{F}_h^D} \frac{\alpha_F}{h_F} \|\llbracket u_h \rrbracket\|_F^2 \right\} \|\nabla v\|_{\mathbf{A}, \Omega}, \end{aligned}$$

and (5.9) follows from (5.10), (5.11) and (5.12) recalling the definition of R_r in (5.1a). \square

We therefore define a residual-based estimator by

$$\begin{aligned} \eta^2 := & \sum_{K \in \mathcal{T}_h} \left(\frac{h_K^2}{\alpha_K} \|f + \omega^2 \mu u_h - \nabla \cdot (\mathbf{A} \nabla u_h)\|_{\mathcal{T}_h}^2 + \frac{h_K}{\alpha_K} \|[[\mathbf{A} \nabla u_h]] \cdot \mathbf{n}_K\|_{\partial K \setminus \partial \Omega}^2 + \frac{\alpha_K}{h_K} \|[[u_h]]\|_{\partial K \setminus \partial \Omega}^2 \right) \\ & + \sum_{F \in \mathcal{F}_h^D} \frac{\alpha_F}{h_F} \|u_h\|_F^2 + \sum_{F \in \mathcal{F}_h^N} \frac{h_F}{\alpha_F} \|\mathbf{A} \nabla u_h \cdot \mathbf{n}\|_F^2 + \sum_{F \in \mathcal{F}_h^R} \frac{h_F}{\alpha_F} \|\mathbf{A} \nabla u_h \cdot \mathbf{n}\|_F^2. \end{aligned}$$

As a direct consequence of (5.3), (5.8) and (5.9), we obtain a reliability estimate stated in Theorem 5.5. It is to be compared with [9, Theorem 2.3] and [33, Theorem 3.6].

Theorem 5.5 (Reliability of the residual estimator). *We have*

$$\|e_h\|_{\omega, \mathcal{T}_h} \leq C \max \left(1, \frac{\omega h_F^*}{\vartheta_F^*}, \left(\frac{\omega h_K^*}{\vartheta_K^*} \right)^2 \right) (1 + \gamma_{\text{ba}}) \eta.$$

5.3. Efficiency. For the sake of shortness, we do not give any efficiency proofs. Actually, such proofs can be found in [14, Section 4] for equilibrated estimators and in [33, Section 3.3] for the residual-based estimators considered here.

REFERENCES

1. R. Adams and J. Fournier, *Sobolev spaces*, Academic Press, 2003.
2. C. Agut and J. Diaz, *Stability analysis of the interior penalty discontinuous Galerkin method for the wave equation*, ESAIM Math. Model. Numer. Anal. **47** (2013), 903–932.
3. D.N. Arnold, F. Brezzi, B. Cockburn, and L.D. Marini, *Unified analysis of discontinuous Galerkin, methods for elliptic problems*, SIAM J. Numer. Anal. **39** (2002), no. 5, 1749–1779.
4. H. Barucq, T. Chaumont-Frelet, and C. Gout, *Stability analysis of heterogeneous Helmholtz problems and finite element solution based on propagation media approximation*, Math. Comp. **86** (2017), no. 307, 2129–2157.
5. M. Bernkopf, S. Sauter, C. Torres, and A. Veit, *Solvability of discrete Helmholtz equations*, arXiv:2105.02273v2, 2022.
6. J. Chand, Z. Wang, A. Modave, J.F. Remacle, and T. Warburton, *GPU-accelerated discontinuous Galerkin methods on hybrid meshes*, J. Comput. Phys. **318** (2016), 142–168.
7. S.N. Chandler-Wilde, E.A. Spence, A. Gibbs, and V.P. Smyshlyaev, *High-frequency bounds for the Helmholtz equation under parabolic trapping and applications in numerical analysis*, SIAM J. Math. Anal. **52** (2020), no. 1, 845–893.
8. T. Chaumont-Frelet, *Mixed finite element discretizations of acoustic Helmholtz problems with high wavenumbers*, Calcolo **56** (2019).
9. T. Chaumont-Frelet, A. Ern, and M. Vohralík, *On the derivation of guaranteed and p -robust a posteriori error estimates for the Helmholtz equation*, Numer. Math. **148** (2021), 525–573.
10. T. Chaumont-Frelet and S. Nicaise, *Wavenumber explicit convergence analysis for finite element discretizations of general wave propagation problems*, IMA J. Numer. Anal. **40** (2020), no. 2, 1503–1543.
11. T. Chaumont-Frelet and E.A. Spence, *Scattering by finely-layered obstacles: frequency-explicit bounds and homogenization*, arXiv preprint arXiv:2109.11267, 2021.
12. P.G. Ciarlet, *The finite element method for elliptic problems*, SIAM, 2002.
13. G. Cohen, P. Joly, J.E. Roberts, and N. Tordjman, *High order triangular finite element with mass lumping for the wave equation*, SIAM J. Numer. Anal. **38** (2001), no. 6, 2047–2078.
14. S. Congreve, J. Gedicke, and I. Perugia, *Robust adaptive hp discontinuous Galerkin finite element methods for the Helmholtz equation*, SIAM J. Sci. Comput. **41** (2019), no. 2, A1121–A1147.
15. M. Costabel, M. Dauge, and S. Nicaise, *Singularities of Maxwell interface problems*, ESAIM Math. Model. Numer. Anal. **33** (1999), no. 3, 627–649.
16. Y. Du and L. Zhu, *Preasymptotic error analysis of high order interior penalty discontinuous Galerkin methods for the Helmholtz equation with high wave number*, J. Sci. Comput. **67** (2016), 130–152.

17. A. Ern and J.L. Guermond, *Finite element quasi-interpolation and best approximation*, ESAIM Math. Model. Numer. Anal. **51** (2017), 1367–1385.
18. ———, *Finite elements I: basic theory and practice*, in preparation, 2019.
19. A. Ern and M. Vohralík, *Polynomial-degree-robust a posteriori estimates in a unified setting for conforming, nonconforming, discontinuous Galerkin, and mixed discretizations*, SIAM J. Numer. Anal. **53** (2015), no. 2, 1058–1081.
20. X. Feng and H. Wu, *Discontinuous Galerkin methods for the Helmholtz equation with large wave number*, SIAM J. Numer. Anal. **47** (2009), no. 4, 2872–2896.
21. ———, *hp-discontinuous Galerkin methods for the Helmholtz equation with large wave number*, Math. Comp. **80** (2011), no. 276, 1997–2024.
22. P. Fernandes and G. Gilardi, *Magnetostatic and electrostatic problems in inhomogeneous anisotropic media with irregular boundary and mixed boundary conditions*, Math. Meth. Appl. Sci. **47** (1997), no. 4, 2872–2896.
23. V. Girault and P.A. Raviart, *Finite element methods for Navier-Stokes equations: theory and algorithms*, Springer-Verlag, 1986.
24. M.J. Grote, A. Schneebeli, and D. Schötzau, *Discontinuous Galerkin finite element method for the wave equation*, SIAM J. Numer. Anal. **44** (2006), no. 6, 2408–2431.
25. R.H.W. Hope and N. Sharma, *Convergence analysis of an adaptive interior penalty discontinuous Galerkin method for the Helmholtz equation*, IMA J. Numer. Anal. **33** (2013), 898–921.
26. P. Houston, D. Schötzau, and T.P. Wihler, *Energy norm a posteriori error estimation of hp-adaptive discontinuous Galerkin methods for elliptic problems*, Math. Models Meth. Appl. Sci. **17** (2006), no. 1, 33–62.
27. D. Lafontaine, E.A. Spence, and J. Wunsch, *For most frequencies, strong tapping has a weak effect in frequency-domain scattering*, Comm. Pure Appl. Math. **LXXIV** (2022), 2025–2063.
28. J.M. Melenk, A. Parsania, and S. Sauter, *General DG-methods for highly indefinite Helmholtz problems*, J. Sci. Comput. **57** (2013), 536–581.
29. J.M. Melenk and S. Sauter, *Convergence analysis for finite element discretizations of the Helmholtz equation with Dirichlet-to-Neumann boundary conditions*, Math. Comp. **79** (2010), no. 272, 1871–1914.
30. A. Moiola and E.A. Spence, *Acoustic transmission problems: wavenumber-explicit bounds and resonance-free regions*, Math. Meth. Appl. Sci. **29** (2019), no. 2, 317–354.
31. P. Monk, *Finite element methods for Maxwell’s equations*, Oxford science publications, 2003.
32. D.A. Di Pietro and A. Ern, *Mathematical aspects of discontinuous Galerkin methods*, Springer, 2012.
33. S. Sauter and J. Zech, *A posteriori error estimation of hp – dg finite element methods for highly indefinite Helmholtz problems*, SIAM J. Numer. Anal. **53** (2015), no. 5, 2414–2440.
34. A.H. Schatz, *An observation concerning Ritz-Galerkin methods with indefinite bilinear forms*, Math. Comp. **28** (1974), no. 128, 959–962.