



**HAL**  
open science

# Stability and accuracy of Runge-Kutta based split-explicit time-stepping algorithms for free-surface ocean models

Florian Lemarié, Nicolas Ducouso, Laurent Debreu, Gervan Madec

► **To cite this version:**

Florian Lemarié, Nicolas Ducouso, Laurent Debreu, Gervan Madec. Stability and accuracy of Runge-Kutta based split-explicit time-stepping algorithms for free-surface ocean models. EGU 2022 - General Assembly on European Geosciences Union, May 2022, Vienna, Austria. pp.1-6. hal-03769267

**HAL Id: hal-03769267**

**<https://inria.hal.science/hal-03769267v1>**

Submitted on 5 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Stability and accuracy of Runge-Kutta based split-explicit time-stepping algorithms for free-surface ocean models

Nicolas Ducousso<sup>1,3</sup>, Florian Lemarié<sup>1</sup>, Laurent Debreu<sup>1</sup> and Gurvan Madec<sup>2,1</sup>

<sup>1</sup> Inria, Grenoble, France

<sup>2</sup> LOCEAN, Paris, France

<sup>3</sup> SHOM, Brest, France

# Runge-Kutta methods

---

## Advantages of RK methods:

- ▶ 2-time-level schemes (no computational mode)
- ▶ Good stability properties for advection and diffusion
- ▶ Abundant literature (e.g. RK-IMEX, RK-SSP, DIRK, etc)
- ▶ Easy to implement (convex combination of Euler steps)

## Main questions:

- ▶ Many variants of RK schemes
- ▶ **RK with mode-splitting ?**

## Existing work:

- ▶ Many RK-based atmospheric models (e.g. WRF, Cosmo, Homme-NH, etc)
- ▶ « *Pseudo-second order Runge-Kutta time stepping scheme* » in HIM/MOM6 [Hallberg, 1997]

# Runge-Kutta methods in the oceanic context

## ► Many variants of RK schemes

- Low-storage and « order=stage » method for advection [Wicker & Skamarock, 2002]

$$\begin{aligned} \partial_t \phi = \mathcal{F}(\phi, t) \quad \longrightarrow \quad & \phi^{n+1/3} = \phi^n + \frac{\Delta t}{3} \mathcal{F}(\phi^n) \\ & \phi^{n+1/2} = \phi^n + \frac{\Delta t}{2} \mathcal{F}(\phi^{n+1/3}) \\ & \phi^{n+1} = \phi^n + \Delta t \mathcal{F}(\phi^{n+1/2}) \end{aligned}$$

- Introduce Shuman averaging for internal gravity waves (« order=stage-1 »)

## ► Additional difficulty inherent to oceanic models


- 2D barotropic / 3D baroclinic mode splitting to improve computational efficiency (fast and slow dynamics are split into separate subproblems)
- Separation of slow and fast modes is non-orthogonal (depth-independent barotropic mode assumption)

➡ Some form of filtering is required to stabilize the splitting


$$\begin{aligned} \partial_t \bar{u} &= -g \partial_x \left( \eta + \frac{1}{\rho_0 g H} \int_{-H}^0 p_h dz \right) \\ \partial_t \eta &= -H \partial_x \bar{u} \end{aligned}$$

# Robust mode-splitting method based on RK3

**Peculiarity of oceanic models:** 2D barotropic / 3D baroclinic mode splitting to improve computational efficiency



Gravity waves integration (without dissipation in 2D mode)



## **Single-first approach:**

- ▶ Only one integration of the 2D within the RK3 time-step
- ▶ Split-explicit approach with a dissipative time-stepping for the barotropic mode
- ▶ Linear stability analysis can be conducted following [Demange et al. \(2019\)](#)


# Robust mode-splitting method based on RK3

**Peculiarity of oceanic models:** 2D barotropic / 3D baroclinic mode splitting to improve computational efficiency


$$\bar{u}^{n+1} = \bar{u}^n - g\Delta t_0 \partial_x \eta^n$$

$$\eta^{n+1} = \eta^n - H\Delta t_0 \partial_x \left( (1+\theta)\bar{u}^{n+1} - \theta\bar{u}^n \right)$$

(a) Barotropic mode amplification



(b) Baroclinic mode amplification



- ▶ Linear stability analysis provides the minimal dissipation to stabilize the splitting
- ▶ Stability constrained by barotropic mode and not by baroclinic ones

# Conclusion and future work

---

## Summary:

- ➔ Design of an RK3 based time-stepping algorithm for split-explicit free-surface ocean models
- ▶ The single-first strategy appears to be more robust than other alternatives
- ▶ Thought for a quasi-Eulerian vertical coordinate
- ▶ A non-linear  $z^*$  2D  $x$ - $z$  model has been developed to check results from stability analysis

## Ongoing work:

- ▶ Implementation in NEMO (see next talk)
- ▶ Accuracy analysis and consequences of mode-splitting

◆ Ducouso, N., Lemarié, F., Debreu, L., Madec, G.: *Stability and accuracy of Runge-Kutta-based time-stepping algorithms for split-explicit free-surface ocean models*. In preparation for J. Adv. Model. Earth Syst.