



HAL
open science

M/EEG networks integration to elicit patterns of motor imagery-based Brain-Computer Interface (BCI) training

Marie-Constance Corsi, Mario Chavez, Denis P Schwartz, Nathalie George, Laurent Hugueville, Ari Kahn, Sophie Dupont, Danielle Bassett, Fabrizio de Vico Fallani

► **To cite this version:**

Marie-Constance Corsi, Mario Chavez, Denis P Schwartz, Nathalie George, Laurent Hugueville, et al.. M/EEG networks integration to elicit patterns of motor imagery-based Brain-Computer Interface (BCI) training. BIOMAG 2022 - 22nd International Conference on Biomagnetism, Aug 2022, Birmingham, United Kingdom. hal-03773296

HAL Id: hal-03773296

<https://inria.hal.science/hal-03773296>

Submitted on 9 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

M/EEG networks integration to elicit patterns of motor imagery-based Brain-Computer Interface (BCI) training

Marie-Constance Corsi¹

Mario Chavez²

Denis Schwartz¹

Nathalie George¹

Laurent Hugueville¹

Ari E. Kahn²

Sophie Dupont¹

Danielle S. Bassett²

Fabrizio De Vico Fallani¹

¹ Sorbonne Université, Institut du Cerveau – Paris Brain Institute, ICM, CNRS, Inria, Inserm, AP-HP, Hôpital de la Pitié-Salpêtrière, F-75013, Paris, France

² University of Pennsylvania, Philadelphia, USA



marie.constance.corsi@gmail.com



@MConstanceCorsi

Non-invasive Brain-Computer Interfaces (BCIs) can exploit the ability of subjects to voluntarily modulate their brain activity through mental imagery. Despite its clinical applications, controlling a BCI appears to be a learned skill that requires several weeks to reach relatively high-performance in control, without being sufficient for 15 to 30 % of the users [1]. This gap has motivated a deeper understanding of mechanisms associated with motor imagery (MI) tasks. Here, we investigated dynamical changes in multimodal network recruitment. We hypothesized that integrating information from EEG and MEG data, show a better description of the core-periphery changes occurring during a motor imagery-based BCI training.


Longitudinal study

The EEG-based BCI task consisted of a standard 1D, two-target box task in which the subjects modulated their α and/or β activity. To hit the target-up, the subjects performed a sustained motor imagery of their right-hand grasping and to hit the target-down, they remained at rest. 20 healthy subjects (aged 27.45 ± 4.01 years, 12 men), all right-handed, participated in the study.

Magnetoencephalography (MEG) and electroencephalography (EEG) signals were simultaneously recorded. M/EEG data were preprocessed using the Independent Component Analysis method, followed by the source reconstruction on the epoched data [2].



Multiplex core-periphery computation



To study the core-periphery properties of the brain networks, we used the probability for a given node to belong to the core, defined as the coreness [3].

Multiplex coreness of node i – C_i

$$C_i = \frac{1}{N-1} \sum_{k=1}^{N-1} \delta_i^k; \quad \delta_i^k = 1, \text{ if nodes } i \text{ in the core, } 0 \text{ otherwise}$$

Optimization of the contribution c of each layer/modality:


$$F(c) = \frac{((c^{MI}(c)) - (c^{Rest}(c)))^2}{(s^{MI})^2 + (s^{Rest})^2}$$

Where:

$$(s^{cond})^2 = \sum_{i \in \{1 \dots N\}} ((c_i^{cond}(c)) - (c^{cond}(c)))^2$$

$(c^{cond}(c))$, averaged coreness over the nodes i

c_i^{cond} , coreness computed in node i , condition $cond$



Studying the network integration changes at the single and multilayer levels provides additional information to characterize dynamic brain reorganization during BCI training. A progressive increase of the integration of somatosensory areas in the α band was paralleled by a decrease of the integration of visual processing and working memory areas in the β band. Such changes were more visible in multiplex in which brain network properties correlated with future BCI scores in the $\alpha 2$ band.

Taken together, our results cast new light on brain network reorganization occurring during BCI training and more generally during human learning.

References


- [1] Allison and C. Neuper, "Could Anyone Use a BCI?," in Brain-Computer Interfaces, Eds. Springer London, 2010,
- [2] Corsi, et al (2020). Functional disconnection of associative cortical areas predicts performance during BCI training. NeuroImage
- [3] Battiston et al. (2018) Multiplex core-periphery organization of the human connectome. Journal of The Royal Society Interface

Acknowledgements

This work was partially supported by the program "Investissements d'avenir" ANR-10-IAIHU-06; "ANR-NIH CRCNS" ANR-15-NEUC-0006-02 and by NICHD 1R01HD086888-01.

Relative coreness ΔC during training


To provide a more detailed description of the evolution of the relative coreness over training, we performed a one-way ANOVA for each layer separately.



ΔC presented a significant session effect involving different brain areas. Within the $\alpha 2$ range, a significant session effect was observed in EEG mostly within the long insular gyrus and the gyrus rectus; a significant session effect was observed in MEG in the supramarginal gyrus (working memory and motor planning); and in the multiplex a significant session effect was observed in areas involved during motor planning and working memory and in learning complex motor skills. In each case, we obtained an increase of ΔC with training.

Multiplex relative coreness correlated with future BCI performance

To assess whether relative coreness could be associated with future BCI performance, we estimated the correlation between ΔC in session i and the BCI score obtained in session $i + 1$.



We observed significant correlations only with multiplex within the $\alpha 2$ band. More precisely, a positive correlation ($p < 0.01$) was observed in the gyrus rectus, the subcentral gyrus, but also the long insular gyrus (involved during somatosensory tasks). A negative correlation was obtained in the superior occipital gyrus associated with visual processing.

Interested in this study?

Scan the QR code to get access to the associated paper!

