HAL open science

A Geometric Summation of the Geometric Series

Josué Tonelli-Cueto

To cite this version:

Josué Tonelli-Cueto. A Geometric Summation of the Geometric Series. The American Mathematical Monthly, 2022, pp.1-1. 10.1080/00029890.2022.2115825 . hal-03779492

HAL Id: hal-03779492

https://inria.hal.science/hal-03779492v1

Submitted on 16 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This is an Accepted Manuscript version of the following article, accepted for publication in *The American Mathematical Monthly*:

Josué Tonelli-Cueto (2022) A Geometric Summation of the Geometric Series, *The American Mathematical Monthly*, DOI: 10.1080/00029890.2022.2115825

It is deposited under the terms of the Creative Commons Attribution-NonCommercial License (`http://creativecommons.org/licenses/by-nc/4.0/`), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

A Geometric Summation of the Geometric Series

Let k be a finite field of size q. The n-dimensional projective space over k, $\mathbb{P}^n(k)$, is the set of orbits of $k^{n+1}\backslash\{0\}$ under the action of k^* given by scalar multiplication. Therefore $|\mathbb{P}^n(k)| = \frac{q^{n+1}-1}{q-1}$ since all orbits have the same size. Now, since the decomposition of $\mathbb{P}^n(k)$ into a disjoint union of affine spaces $\mathbb{P}^n(k) = A_0 \cup A_1 \cup \cdots \cup A_n$ where

$$A_i := \{(x_0 : \cdots : x_{i-1} : 1 : 0 : \cdots : 0) \,|\, x_j \in k\}$$

is in bijection with the affine space k^i, then $|\mathbb{P}^n(k)| = \sum_{i=0}^n q^i$. Thus for all n and power of primes q,

$$\sum_{i=0}^{n} q^i = \frac{q^{n+1}-1}{q-1}.$$

Since this identity of univariate rational functions over $\mathbb{Q}$ is true for an infinite amount of numbers, it translates into an equality of rational functions. Hence

$$\sum_{i=0}^{n} X^i = \frac{X^{n+1}-1}{X-1}$$

with X indeterminate. Can this method be used to prove other identities by interpreting them geometrically using algebraic varieties over finite fields?

—Submitted by Josué Tonelli-Cueto, Inria Paris & IMJ-PRG

MSC: Primary 05A19