



# Modelling the dynamics of Salmonella infection in the gut at the bacterial and the host level

Coralie Muller, Arie Wortsman, Pablo Ugalde-Salas, Clémence Frioux, Simon Labarthe

## ► To cite this version:

Coralie Muller, Arie Wortsman, Pablo Ugalde-Salas, Clémence Frioux, Simon Labarthe. Modelling the dynamics of Salmonella infection in the gut at the bacterial and the host level. Microbial communities: current approaches and open challenges, Oct 2022, Cambridge, United Kingdom. hal-03839405

**HAL Id: hal-03839405**

**<https://inria.hal.science/hal-03839405>**

Submitted on 4 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Modelling the dynamics of *Salmonella* infection in the gut at the bacterial and host levels

Coralie MULLER<sup>1</sup>, Arie WORTSMAN<sup>1</sup>, Pablo Andres UGALDES SALAS<sup>1</sup>, Clémence FRIOUX<sup>1</sup>, and Simon LABARTHE<sup>2,1</sup>

<sup>1</sup>Inria, Univ. Bordeaux, INRAE, F-33400 Talence, France

<sup>2</sup>INRAE, Univ. Bordeaux, BIOGECO, Pessac, France

**Keywords:** Gut microbiota - Metabolic network - Biological system - System of ODE - Metamodelling - Numerical metabolic modelling

## Context : a multidisciplinary project

We want to model the dynamics of a host-microbiota-pathogen system using the Dynamic Flux Balance Analysis method. dFBA integrates the fluxes calculated in FBA into a system of **ordinary differential equations (ODEs)**, enabling us to model the temporal dimension of metabolic simulations. In order to speed up the computation of the FBA models needed for a thorough numerical exploration, the metabolic models will be approximated by a **metamodel method [1]**.

Three main challenges:

curation, harmonization, optimization

Human colonocyte network

Three metabolites networks to harmonize and interconnect

Speed up the computation of the FBA models

Bioinformatics


Mathematics

Dynamic simulation fit better to the biological reality

Reduced simulation time

## Colonocyte metabolic network curation

A **manual curation** process was performed in order to clean up and make the model usable. It consists of **five main steps**:




## Simulation: FBA - dFBA

For the simulation, we consider that the system works by maximizing biomass production. Also, we assume that the system is in a stationary state. This way, can model the system by solving the linear **optimization problem**

$$\min_{v \in \mathbb{R}^{N_r}} v_{\text{biomass}} \quad A \cdot v = 0 \quad c_{\min} \leq v \leq c_{\max}$$

Our model assumes that FBA holds for each instant [2]. Then, for simulating it for a **period of time**, it is necessary to solve FBA on each compartment. After this timestep, an **ODE system models** the fluxes between different compartments, until the next step...



## Harmonization of the models

We model the host-microbiota-pathogen system by gathering three metabolic models: *Faecalibacterium prausnitzii* [3] as a representative butyrate producing **commensal bacterium**, *Salmonella Enterica* serovar *Thiphimurium* [4] for the **enteric pathogen**, these models previously used in CEMRACS[1] project, and a host genome scale metabolic network of human [5] called iHsa, to simulate the **epithelial cell**.

In order to make the simulation with the 3 models, we have **harmonized** the networks between them on **two points**:

Different IDs for the same metabolite and differentiate name for compartments between networks

Fix fluxes for essential reactions in the right compartments for the exchanges




## Simulation of growth in the human

We performed simulations with a custom medium to observe **biomass growth** as a function of O<sub>2</sub> and butyrate uptake.

We observed a **dependence of biomass growth on O<sub>2</sub> uptake** but **not on butyrate uptake**.

These results are not surprising because biomass includes many are **not directed related to the energy metabolism** and therefore **do not depend on butyrate**.




## Effect of but and O<sub>2</sub> on energy production

The **electron transport chain (ETC)** is the source of **energy production** of the cell, it is located in the **mitochondria** and **allows to produce at the end of the chain ATP**. To function, ETC needs in part **O<sub>2</sub>** and **FADH<sub>2</sub>** to enable electron transport. **FADH<sub>2</sub>** is a product of the **Krebs cycle reactions (TCA)** whose main reactant is **butyrate** in enterocytes. **Butyrate** is produced by the **beta oxidation of fatty acids** in the cell.



The **ATP synthase** is the main energy production reaction of the mitochondria and reflects the **energy production [6]** of the cell. With this activity, we want to **monitor the Salmonella infection**.



The butyrate/O<sub>2</sub> uptake has an impact on the ATP synthase activity, when the other uptake is stable.

The effect of O<sub>2</sub> uptake has a greater impact on ATP synthase activity than butyrate uptake.

## Next steps

Each time iteration of the model implies an optimization step in a high dimensional space, which means that the simulation has a very high computational cost.

Are there ways to solve this? **?** Is there a big trade-off between computation cost and precision?

Simulations involving the uptake of O<sub>2</sub> and butyrate on ATP synthase activity are encouraging. The **next steps** of the project will be:

- Simulate with dFBA to **validate** that the reduced of butyrate uptake lead to O<sub>2</sub> accumulation in the lumen [luLI].
- Learn the **behaviour** of the system and **build a metamodel** able to efficiently approximate the dynamics.
- Generalize the approach to other ecosystems **which functions** are important for dynamics of the **community**.

## References

- Clémence Frioux et al. Accelerating metabolic models evaluation with statistical metamodels: application to *Salmonella* infection models. working paper or preprint, April 2022.
- Ove Oyås and Jörg Satellin. Genome-scale metabolic networks in time and space. *Current Opinion in Systems Biology*, 8:51-8, 2018.
- El-Semman et al. "Genome-scale metabolic reconstructions of *Bifidobacterium adolescentis* L2-32 and *Faecalibacterium prausnitzii* A2-165 and their interaction." *BMC systems biology* 8.1 (2014): 1-11.
- Raghuathan et al. Constraint-based analysis of metabolic capacity of *Salmonella typhimurium* during host-pathogen interaction. *BMC Syst Biol* 3, 38 (2009). <https://doi.org/10.1186/1752-0509-3-38>
- Kocabaş, K., Arif, A., Uddin, R., & Cakir, T. (2022). Dual transcriptome based reconstruction of salmonella human integrated metabolic network to screen potential drug targets. *PLOS ONE*, 17, e0268889 <https://doi.org/10.1371/JOURNAL.PONE.0268889>
- Schuetz, R., Kuepfer, L., & Sauer, U. (2007). Systematic evaluation of objective functions for predicting intracellular fluxes in *Escherichia coli*. *Molecular Systems Biology*, 3, 119. <https://doi.org/10.1038/MSB4100162>