



**HAL**  
open science

## **Surgical management of insulinoma over three decades**

Anne de Carbonnières, Alexandre Challine, Anne Ségolène Cottureau, Romain Coriat, Philippe Soyer, Einas Abou Ali, Frédéric Prat, Benoit Terris, Jérôme Bertherat, Bertrand Dousset, et al.

► **To cite this version:**

Anne de Carbonnières, Alexandre Challine, Anne Ségolène Cottureau, Romain Coriat, Philippe Soyer, et al.. Surgical management of insulinoma over three decades. *HPB*, 2021, 23 (12), pp.1799-1806. 10.1016/j.hpb.2021.04.013 . hal-03864941

**HAL Id: hal-03864941**

**<https://inria.hal.science/hal-03864941>**

Submitted on 5 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## **Surgical management of insulinoma over three decades**

A. de Carbonnières<sup>1,2</sup>; A. Challine<sup>1,2</sup>; A.S. Cottereau<sup>2,3</sup>; R. Coriat<sup>2,4</sup>; P. Soyer<sup>2,5</sup>; E. Abou Ali<sup>2,4</sup>; F. Prat<sup>2,4</sup>; B. Terris<sup>2,6</sup>; J. Bertherat<sup>2,7</sup>; B. Dousset<sup>1,2</sup>; S. Gaujoux<sup>1,2</sup>

*ADC and AC contributed equally to the work*

1. Department of Digestive, Hepato-biliary and Endocrine Surgery, Cochin Hospital, APHP, 75014 Paris, France
2. Université de Paris, 75006 Paris, France
3. Department of Nuclear Medicine, Cochin Hospital, APHP, 75014 Paris, France
4. Department of Gastroenterology, Cochin Hospital, APHP, 75014 Paris, France
5. Department of Radiology, Cochin Hospital, APHP, 75014 Paris, France
6. Department of Pathology, Cochin Hospital, APHP, 75014 Paris, France
7. Department of Endocrinology, Cochin Hospital, APHP, 75014 Paris, France

**Corresponding author:**

Pr Sébastien Gaujoux, MD, PhD

Department of Digestive, Hepato-biliary and Endocrine Surgery, Cochin Hospital, APHP,

Université Paris Descartes, Paris, France

27, rue du Faubourg Saint Jacques, 75014, Paris, France.

Professional number: +33 1 58 41 29 61

Fax number: +33 1 58 41 29 58

E-mail: [sebastien.gaujoux@aphp.fr](mailto:sebastien.gaujoux@aphp.fr)

**Key words**

Insulinoma; Pancreas; Pancreatectomy; Laparoscopy; Tomography; X-Ray computed;

**Parenchyma-sparing pancreatectomy;** Enucleation; Neuroendocrine tumours; Postoperative complications.

## Abstract

**Introduction:** This paper reports our experience of the perioperative management of patients with sporadic, non-malignant, pancreatic insulinoma.

**Methods:** A retrospective monocentric cohort study was performed from January 1989 to July 2019 using a prospectively maintained database, including all the patients who had been operated on for pancreatic insulinoma as of the time of the study. The preoperative work-up, surgical management, and postoperative outcome were analyzed.

**Results:** Eighty patients underwent surgery for sporadic pancreatic insulinoma, 50 of which were female (62%), with a median age of 50 (36–70) years. Preoperatively, the tumors were localized in 76 patients (95%). Computed tomography (CT) and magnetic resonance imaging allowed exact preoperative tumor localization in 76% of the patients (64–85 and 58–88 patients, respectively), increasing to 96% when endoscopic ultrasonography was performed in addition to CT. Forty-one **parenchyma-sparing pancreatectomies** (including enucleation, **caudal** pancreatectomy, and uncinata process resection) and 39 pancreatic resections were performed, 32% via laparoscopy. The mortality rate was 6% (n = 5, 4 after pancreaticoduodenectomy), with a morbidity rate of 72%, including 24 severe complications (30%) and 35 pancreatic fistulas (44%). No differences were found between **formal pancreatectomy** and **parenchyma-sparing pancreatectomy** in terms of postoperative outcome procedures. The surgery was curative in all the patients, with long-term postoperative normalization of the glycemia and insulin blood levels.

**Conclusion:** CT used in combination with endoscopic ultrasonography allows accurate localization of insulinomas in almost all patients. When possible, a **parenchyma-sparing pancreatectomy** should be proposed as the first-line surgical strategy.

## Introduction

Insulinoma is the most common functional pancreatic neuroendocrine tumor. In most patients, it is a small, sporadic, solitary, and intrapancreatic benign tumor,<sup>1-3</sup> but its diagnosis can be difficult and is often delayed because of the highly variable and non-specific symptoms of hypoglycemia. When a diagnosis of organic hypoglycemia **with inappropriately elevated endogenously produced insulin** after a 48–72 h fasting test<sup>4</sup> is confirmed, the next issue is to accurately localize the insulinoma to anticipate the surgical approach, which has long been exclusively based on intraoperative ultrasonography and thorough pancreatic palpation. This is often challenging because the tumors distributed throughout the pancreatic gland are usually small. The combination of cross-sectional imaging techniques, nuclear medicine, and endoscopic ultrasonography, however, has dramatically changed the preoperative assessment of insulinomas.<sup>5-7</sup> Thus, blind pancreatectomy is no longer recommended.<sup>8</sup> Surgery remains the cornerstone of insulinoma treatment.<sup>9</sup> As it is benign in most patients, it should ideally be done through **parenchyma-sparing pancreatectomy**<sup>10</sup> ideally enucleation<sup>11,12</sup> to preserve both the exocrine and endocrine functions.<sup>1</sup> Nevertheless, these procedures often expose patients to substantial postoperative morbidity.<sup>13</sup>

Reported herein are **the evolution** in the pre- and perioperative management of patients with sporadic non-malignant insulinoma and our experience in relation to such, with specific attention given to preoperative localization and postoperative outcome following pancreatectomy **and the development of the laparoscopic approach.**

## Methods

### Patients and methods

#### *Data collection*

All the patients who underwent pancreatectomy for sporadic non-malignant insulinoma within the period from 1989 to 2019 at Cochin Hospital, Paris, France, a tertiary referral center for endocrine diseases, were included in this study. Demographic, clinical, radiological, pathologic, and follow-up data were obtained from a prospectively maintained database, with additional retrospective review. For each patient, the diagnosis was confirmed by an experienced pathologist. Malignant (i.e., metastatic insulinoma) and multiple endocrine neoplasia type 1 (MEN1) insulinoma were excluded from the analysis. Postoperative mortality included all the deaths that occurred within 90 days from the surgery. Morbidity included all the complications that occurred within 90 days from the surgery and was graded according to the Clavien-Dindo classification<sup>13</sup>. Postoperative pancreatic fistula, hemorrhage, and delayed gastric emptying were defined according to the International Study Group of Pancreatic Surgery<sup>14-17</sup>. The follow-up was based on clinical, radiological, and laboratory assessments and was updated on the basis of outpatient reviews and correspondence. All the patients were followed up at least 1 year after surgery. The year 2000 was considered a time landmark because it marked the beginning of the evolution of imaging, endoscopic, and laparoscopic techniques. The study protocol was approved by the hospital's medical ethics committee.

#### *Preoperative work-up and surgical indications*

The preoperative work-up (all performed in Cochin Hospital) included a clinical evaluation with patient questioning and a complete hormonal and radiological evaluation with triphasic computed tomography (CT) examination (unenhanced, arterial, and portal phases)

and endoscopic ultrasound (EUS) since it became available. Additional investigations with magnetic resonance imaging (MRI) including standard sequences and cholangiopancreatography images and somatostatin receptor scintigraphy or positron emission tomography (PET)/CT either with  $^{111}\text{In}$ -DTPA-octreotide or more recently  $^{68}\text{Ga}$ -DOTATOC or EUS with fine needle aspiration (FNA) were left to the surgeon's discretion. The policy regarding the use of nuclear medicine imaging has evolved over time.  $^{111}\text{In}$ -DTPA-octreotide, despite its low sensibility, was initially used but was later replaced by  $^{68}\text{Ga}$ -DOTATOC when it became more easily available. All surgical indications were discussed in a multidisciplinary tumor board that included surgeons, radiologists, nuclear physicians, endocrinologists, pathologists, and gastroenterologists.

### *Surgical techniques*

The type of surgical resection that was performed was based on the tumor location and the distance to the main pancreatic duct. Distal pancreatectomy was defined as a pancreatectomy with pancreatic resection at the level of the isthmus. Caudal pancreatectomy, on the other hand, which is considered parenchyma-sparing pancreatectomy, was defined as a pancreatectomy with pancreatic resection on the left side of the isthmus. Enucleation and parenchyma-sparing pancreatectomy (central pancreatectomy, resection of the uncinate process, and caudal pancreatectomy) was preferred whenever possible. During surgery, pancreatic exploration after mobilization of the transverse colon root was first performed through visual inspection and surgical ultrasonography to confirm the tumor location.

### *Statistical analysis*

The variables were expressed as medians, interquartiles, or numbers and percentages of patients, as appropriate. The continuous variables were evaluated for normal distribution

using the Komogorov-Smirnov test. Chi square or Fisher's exact test was used to evaluate the categorical variables. The continuous variables were analyzed using Student's t-test or Mann-Whitney test, as appropriate. The capability of each imaging test to determine the exact tumor location was calculated in terms of the true location rate, using the surgical and histopathological findings as the standards of reference. The McNemar test was used to compare the diagnostic tests. All the tests were two-tailed, and significance was set at  $p < 0.05$ . The analysis was conducted with the R software.<sup>14</sup>

## Results

### *Characteristics of the population and preoperative imaging*

The final study population included 80 patients who underwent surgery for pancreatic insulinoma, 50 (62%) of which were women and 30 (38%) men, with a median age of 50 (36–70) years (**Table 1**). Most of the patients were ASA 1 or 2 (n = 76; 95%) and presented neuroglycopenic symptoms (n = 74; 92%), including fainting (40%; n = 32). The median time interval between the first symptoms and the surgery was 13 months (4–26 months).

The proportions of tumors that were correctly localized using the different preoperative imaging tests are shown in **Table 2**. The exact preoperative location was obtained in 76/80 patients (95%), and the preoperative tumor location was accurate in all the patients who underwent CT, MRI, EUS, and somatostatin receptor imaging (19/19; 100%; 95% confidence interval [CI]: 79–100). When CT was used in combination with EUS, an accurate preoperative tumor location was obtained in 65/68 patients (96%; 95% CI: 87–99). The results of the other combinations are given in **Table 2**. CT, MRI, and EUS showed more accurate preoperative tumor locations (p = 0.009, 0.01, and 0.02, respectively) than octreoscan, with only four DOTAPETs realised after the year 2000 and only one positive for a 7 mm lesion. There were no differences in size between the negative and positive octreoscans and before and after the year 2000. No significant differences were found between CT, MRI, and EUS. Four patients were operated on without a preoperative location, all before 1995. For the lesion in the tail of the pancreas detected via intraoperative ultrasonography or palpation, none of the patients had preoperative MRI or EUS.

### *Surgical procedure*

In most cases, the actual surgery follows the preoperative planning. In 2 cases in this study, however, the planned enucleation of a large cephalic lesion was not possible because of

the intraoperative lateral wound in the main pancreatic duct; as such, pancreaticoduodenectomy was performed instead. Overall, 25/50 patients (31%) underwent surgery via the laparoscopic approach, and 5 (20%) required conversion to accurately locate and resect the tumor. Twelve (48%) patients underwent pancreatic enucleation; 6 (40%), distal pancreatectomy; 3 (12%), distal pancreatectomy with splenectomy; and 4 (16%), caudal pancreatectomy. In addition, an open approach was adopted in 55/80 patients (69%); among them, 24/55 (44%) had pancreatic enucleation; 9/55 (16%), pancreaticoduodenectomy; 15/55 (27%), distal pancreatectomy; 6/55 (11%), distal pancreatectomy with splenectomy; and 1/55 (2%), resection of the uncinate process. Regarding the surgical approach, 34/80 patients (43%) had a tumor located in the head of the pancreas, 32/34 (94%) of whom were operated on using the open approach whereas 22/46 patients (48%) with tumors in other locations underwent laparoscopy ( $p < 0.001$ ). In summary, 41 parenchyma-sparing pancreatectomies, including enucleation, caudal pancreatectomy, and uncinate process resection, were realised, with a 41% laparoscopic rate. The rate of laparoscopy was 28% before the year 2000 and increased to 55% after 2000 ( $p = 0.09$ ) for the patients who underwent parenchyma-sparing pancreatectomy. In all the cases, the pancreas was considered soft.

### *Postoperative outcomes*

The postoperative course is summarised in **Table 3**. The mortality rate was 6% (5/80), including 1 death from septic shock from the venous central catheter, 2 from grade C pancreatic fistula (bleeding and septic shock), 1 from liver abscesses due to hepatic artery thrombosis, and 1 from fungal septicemia. Four of these deaths occurred after pancreaticoduodenectomy. The morbidity rate was 72% ( $n = 58$ ), including 24 severe (grade III/IV or V) complications (30%), and a pancreatic fistula occurred in 35 patients (44%), with 19 (24%) grade B or C. No significant differences were found between the patients who

underwent **parenchyma-sparing pancreatectomy** and those who underwent **formal pancreatectomy**. The surgery was successful in all the patients, with long-term postoperative normalization of the glycemia and insulin blood levels. One patient presented a distant liver recurrence 6 years after the initial pancreatic resection of a 2 cm insulinoma.

#### *Evolution over time*

The use of MRI increased after 2000 (< 2000: 24% vs. > 2000: 71%;  $p < 0.001$ ) along with the rate of detected lesions (< 2000: 93% vs. > 2000: 97%;  $p = 0.6$ ). The rate of laparoscopy also increased after 2000 (< 2000: 22% vs. > 2000: 41%;  $p = 0.1$ ). The severe complications (death, severe complication, or severe leakage) did not significantly differ between the two study periods.

## Discussion

Reported herein is the pre- and perioperative management of patients with sporadic non-malignant insulinoma over three decades, and several questions were raised. Once the hormonal diagnosis of organic hypoglycemia has been confirmed by a 72 h fasting test, the main challenge for the surgeon is *to accurately determine the tumor location*. The second challenge is *to define the best surgical strategy* with regard to the type of resection (**formal pancreatectomy** vs. **parenchyma-sparing pancreatectomy**) and the surgical approach (laparotomy vs. laparoscopy). The final objective is *to decrease the perioperative morbidity and minimize the late functional consequences of surgery*.

Accurate preoperative localization of insulinoma is one of the main issues in the management of this tumor. Indeed, insulinomas are small lesions (with 84% typically < 2 cm) randomly distributed in a fragile retroperitoneal organ<sup>6</sup>. Our study reported high accuracy in the detection and localization of insulinoma. We previously reported that dual-phase thin-section multidetector CT showed 93% localization sensitivity<sup>5</sup>. The performances of MRI reported in our study are also in line with those reported in previous studies<sup>15,16</sup> and had been increased by the addition of diffusion-weighted MRI sequences<sup>15,17</sup>. EUS has also been a major improvement in insulinoma location, with very high sensibility and specificity<sup>15,18</sup>. **Nevertheless, as the results of the other investigation methods are usually available when EUS is done, the investigator is often helped by such results.**

EUS also allows contrast-enhanced ultrasonography, which began to be used in our department only after 2010, improving the localization of the pancreatic mass to approximately 8–10%<sup>19,20</sup> and providing additional information about the vascularization of the tumor, FNA or biopsy (uncommonly needed for insulinoma), or EUS fiducial placement for an accurate intraoperative location of smaller lesions<sup>15,18,21</sup>. Nuclear medicine has also improved the location and characterization of insulinoma since the disappointing sensitivity of

<sup>111</sup>In-pentetreotide somatostatin receptor scintigraphy. Other somatostatin receptor imaging methods, such as <sup>68</sup>Ga-DOTATATE PET/CT, have shown sufficiently high sensitivity for identifying most insulinomas, and are considered by some adjunct imaging studies when the previous imaging findings were negative<sup>7</sup>. <sup>18</sup>F-FDOPA PET/CT with carbidopa premedication has also shown interesting results, accurately **locating more than 80% of lesions when combined with carbidopa premedication and early pancreatic acquisitions**<sup>22,23</sup>. **The policy regarding the use of nuclear-medicine imaging has evolved over time**, and more recently, <sup>68</sup>Ga-NOTA-exendin-4 PET/CT scan has shown high sensitivity and specificity for the detection and localization of insulinoma<sup>7,24</sup>. With modern imaging tools combining CT, MRI, EUS, and nuclear medicine, insulinoma detection and localization can be adequately achieved preoperatively in nearly 100% of patients. In this setting, blind pancreatectomy or exploratory laparotomy is no longer advised<sup>8</sup>.

When the lesion has been identified, the next step for the surgical management of insulinoma is to define the best surgical procedure (**parenchyma-sparing pancreatectomy vs. formal pancreatectomy**) and surgical approach (laparoscopy vs. laparotomy). It is important to emphasize that the open approach to pancreatectomy remains the standard of care, and that there is no clear consensus regarding the indications for a minimally invasive approach in pancreatic surgery. Regarding pancreaticoduodenectomy, the laparoscopic approach did not show any clear advantage over open pancreaticoduodenectomy<sup>25</sup>. Additionally, a Dutch multicentric randomized controlled study comparing laparoscopic and open pancreaticoduodenectomy was prematurely stopped because of increased mortality in the laparoscopic arm<sup>26</sup>. In contrast, pancreatectomy without reconstruction (i.e., distal **or caudal** pancreatectomy and enucleation) remains the best procedure for insulinoma when feasible. Several studies and meta-analyses have reported that laparoscopic distal pancreatectomy is a safe and effective procedure associated with a lower rate of complications and a shorter length

of hospital stay compared to open surgery<sup>25,26</sup>, but very few of these studies specifically focused on pancreas neuroendocrine tumors (PNETs). Similarly, laparoscopic enucleation is safe and effective for the treatment of benign and borderline pancreatic neoplasms<sup>27</sup>, as confirmed by a meta-analysis of more than 400 patients, which showed that a minimally invasive approach does not induce a higher risk of pancreatic fistula and offers numerous advantages over the conventional surgery for pancreatic enucleation<sup>28</sup>. This, along with the increasing relevant technical skills and experience, explains the more frequent use of the laparoscopic approach for **parenchyma-sparing pancreatectomy**. The choice of surgical technique between **parenchyma-sparing pancreatectomy** and **formal pancreatectomy** for the treatment of insulinoma is a crucial issue. The decision should be made according to anatomical considerations (size and location) and should take into account the distance between the lesion and the main pancreatic duct, which should be greater than 2–3 mm<sup>8</sup>. Nevertheless, **parenchyma-sparing pancreatectomy** has a similar postoperative mortality rate as and a higher risk of developing pancreatic fistula than **formal pancreatectomy**, especially when performed on a soft pancreas<sup>29</sup>. Moreover, it is associated with excellent long-term functional results<sup>29</sup>. In the present study, its overall morbidity rate was high, with 30% severe complications, 24% grade B or C pancreatic fistulas, and a 6% postoperative death rate, mainly observed after pancreaticoduodenectomy, with **an unusually high mortality rate. This underlines the risk posed by pancreatic surgery on a soft pancreas without main-pancreatic-duct dilatation even if performed in young and fit patients.** The **high** morbidity in recent years might have also been the consequence of the greater use of laparoscopic **parenchyma-sparing pancreatectomy**.

The major challenge for pancreatic surgeons is the significant rate of postoperative complications. In large surgical series for insulinoma, the reported mortality rate is 0–6%, with a 33–56% morbidity rate, whereas the mortality rate of pancreatic surgery ranges from

1% to 3% in high-volume centers<sup>38,39</sup>, increasing to 6–10% when nationwide centers are considered<sup>40–42</sup>. Considering these heterogeneous results, being treated in expert-organized high-volume centers presents a clear benefit for patients<sup>40–42</sup>. Considering that the mortality rate of pancreatectomy is mainly related to pancreaticoduodenectomy, other therapeutic options have emerged in the last few years. From a technical view point, the prophylactic endoscopic placement of a pancreatic stent for “deep” pancreatic enucleations<sup>43</sup> may help extend the surgical indications for pancreatic enucleation, but this advantage may be overshadowed by the risk of developing post-endoscopic retrograde cholangiopancreatography pancreatitis. Controlled studies, although desirable, are unlikely to be feasible due to the low incidence of such lesions. If surgical resection has been advocated for years as the only curative option for insulinoma, local destruction of tumors through radiofrequency ablation (RFA) may be proposed as an alternative despite the fact that very few such procedures have been performed to date. In the literature, only about 60 patients with PNETs have been treated with RFA<sup>44–53</sup>, but it has a clinical success rate of 83.3–100%<sup>44–53</sup>. Nevertheless, the morbidity rate, technical constraints, and long-term results of this emerging treatment should be investigated in depth. Considering the experience with RFA so far, we believe that local ablation of tumors through it may be considered a suitable alternative for insulinomas located in the head of the pancreas.

The main limitation of this study is its retrospective design spanning over three decades due to the low incidence of insulinoma. This long inclusion period saw improvements in imaging techniques and perioperative management with the development of the laparoscopic approach. Even though this study’s sample consisting of 80 patients operated on for insulinoma may be viewed as restricted, this study was one of the largest monocentric studies that have been conducted to date, providing an interesting overview of the management of the rare disease of insulinoma<sup>8,16,18,30–34</sup>.

In conclusion, the management of insulinoma has evolved dramatically over the recent decades with the emergence of modern imaging techniques, including cross-sectional imaging, EUS, and nuclear medicine. As such, accurate preoperative detection and localization is obtained in most patients. A major challenge remains, however: the significant postoperative morbidity rate, which is **high** after **parenchyma-sparing pancreatectomy**, counterbalancing the long-term benefits on both the exocrine and endocrine functions.

## References

1. Crippa S, Zerbi A, Boninsegna L, et al. Surgical management of insulinomas: short- and long-term outcomes after enucleations and pancreatic resections. *Arch Surg Chic Ill 1960*. 2012;147(3):261-266. doi:10.1001/archsurg.2011.1843
2. Tucker ON, Crotty PL, Conlon KC. The management of insulinoma. *Br J Surg*. 2006;93(3):264-275. doi:10.1002/bjs.5280
3. Davì MV, Falconi M. Pancreas: Insulinoma--new insights into an old disease. *Nat Rev Endocrinol*. 2009;5(6):300-302. doi:10.1038/nrendo.2009.99
4. Guettier J-M, Lungu A, Goodling A, Cochran C, Gorden P. The role of proinsulin and insulin in the diagnosis of insulinoma: a critical evaluation of the Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2013;98(12):4752-4758. doi:10.1210/jc.2013-2182
5. Gouya H, Vignaux O, Augui J, et al. CT, Endoscopic Sonography, and a Combined Protocol for Preoperative Evaluation of Pancreatic Insulinomas. *Am J Roentgenol*. 2003;181(4):987-992. doi:10.2214/ajr.181.4.1810987
6. Mehrabi A, Fischer L, Hafezi M, et al. A Systematic Review of Localization, Surgical Treatment Options, and Outcome of Insulinoma. *Pancreas*. 2014;43(5):675-686. doi:10.1097/MPA.000000000000110
7. Nockel P, Babic B, Millo C, et al. Localization of Insulinoma Using <sup>68</sup> Ga-

DOTATATE PET/CT Scan. *J Clin Endocrinol Metab.* Published online November 2, 2016;jc.2016-3445. doi:10.1210/jc.2016-3445

8. Zhao Y, Zhan H, Zhang T, et al. Surgical management of patients with insulinomas: Result of 292 cases in a single institution: Surgical Management of Insulinoma. *J Surg Oncol.* 2011;103(2):169-174. doi:10.1002/jso.21773

9. Placzkowski KA, Vella A, Thompson GB, et al. Secular trends in the presentation and management of functioning insulinoma at the Mayo Clinic, 1987-2007. *J Clin Endocrinol Metab.* 2009;94(4):1069-1073. doi:10.1210/jc.2008-2031

10. Goudard Y, Gaujoux S, Dokmak S, et al. Reappraisal of central pancreatectomy a 12-year single-center experience. *JAMA Surg.* 2014;149(4):356-363. doi:10.1001/jamasurg.2013.4146

11. Cherif R, Gaujoux S, Sauvanet A. Enucleation of pancreatic lesions through laparotomy. *J Visc Surg.* 2012;149(6):395-399. doi:10.1016/j.jviscsurg.2012.10.014

12. Faitot F, Gaujoux S, Barbier L, et al. Reappraisal of pancreatic enucleations: A single-center experience of 126 procedures. *Surgery.* 2015;158(1):201-210. doi:10.1016/j.surg.2015.03.023

13. Strobel O, Cherrez A, Hinz U, et al. Risk of pancreatic fistula after enucleation of pancreatic tumours. *Br J Surg.* 2015;102(10):1258-1266. doi:10.1002/bjs.9843

14. R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
15. Okabayashi T. Diagnosis and management of insulinoma. *World J Gastroenterol*. 2013;19(6):829. doi:10.3748/wjg.v19.i6.829
16. Thoeni RF, Mueller-Lisse UG, Chan R, Do NK, Shyn PB. Detection of Small, Functional Islet Cell Tumors in the Pancreas: Selection of MR Imaging Sequences for Optimal Sensitivity. *Radiology*. 2000;214(2):483-490. doi:10.1148/radiology.214.2.r00fe32483
17. Kulali F, Semiz-Oysu A, Demir M, Segmen-Yilmaz M, Bukte Y. Role of diffusion-weighted MR imaging in predicting the grade of nonfunctional pancreatic neuroendocrine tumors. *Diagn Interv Imaging*. 2018;99(5):301-309. doi:10.1016/j.diii.2017.10.012
18. Goh BKP, Ooi LLPJ, Cheow P-C, et al. Accurate preoperative localization of insulinomas avoids the need for blind resection and reoperation: analysis of a single institution experience with 17 surgically treated tumors over 19 years. *J Gastrointest Surg Off J Soc Surg Aliment Tract*. 2009;13(6):1071-1077. doi:10.1007/s11605-009-0858-y
19. Gincul R, Palazzo M, Pujol B, et al. Contrast-harmonic endoscopic ultrasound for the diagnosis of pancreatic adenocarcinoma: a prospective multicenter trial. *Endoscopy*. 2014;46(5):373-379. doi:10.1055/s-0034-1364969
20. Fusaroli P, Spada A, Mancino MG, Caletti G. Contrast harmonic echo-endoscopic

ultrasound improves accuracy in diagnosis of solid pancreatic masses. *Clin Gastroenterol Hepatol Off Clin Pract J Am Gastroenterol Assoc.* 2010;8(7):629-634.e1-2. doi:10.1016/j.cgh.2010.04.012

21. Gaujoux S, Genc S, Leblanc S, Dousset B, Prat F. Insulinoma enucleation after echoendoscopic fiducial placement. *Gastrointest Endosc.* 2018;87(2):615-616. doi:10.1016/j.gie.2017.08.005

22. Imperiale A, Sebag F, Vix M, et al. 18F-FDOPA PET/CT imaging of insulinoma revisited. *Eur J Nucl Med Mol Imaging.* 2015;42(3):409-418. doi:10.1007/s00259-014-2943-z

23. Leroy-Freschini B, Amodru V, Addeo P, et al. Early 18F-FDOPA PET/CT imaging after carbidopa premedication as a valuable diagnostic option in patients with insulinoma. *Eur J Nucl Med Mol Imaging.* 2019;46(3):686-695. doi:10.1007/s00259-018-4245-3

24. Luo Y, Pan Q, Yao S, et al. Glucagon-Like Peptide-1 Receptor PET/CT with 68Ga-NOTA-Exendin-4 for Detecting Localized Insulinoma: A Prospective Cohort Study. *J Nucl Med Off Publ Soc Nucl Med.* 2016;57(5):715-720. doi:10.2967/jnumed.115.167445

25. Drymoussis P, Raptis DA, Spalding D, et al. Laparoscopic versus open pancreas resection for pancreatic neuroendocrine tumours: a systematic review and meta-analysis. *HPB.* 2014;16(5):397-406. doi:10.1111/hpb.12162

26. Han SH, Han IW, Heo JS, et al. Laparoscopic versus open distal pancreatectomy for nonfunctioning pancreatic neuroendocrine tumors: a large single-center study. *Surg Endosc.*

2018;32(1):443-449. doi:10.1007/s00464-017-5702-y

27. Song KB, Kim SC, Hwang DW, et al. Enucleation for benign or low-grade malignant lesions of the pancreas: Single-center experience with 65 consecutive patients. *Surgery*. 2015;158(5):1203-1210. doi:10.1016/j.surg.2014.10.008

28. Guerra F, Giuliani G, Bencini L, Bianchi PP, Coratti A. Minimally invasive versus open pancreatic enucleation. Systematic review and meta-analysis of surgical outcomes. *J Surg Oncol*. 2018;117(7):1509-1516. doi:10.1002/jso.25026

29. Cherif R, Gaujoux S, Couvelard A, et al. Parenchyma-Sparing Resections for Pancreatic Neuroendocrine Tumors. *J Gastrointest Surg*. 2012;16(11):2045-2055. doi:10.1007/s11605-012-2002-7

30. Cunha AS, Beau C, Rault A, Catargi B, Collet D, Masson B. Laparoscopic versus open approach for solitary insulinoma. *Surg Endosc*. 2007;21(1):103-108. doi:10.1007/s00464-006-0021-8

31. Daneshvar K, Grenacher L, Mehrabi A, Kauczor H-U, Hallscheidt P. Preoperative tumor studies using MRI or CT in patients with clinically suspected insulinoma. *Pancreatol Off J Int Assoc Pancreatol IAP Al*. 2011;11(5):487-494. doi:10.1159/000330208

32. Morera J, Guillaume A, Courtheoux P, et al. Preoperative localization of an insulinoma: selective arterial calcium stimulation test performance. *J Endocrinol Invest*. 2016;39(4):455-463. doi:10.1007/s40618-015-0406-4

33. Sotoudehmanesh R, Hedayat A, Shirazian N, et al. Endoscopic ultrasonography (EUS) in the localization of insulinoma. *Endocrine*. 2007;31(3):238-241. doi:10.1007/s12020-007-0045-4
34. Peltola E, Hannula P, Huhtala H, et al. Characteristics and Outcomes of 79 Patients with an Insulinoma: A Nationwide Retrospective Study in Finland. *Int J Endocrinol*. 2018;2018:1-10. doi:10.1155/2018/2059481
35. Tian F, Hong X-F, Wu W-M, et al. Propensity score-matched analysis of robotic versus open surgical enucleation for small pancreatic neuroendocrine tumours. *Br J Surg*. 2016;103(10):1358-1364. doi:10.1002/bjs.10220
36. Xu J, Li F, Zhan H, et al. Laparoscopic enucleation of pancreatic tumours: a single-institution experience of 66 cases. *ANZ J Surg*. 2021;91(1-2):106-110. doi:10.1111/ans.16450
37. Qu Y, Li H, Wang X, et al. Clinical Characteristics and Management of Functional Pancreatic Neuroendocrine Neoplasms: A Single Institution 20-Year Experience with 286 Patients. *Int J Endocrinol*. 2020;2020:1030518. doi:10.1155/2020/1030518
38. Cameron JL, Riall TS, Coleman J, Belcher KA. One thousand consecutive pancreaticoduodenectomies. *Ann Surg*. 2006;244(1):10-15. doi:10.1097/01.sla.0000217673.04165.ea
39. Cameron JL, He J. Two thousand consecutive pancreaticoduodenectomies. *J Am Coll*

*Surg.* 2015;220(4):530-536. doi:10.1016/j.jamcollsurg.2014.12.031

40. Swanson RS, Pezzi CM, Mallin K, Loomis AM, Winchester DP. The 90-Day Mortality After Pancreatectomy for Cancer Is Double the 30-Day Mortality: More than 20,000 Resections From the National Cancer Data Base. *Ann Surg Oncol.* 2014;21(13):4059-4067. doi:10.1245/s10434-014-4036-4

41. El amrani M, Clément G, Lenne X, et al. Should all pancreatic surgery be centralized regardless of patients' comorbidity? *HPB.* Published online November 2019:S1365182X19331879. doi:10.1016/j.hpb.2019.10.2443

42. El Amrani M, Lenne X, Clement G, et al. Specificity of Procedure volume and its Association With Postoperative Mortality in Digestive Cancer Surgery: A Nationwide Study of 225,752 Patients. *Ann Surg.* 2019;270(5):775-782. doi:10.1097/SLA.0000000000003532

43. Giuliani T, Marchegiani G, Girgis MD, et al. Endoscopic placement of pancreatic stent for “Deep” pancreatic enucleations operative technique and preliminary experience at two high-volume centers. *Surg Endosc.* 2020;34(6):2796-2802. doi:10.1007/s00464-020-07501-y

44. Rossi S, Viera FT, Ghittoni G, et al. Radiofrequency ablation of pancreatic neuroendocrine tumors: a pilot study of feasibility, efficacy, and safety. *Pancreas.* 2014;43(6):938-945. doi:10.1097/MPA.000000000000133

45. Pai M, Habib N, Senturk H, et al. Endoscopic ultrasound guided radiofrequency ablation, for pancreatic cystic neoplasms and neuroendocrine tumors. *World J Gastrointest*

*Surg.* 2015;7(4):52-59. doi:10.4240/wjgs.v7.i4.52

46. Limmer S, Huppert PE, Juette V, Lenhart A, Welte M, Wietholtz H. Radiofrequency ablation of solitary pancreatic insulinoma in a patient with episodes of severe hypoglycemia. *Eur J Gastroenterol Hepatol.* 2009;21(9):1097-1101. doi:10.1097/meg.0b013e328323d70e

47. Armellini E, Crinò SF, Ballarè M, Occhipinti P. Endoscopic ultrasound-guided radiofrequency ablation of a pancreatic neuroendocrine tumor. *Endoscopy.* 2015;47 Suppl 1 UCTN:E600-601. doi:10.1055/s-0034-1393677

48. Barthet M, Giovannini M, Lesavre N, et al. Endoscopic ultrasound-guided radiofrequency ablation for pancreatic neuroendocrine tumors and pancreatic cystic neoplasms: a prospective multicenter study. *Endoscopy.* 2019;51(9):836-842. doi:10.1055/a-0824-7067

49. Waung JA, Todd JF, Keane MG, Pereira SP. Successful management of a sporadic pancreatic insulinoma by endoscopic ultrasound-guided radiofrequency ablation. *Endoscopy.* 2016;48 Suppl 1:E144-145. doi:10.1055/s-0042-104650

50. Lakhtakia S, Ramchandani M, Galasso D, et al. EUS-guided radiofrequency ablation for management of pancreatic insulinoma by using a novel needle electrode (with videos). *Gastrointest Endosc.* 2016;83(1):234-239. doi:10.1016/j.gie.2015.08.085

51. Choi J-H, Seo D-W, Song TJ, et al. Endoscopic ultrasound-guided radiofrequency ablation for management of benign solid pancreatic tumors. *Endoscopy.* 2018;50(11):1099-

1104. doi:10.1055/a-0583-8387

52. Bas-Cutrina F, Bargalló D, Gornals JB. Small pancreatic insulinoma: Successful endoscopic ultrasound-guided radiofrequency ablation in a single session using a 22-G fine needle. *Dig Endosc Off J Jpn Gastroenterol Endosc Soc.* 2017;29(5):636-638. doi:10.1111/den.12866

53. Procházka V, Hlavsa J, Andrašina T, et al. Laparoscopic radiofrequency ablation of functioning pancreatic insulinoma: video case report. *Surg Laparosc Endosc Percutan Tech.* 2012;22(5):e312-315. doi:10.1097/SLE.0b013e318264b607

## Table and figures

**Table 1:** Characteristics of the study population (n = 80)

Characteristics		Median [IQR]; (range) or n (%)
Comorbidity	Age (years)	50 [36, 70]
	Gender (female)	50 (62%)
	BMI (kg/m <sup>2</sup> )	26 [23, 28]
	ASA score	
	<i>I/II</i>	76 (95%)
	<i>III/IV</i>	4 (5%)
Diagnostic	Duration of symptoms before surgery (months)	13 [4–26]
	Glycopenic symptoms	74 (92%)
	<i>Coma</i>	10 (12%)
	<i>Faint</i>	32 (40%)
	<i>Epilepsy</i>	8 (10%)
	Adrenergic symptoms	46 (58%)
	Weight gain	40 (50%)
Surgery	Surgery after 2000	39 (49%)
	Laparoscopic approach	25 (31%)
	<b>Parenchyma-sparing pancreatectomy</b>	41 (51%)
	Type of surgery	
	<i>Enucleation</i>	37 (46%)
	<i>Pancreaticoduodenectomy</i>	9 (11%)
	<i>Distal pancreatectomy</i>	21 (26%)
	<i>Distal pancreatectomy with splenectomy</i>	9 (11%)
	<i>Caudal pancreatectomy</i>	4 (5%)
	<i>Uncinate process resection</i>	1 (1%)
	Duration of surgery (min)	210 [162, 240]
Accurate location before surgery	76 (95%)	

BMI = Body Mass Index; ASA = American Society of Anesthesiologists

**Table 2:** Capabilities of imaging tests for preoperative tumour detection

<b>Imaging tests</b>	<b>Number of patients (%)</b>	<b>Number of patients with a correctly localised lesion</b>	<b>True localisation rate (%)</b>	<b>95% Confidence interval</b>
<b>CT</b>	76 (96%)	58	76%	64–85
<b>MRI</b>	37 (47%)	28	76%	58–88
<b>EUS</b>	71 (89%)	64	90%	80–96
<b>Somatostatin receptor imaging</b>	26 (34%)	10	38%	21–59
<b>CT and MRI</b>	35 (44%)	32	91%	76–98
<b>CT and EUS</b>	68 (85%)	65	96%	87–99
<b>CT and IRM and EUS and somatostatin receptor imaging</b>	19 (34%)	19	100%	79–100

CT = computed tomography; MRI = magnetic resonance imaging; EUS = Endoscopic ultrasonography

**Table 3:** Outcome after pancreatic surgery for insulinoma in 80 patients

<b>Outcomes</b>	<b>Total (n = 80)</b>	<b>Parenchyma- sparing pancreatectomy (n = 41)</b>	<b>Formal resection (n = 39)</b>	<b>p</b>
<b>90-day mortality</b>	5 (6%)	1 (2.5%)	4 (10%)	0.95
<b>90-day morbidity</b>	58 (72%)	32 (78%)	26 (67%)	0.37
<i>Grade III, IV, V complications</i>	24 (30%)	13 (32%)	11 (28%)	0.92
<b>Pancreatic fistula</b>	35 (44%)	20 (49%)	15 (38%)	0.48
<b>Grade B/C</b>	19(24%)	13 (32%)	6 (15%)	0.15

**Table 4:** Outcome after pancreatic surgery for insulinoma in 80 patients

	Variables	Total (n=80)	Before 2000 (n=41)	After 2000 (n=39)	p	
Preoperative	Age	50 [36, 70]	50 [33, 62]	59 [36, 72]	0.28	
	Male	30 (38%)	16 (39%)	14 (36%)	0.95	
	ASA score	<i>I and II</i>	76 (95%)	40 (98%)	36 (92%)	
		<i>III and IV</i>	4 (5%)	1 (2%)	3 (8%)	0.57
	Neuroglycopenic symptoms	46 (58%)	17 (41%)	29 (74%)	<b>0.006</b>	
	CT	76 (96%)	38 (95%)	38 (97%)	1	
	MRI	37 (47%)	10 (24%)	27 (71%)	<b>&lt;0.001</b>	
	Somatostatin receptor imaging	26 (32%)	9 (22%)	17 (44%)	0.07	
	EUS	71 (89%)	36 (88%)	35 (90%)	1	
	<b>Non-located tumour</b>	<b>4 (5%)</b>	<b>3 (7%)</b>	<b>1 (3%)</b>	<b>0.64</b>	
Intraoperative	<b>Parenchyma-sparing pancreatectomy</b>	41 (51%)	21 (51%)	20 (51%)	1.00	
	<b>Laparoscopy</b>	25 (31%)	9 (22%)	16 (41%)	0.11	
	Type of surgery	<i>Enucleation</i>	36 (45%)	21 (51%)	<b>15 (38%)</b>	0.14
		<i>Pancreaticoduodenectomy</i>	9 (11%)	3 (7%)	6 (15%)	
		<i>Distal pancreatectomy</i>	<b>21 (26%)</b>	<b>14 (34%)</b>	<b>7 (18%)</b>	
		<i>Distal pancreatectomy with splenectomy</i>	9 (11%)	2 (5%)	7 (18%)	
		<i>Caudal pancreatectomy</i>	<b>4 (5%)</b>	<b>1 (2%)</b>	<b>3 (8%)</b>	
		<i>Uncinate process resection</i>	<b>1 (1%)</b>	.	<b>1 (3%)</b>	
<b>Drainage</b>	74 (92%)	41 (100%)	33 (85%)	<b>0.03</b>		
Postoperative	<b>Postoperative death</b>	5 (6%)	4 (10%)	1 (3%)	0.39	
	<b>Overall morbidity</b>	58 (72%)	24 (59%)	34 (87%)	<b>0.009</b>	
	<b>Severe complications</b>	24 (30%)	12 (29%)	12 (31%)	1	
	<b>Pancreatic fistula</b>	35 (44%)	11 (27%)	24 (62%)	<b>0.004</b>	
	<b>Severe fistula</b>	19 (24%)	7 (17%)	12 (31%)	0.24	

CT = computed tomography; ASA = American Society of Anesthesiologists; MRI = Magnetic resonance imaging; EUS = Endoscopic Ultrasound