



**HAL**  
open science

## Impact of missing data on mixtures and clustering with illustrations in Biology and Medicine

Christophe Biernacki, Claire Boyer, Gilles Celeux, Julie Josse, Fabien Laporte, Matthieu Marbac, Aude Sportisse, Vincent Vandewalle

### ► To cite this version:

Christophe Biernacki, Claire Boyer, Gilles Celeux, Julie Josse, Fabien Laporte, et al.. Impact of missing data on mixtures and clustering with illustrations in Biology and Medicine. ASA 2022 - Apprentissage Statistiques et Applications, Jun 2022, Poitiers, France. hal-03936781

**HAL Id: hal-03936781**

<https://inria.hal.science/hal-03936781v1>

Submitted on 12 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Impact of missing data on mixtures and clustering with illustrations in Biology and Medicine

C. Biernacki

with C. Boyer, G. Celeux, J. Josse, F. Laporte, M. Marbac, A. Sportisse, V. Vandewalle

June 23rd 2022, Poitiers

## Take home message

Missing data may change preconceptions

- **Mixtures**: EM has unexpected behaviour concerning degeneracy dynamic
- **Clustering**: the missing data pattern may convey some information on partition

These topics are in the research agenda of Statisticians since:

The larger the datasets, the more missing data may appear. . .

○○○○○  
○○○○○○○○○○  
○○○○○○○○○○○○  
○○○

# Outline

## 1 Introduction

## 2 Impact of MAR data on the EM algorithm

- Gaussian mixture degeneracy *without* missing data
- Gaussian mixture degeneracy *with* missing data

## 3 Impact of MAR data on clustering

- A model-based MAR approach
- Biology study illustration

## 4 Impact of MNAR data on clustering

- A model-based MNAR clustering approach
- Medical study illustration

## 5 Concluding remarks

○○○○○  
○○○○○○○○○○  
○○○○○○○○○○○○  
○○○

## Missing data: notations

- $y = \{y_1 | \dots | y_n\}^T$ : **full dataset** with  $n$  individuals
- $y_i = (y_{i1}, \dots, y_{id}) \in \mathcal{Y} = \mathbb{R}^d$  with individual  $i \in \{1, \dots, n\}$
- $c = \{c_1 | \dots | c_n\}^T \in \{0, 1\}^{n \times d}$ : **pattern of missing data** for the full dataset
- $c_i = (c_{i1}, \dots, c_{id}) \in \{0, 1\}^d$ : pattern of missing data for individual  $i \in \{1, \dots, n\}$

$$c_{ij} = 1 \Leftrightarrow y_{ij} \text{ is missing}$$

- $y_i^{\text{obs}}$ : the observed variables values for indiv.  $i$  (and  $y^{\text{obs}} = \{y_1^{\text{obs}} | \dots | y_n^{\text{obs}}\}^T$ )
- $y_i^{\text{mis}}$ : the missing variables values for individual  $i$  (and  $y^{\text{mis}} = \{y_1^{\text{mis}} | \dots | y_n^{\text{mis}}\}^T$ )

○○○○○  
○○○○○○○○○○○  
○○○○○○○○○○○○  
○○○

## Typology of the missingness mechanisms

- Missing completely at random (**MCAR**):

$$p(c|y; \psi) = p(c; \psi) \quad \forall y$$

- Missing at random (**MAR**):

$$p(c|y; \psi) = p(c|y^{\text{obs}}; \psi) \quad \forall y^{\text{mis}}$$

- Missing not at random (**MNAR**): the mechanism is not MCAR nor MAR

## Ignorable vs. non ignorable model

A missing mechanism is ignorable if likelihoods can be decomposed as

$$L(\theta, \psi; \underbrace{y^{\text{obs}}, c}_{\text{observed data}}) = L(\psi; c | y^{\text{obs}}) \times L(\theta; y^{\text{obs}})$$

Some simple algebra show that this occurs when missing mechanism is not MNAR

### Inference of $\theta$

“If the missing mechanism is **ignorable** then likelihood-based inferences for  $\theta$  from  $L(\theta; y^{\text{obs}})$  will be the same as likelihood based inference for  $\theta$  from  $L(\theta, \psi; y^{\text{obs}}, c)$ .”

[Little and Rubin, 2002 Section 6.2]

- M(C)AR is ignorable
- MNAR is not ignorable

○○○○  
○○○○○

○○○○  
○○○

○○○○○○○  
○○○

## Mixture model and clustering (Gaussian case)

- **Partition ( $K$  clusters):**  $z = (z_1 | \dots | z_n)^T \in \{0, 1\}^{n \times K}$  where
  - $z_i = (z_{i1}, \dots, z_{iK}) \in \{0, 1\}^K$
  - $z_{ik} = 1$  if  $y_i$  belongs to cluster  $k$ ,  $z_{ik} = 0$  otherwise
- **Mixture model:**  $y_1, \dots, y_n$  are i.i.d. from the  $d$ -variate Gaussian mixture

$$f(y_i; \theta) = \sum_{k=1}^K \pi_k f_k(y_i; \alpha_k)$$

- $\pi_k = \mathbb{P}(z_{ik} = 1)$
  - $f_k(\cdot; \alpha_k) = \phi(\cdot; \mu_k, \Sigma_k)$  is the  $d$ -variate **Gaussian distribution** with mean vector  $\mu_k$  and covariance matrix  $\Sigma_k$
  - $\theta = (\pi_1, \dots, \pi_K, \alpha_1, \dots, \alpha_K)$  is the whole mixture parameter
- **Clustering:** MAP principle from the mixture output to estimate the partition

**And also:** algorithm estimation ( EM, SEM... ) and model selection ( BIC, ICL... )



# Outline

## 1 Introduction

## 2 Impact of MAR data on the EM algorithm

- Gaussian mixture degeneracy *without* missing data
- Gaussian mixture degeneracy *with* missing data

## 3 Impact of MAR data on clustering

- A model-based MAR approach
- Biology study illustration

## 4 Impact of MNAR data on clustering

- A model-based MNAR clustering approach
- Medical study illustration

## 5 Concluding remarks



## Outline

- 2 Impact of MAR data on the EM algorithm
  - Gaussian mixture degeneracy *without* missing data
  - Gaussian mixture degeneracy *with* missing data



## Degeneracy genesis: unbounded likelihood

- $d$ -variate Gaussian mixture**

$$f(y_i; \theta) = \sum_{k=1}^K \pi_k \underbrace{\frac{1}{(2\pi)^{d/2} |\Sigma_k|^{1/2}} \exp\left(-\frac{1}{2}(y_i - \mu_k)^T \Sigma_k^{-1} (y_i - \mu_k)\right)}_{\phi(y_i; \mu_k, \Sigma_k)}$$


- Sampling:**  $y_1, \dots, y_n \stackrel{i.i.d.}{\sim} p(\cdot; \theta)$  without any missing data ( $y^{\text{obs}} = y$ )
- Likelihood:**  $\ell(\theta; y) = \ln L(\theta; y) = \sum_{i=1}^n \ln f(y_i; \theta)$

$$\text{particular center } \mu_2 = y_i \quad \Rightarrow \quad \lim_{|\Sigma_2| \rightarrow 0} \ell(\theta; y) = +\infty$$

[Kiefer and Wolfowitz, 1956] [Day, 1969]




## EM behaviour: illustration



- degeneracy may occur even when starting from large variances
- convergence can be slow when far from the degenerate limit
- convergence extremely fast near degeneracy

## EM behaviour: results



$$u_0 = \left[ \frac{1}{p_{i_0 k_0}}, \{p_{ik_0}\}_{i \neq i_0} \right]$$

degeneracy of component  $k_0$  at  $y_{i_0}$

$\Leftrightarrow$

$$\|u_0\| \rightarrow 0$$

[Biernacki and Chrétien, 2003]

[Ingrassia and Rocci, 2009]

### Proposition 1: Existence of a basin of attraction

$\exists \epsilon > 0$  s.t. if  $\|u_0\| \leq \epsilon$  then  $\|u_0^+\| = o(\|u_0\|)$  with probability 1.

### Proposition 2: Speed towards degeneracy is exponential

$\exists \epsilon > 0, \alpha > 0$  and  $\beta > 0$  s.t. if  $\|u_0\| \leq \epsilon$  then, with probability 1,

$$|\Sigma_{k_0}^+| \leq \alpha / |\Sigma_{k_0}| \cdot \exp(-\beta / |\Sigma_{k_0}|).$$

# Outline


- 2 Impact of MAR data on the EM algorithm
  - Gaussian mixture degeneracy *without* missing data
  - Gaussian mixture degeneracy *with* missing data




## EM behaviour illustration

- Breast cancer tissue of the UCI database repository: 106 units, 9 variables.
- 10% of missing data randomly generated
- $K = 4$  clusters

Log-likelihood according to the number of iterations



Decrease of the log-determinant of the degenerated component





## Detail from the illustration

	1	2	3	4	5	6	7	8	9
1	211.00		0.09	30.75	151.98	4.94	14.27	27.24	217.13
2	196.86	0.02	0.09	28.59	82.06	2.87	7.97	27.66	200.75
3	144.00	0.12	0.05	19.65	70.43	3.58		7.57	160.37
4	172.52	0.13	0.04		192.22	5.12	19.32	32.19	174.93
5	121.00	0.17	0.09	24.44	144.47	5.91	22.02	10.59	141.77
6	223.00	0.12	0.08	33.10	197.01	5.95	30.45	12.96	252.48
7		0.17	0.23	34.22	94.35	2.76	31.28	13.88	180.61
8	303.00	0.06	0.04	22.57		4.54	21.83	5.72	321.65
9	250.00	0.09	0.09	29.64	180.76	6.10	26.14	13.96	280.12
10	391.00	0.06	0.01	35.78		7.41	22.13	28.11	400.99
11	176.00	0.09	0.08	20.59	79.71		18.23	9.58	191.99
12	145.00		0.11	21.22	82.46	3.89	20.30	6.17	162.51
13	124.13	0.13	0.11	20.59			18.46	9.12	134.89
14	103.00	0.16	0.29	23.75	78.26	3.29	22.32	8.12	124.98

Table : Data belonging to the degenerated component.

- Cvg. towards a degenerated component (no plateau of the log-likelihood)
- Degeneracy relatively slow: log-likelihood linear according to the nb of it.
- Number of points of the degenerated solution greater than the space dimension  $d$  (but the number of complete points lower than  $d$ )





## Intermediate conclusion on missing data

### Like the complete data $y$ case

- Likelihood is unbounded
- EM can be attracted by degenerate solutions

### Unlike the complete data $y$ case

- Risk to consider a degenerated solution as valid
- Risk of losing a lot of time in useless iterations

**Statisticians should be aware of such dangerous EM behaviour. . .**

. . . since missing data are more and more frequent



## Understanding degeneracy speed on a toy example

- Univariate framework, no mixture, only one observed data:  $y$
- Maximum likelihood estimator (**Unbounded likelihood!**):  $\mu = y, \sigma^2 = 0$
- Suppose equivalently that  $n - 1$  data are unobserved (unchanged likelihood)
- Here is one iteration of a (useless) EM algorithm (it.  $q$ )

$$\mu^{(q+1)} = \frac{(n-1)\mu^{(q)} + y}{n} \quad \text{and} \quad \sigma^{2(q+1)} = \frac{(n-1)\sigma^{2(q)} + (y - \mu^{(q+1)})^2}{n}$$

Linear grow of the log-likelihood (have a look also when  $n$  increases!)

$$\ell(\theta^{(q)}; x) \sim -0.5q \log \frac{n-1}{n}$$

Geometrical convergence rate towards 0 for the variance

$$\sigma^{2(q)} \sim \sigma^{2(0)} \left( \frac{n-1}{n} \right)^q$$



## Influence of the missing data rate

% missing data	0	5	10	15	20	25	30
% deg.	16	4	12	11	46	51	100
Average nb of iterations before deg.	2	13	13	82	304	138	215

**Table :** Frequency and speed of degeneracy (deg.) according to the rate of missing data on the breast cancer data set.

### When the rate of missing data increases:

- The rate of degeneracy increases
- The number of iterations before degeneracy seems to (globally) increase

**Again, statisticians should be aware of such dangerous EM behaviour...**

... since missing data are more and more frequent

○○○○○  
○○○○○○○○○○  
○○○○○○○○○○○○  
○○○

# Outline

## 1 Introduction

## 2 Impact of MAR data on the EM algorithm

- Gaussian mixture degeneracy *without* missing data
- Gaussian mixture degeneracy *with* missing data

## 3 Impact of MAR data on clustering

- A model-based MAR approach
- Biology study illustration

## 4 Impact of MNAR data on clustering

- A model-based MNAR clustering approach
- Medical study illustration

## 5 Concluding remarks

# Outline

- 3 **Impact of MAR data on clustering**
  - **A model-based MAR approach**
  - **Biology study illustration**

## Full mixed data: conditional independence everywhere<sup>1</sup>

The aim is to combine continuous, categorical, integer data, ordinal, ranking and functional data

$$y_1 = (y_1^{cont}, y_1^{cat}, y_1^{int}, \dots)$$

The proposed solution is to mixed all types by **inter-type conditional independence**

$$p(y_1; \alpha_k) = p(y_1^{cont}; \alpha_k^{cont}) \times p(y_1^{cat}; \alpha_k^{cat}) \times p(y_1^{int}; \alpha_k^{int}) \times \dots$$

In addition, for symmetry between types, **intra-type conditional independence**

Only need to define the univariate pdf for each variable type!

- **Continuous**: Gaussian
- **Categorical**: multinomial
- **Integer**: Poisson
- ...

<sup>1</sup>MixtComp package: <https://cran.r-project.org/web/packages/RMixtComp/index.html>

## Missing data: MAR assumption and estimation

### Assumption on the missingness mechanism

**Missing At Random (MAR):** the probability that a variable is missing does not depend on its own value given the observed variables.

### Observed log-likelihood...

$$\ell(\theta; y^{\text{obs}}) = \sum_{i=1}^n \log \left( \sum_{k=1}^K \pi_k p(y_i^{\text{obs}}; \alpha_k) \right) = \sum_{i=1}^n \ln \left[ \sum_{k=1}^K \pi_k \underbrace{\int_{y_i^{\text{mis}}} p(y_i^{\text{obs}}, y_i^{\text{mis}}; \alpha_k) dy_i^{\text{mis}}}_{\text{MAR assumption}} \right]$$



## SEM algorithm

A SEM algorithm to estimate  $\theta$  by maximizing the **observed**-data log-likelihood

- Initialisation:  $\theta^{(0)}$
- Iteration nb  $q$ :
  - **E-step**: compute conditional probabilities  $p(y^{\text{mis}}, z | y^{\text{obs}}; \theta^{(q)})$
  - **S-step**: draw  $(y^{\text{mis}(q)}, z^{(q)})$  from  $p(y^{\text{mis}}, z | y^{\text{obs}}; \theta^{(q)})$
  - **M-step**: maximize  $\theta^{(q+1)} = \arg \max_{\theta} \ln p(y^{\text{obs}}, y^{\text{mis}(q)}, z^{(q)}; \theta)$
- Stopping rule: iteration number

**Properties: simpler than EM and interesting properties!**

- Avoid possibly difficult E-step in an EM
- Classical M steps
- Avoids local maxima
- The mean of the sequence  $(\theta^{(q)})$  approximates  $\hat{\theta}$
- The variance of the sequence  $(\theta^{(q)})$  gives confidence intervals



# Outline

- 
- 
- 3 Impact of MAR data on clustering**
  - A model-based MAR approach
  - Biology study illustration



## Prostate cancer data (with missing data)<sup>3</sup>

- **Individuals:** 506 patients with prostatic cancer grouped on clinical criteria into two Stages 3 and 4 of the disease
- **Variables:**  $d = 12$  pre-trial variates were measured on each patient, composed by **eight continuous** variables (age, weight, systolic blood pressure, diastolic blood pressure, serum haemoglobin, size of primary tumour, index of tumour stage and histologic grade, serum prostatic acid phosphatase) and **four categorical** variables with various numbers of levels (performance rating, cardiovascular disease history, electrocardiogram code, bone metastases)
- Some **missing data:** 62 missing values ( $\approx 1\%$ )

We forget the classes (Stages of the disease) for performing **clustering**

### Questions

- How many clusters?
- Which partition?

<sup>3</sup>Byar DP, Green SB (1980): Bulletin Cancer, Paris 67:477-488

○○○○○  
○○○○○○○

○○○○  
○○●○

○○○○○○○○  
○○○

## Two strategies in competition

- **Strategy “mice<sup>4</sup> + MixtComp”**: MixtComp on the dataset completed by mice

```
> data.imp=mice(data)
> data.comp.mice=complete(data.imp)
```

- **Strategy “full MixtComp”**: MixtComp on the observed (no completed) dataset


### Partition quality with $K = 2$

Strategy	mice + MixtComp	full MixtComp
% misclassified	12.8	<b>8.1</b>


<sup>4</sup><http://cran.r-project.org/web/packages/mice/mice.pdf>



## Choosing $K$ with the ICL criterion



mice + MixtComp  
 $\hat{K} = 7$



full MixtComp  
 $\hat{K} = 2$

... may lose some cluster information when imputation before clustering

# Outline

- 1 Introduction
- 2 Impact of MAR data on the EM algorithm
  - Gaussian mixture degeneracy *without* missing data
  - Gaussian mixture degeneracy *with* missing data
- 3 Impact of MAR data on clustering
  - A model-based MAR approach
  - Biology study illustration
- 4 Impact of MNAR data on clustering**
  - A model-based MNAR clustering approach**
  - Medical study illustration**
- 5 Concluding remarks



## Outline

- 4 Impact of MNAR data on clustering
  - A model-based MNAR clustering approach
  - Medical study illustration

## Proposed MNAR models in clustering

### Question we address now

Since MNAR is not ignorable, which distribution  $p(c|y, z; \psi)$  to propose?

### Hypothesis 1: conditional independence

$$p(c_i | y_i, z_{ik} = 1; \psi) = \prod_{j=1}^d p(c_{ij} | y_i, z_{ik} = 1; \psi)$$

### Hypothesis 2: linear function within canonical link functions $\rho$

$$p(c_{ij} = 1 | y_i, z_{ik} = 1; \psi) = \rho(\alpha_{kj} + \beta_{kj} y_{ij})$$

- $\psi = (\alpha, \beta)$  where  $\alpha = (\alpha_{11}, \dots, \alpha_{1d}, \dots, \alpha_{K1}, \dots, \alpha_{Kd})^T \in \mathbb{R}^{Kd}$  and  $\beta = (\beta_{11}, \dots, \beta_{1d}, \dots, \beta_{K1}, \dots, \beta_{Kd})^T \in \mathbb{R}^{Kd}$
- $\rho$  is the cdf of any continuous distribution (logit, probit)



## A by-product zoology of MNAR models

	Effect on the variable $j$		Effect on the class membership $k$		Nb parameters
	Depends on $j$	Depends on $k$	Depends on $j$	Depends on $k$	Continuous
MNAR $z^j y^k$	✓	✓	✓	✓	$2Kd$
MNAR $yz^j$	✓	✗	✓	✓	$(K + 1)d$
MNAR $y^k z$	✓	✓	✗	✓	$K(d + 1)$
MNAR $yz$	✓	✗	✗	✓	$(K + d)$
MNAR $y$	✓	✗	✗	✗	$d$
MNAR $y^k$	✓	✓	✗	✗	$Kd$
MNAR $z$	✗	✗	✗	✓	$K$
MNAR $z^j$	✗	✗	✓	✓	$Kd$

Remarks:


- MNAR $z^j y^k$  is the most complex model
- MNAR $z$ , MNAR $z^j$ : the only effect of missingness is on the class membership,  $\psi = (\alpha_{11}, \dots, \alpha_{1d}, \dots, \alpha_{K1}, \dots, \alpha_{Kd})^T$ ,  $p(c_{ij} = 1 \mid y_i, z_{ik} = 1; \psi) = \rho(\alpha_{kj})$
- MCAR is a specific and simple case



## MNAR<sub>z</sub> analysis: pattern $c$ gives information on partition $z$ !

Draw Bayes error of a MNAR<sub>z</sub> model with two components and 20% of missing data

$$\pi_k = 0.5, \|\mu_2 - \mu_1\| \text{ varies}, \Sigma_1 = \Sigma_2 = \mathbf{I}, |\alpha_2 - \alpha_1| \text{ varies}$$



Both  $\mu_k$  and  $\alpha_k$  act on the Bayes error



## Reinterpretation of the MNAR<sub>z</sub> and MNAR<sub>z</sub><sup>j</sup> models as MAR

Commonly used in Machine Learning [Jones, 1996], [Little and Rubin, 2002], [Josse *et al.*, 2019]

Mixture model for  $y^{\text{obs}}$  and Bernoulli distribution for  $C$   
 $\Leftrightarrow$  MAR mixture model for  $\tilde{y}^{\text{obs}} = (y^{\text{obs}}|c)$

For example,

$$y^{\text{obs}} = \begin{pmatrix} ? & 2.6 & 5 \\ \text{blue} & 1.9 & 4 \\ \text{red} & 2.3 & ? \end{pmatrix}, \quad c = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

then  $\tilde{y}^{\text{obs}}$  is expressed as

$$\tilde{y}^{\text{obs}} = \begin{pmatrix} ? & 2.6 & 5 & 1 & 0 & 0 \\ \text{blue} & 1.9 & 4 & 0 & 0 & 0 \\ \text{red} & 2.3 & ? & 0 & 0 & 1 \end{pmatrix}.$$

### Proposition 3: in terms of maximum likelihood

The maximum likelihood estimate associated to the dataset  $\tilde{y}^{\text{obs}}$  under MAR model is the one associated to the dataset  $y^{\text{obs}}$  under MNAR<sub>z</sub> or MNAR<sub>z</sub><sup>j</sup> models.

## Identifiability

Previous works: [Teicher, 1963] (without NA), [Miao et al., 2016] (for MNAR data)

### Proposition 4

Assume that

- 1 The marginal mixture  $\sum_{k=1}^K \pi_k f_k(y_i; \theta_k)$  is identifiable
- 2 There exists a total ordering  $\preceq$  of  $\mathcal{F}_j \times \mathcal{R}$ , for  $j \in \{1, \dots, d\}$  fixed, where  $\mathcal{F}_j = \{f_{1j}, \dots, f_{Kj}\}$  and  $\mathcal{R} = \{\rho_1, \dots, \rho_K\} = \{\rho(\cdot; \psi_1), \dots, \rho(\cdot; \psi_K)\}$ . The total ordering is s.t.  $\forall k < \ell, F_k = \rho_k f_{kj} \preceq F_\ell = \rho_\ell f_{\ell j}$  implies

$$\lim_{u \rightarrow +\infty} \frac{\rho_\ell(u) f_{\ell j}(u)}{\rho_k(u) f_{kj}(u)} = 0$$

Then the mixture model with one of the MNAR\* mechanisms is identifiable up to label swapping

All MNAR models are identifiable (or at least generically identifiable) for probit/logit



## Estimation procedure overview

- Use EM or Stochastic EM (SEM) algorithms
- MNAR<sub>z</sub> and MNAR<sub>z</sub><sup>j</sup>: EM and SEM are very simple
- MNAR<sub>y</sub>\*: the SE step requires a within Gibbs loop, sometimes involving itself a Sampling Importance Resampling (SIR)

	EM		SEM	
MNAR <sub>z</sub> MNAR <sub>z</sub> <sup>j</sup>	✓		✓	
	Probit	Logit	Probit	Logit
MNAR <sub>y</sub> *	no closed form	no closed form, optim. pb	✓	require algorithms as SIR (costly)



## Model selection

Can select between MCAR and MNAR\* with any information criterion (BIC, ICL)

Even if the missing mechanism is ignorable for MCAR. . .

. . . need to model  $c$  to compare a MCAR and a MNAR model

### CAUTION

- It is just a selection between several proposed MNAR models
- It is not deciding if missingness procedure is “genererically” MNAR or not



## Outline

- 4 Impact of MNAR data on clustering
  - A model-based MNAR clustering approach
  - Medical study illustration




## Hospital data description

- Number of patients:  $n = 5\,146$
- Number of features:  $d = 7$ 
  - Age
  - Size
  - Weight
  - Cardiac frequency
  - Hemoglobin concentration
  - Temperature
  - Minimum Diastolic and Systolic Blood Pressure
- Percentage of missing data: 6.4%

Doctors are convinced that their missing data are MNAR



## ICL comparison



- MCAR, MNARy and MNARz are equivalent until  $K = 3$
- MNARz and MNARyz clearly indicate presence of an additional cluster ( $K = 4$ )

It seems to be an illustration of the effect of  $c$  through MNARz and MNARyz



# Outline

## 1 Introduction

## 2 Impact of MAR data on the EM algorithm

- Gaussian mixture degeneracy *without* missing data
- Gaussian mixture degeneracy *with* missing data

## 3 Impact of MAR data on clustering

- A model-based MAR approach
- Biology study illustration

## 4 Impact of MNAR data on clustering

- A model-based MNAR clustering approach
- Medical study illustration

## 5 Concluding remarks



## To conclude

### Summary

- Statisticians should properly consider the potential missing data impact both from an **algorithmic** and from a **modeling** point of view
- EM: be careful about degeneracy which seems to exacerbated/masqued
- MAR: the minimal approach to be used by statisticians. . .
- MNAR: interest of the simple but meaningful model MNARz, link with usual methods

### Ongoing works

- EM: propose mechanism to identify/discard degenerate runs
- MNAR: extend to categorical, count and mixed data

Thanks!