



HAL
open science

Mathematical modelling of cancer growth and drug treatments: taking into account cell population heterogeneity and plasticity

Jean Clairambault

► **To cite this version:**

Jean Clairambault. Mathematical modelling of cancer growth and drug treatments: taking into account cell population heterogeneity and plasticity. 2022. hal-03938411v1

HAL Id: hal-03938411

<https://inria.hal.science/hal-03938411v1>

Preprint submitted on 13 Jan 2023 (v1), last revised 5 Apr 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mathematical modelling of cancer growth and drug treatments: taking into account cell population heterogeneity and plasticity

Jean Clairambault*

Abstract—Mathematical models of cancer growth and evolution of cancer cell characteristics, aka phenotypes, together with optimisation and optimal control methods to contain them, in the framework of adaptive dynamics of cell populations, are presented, that take into account the *heterogeneity* of cancer cell populations, i.e., their biological variability, and their intrinsic *plasticity*, i.e., their nongenetic instability that allows them to quickly adapt to changing environments. The presented vision of the cancer disease, which is specific to multicellular organisms, relies on a relatively novel vision, consistent with a billion-year evolutionary perspective. Based on recent contributions from *philosophy of cancer*, these mathematical models aim at designing theoretical therapeutic strategies to simultaneously contain tumour progression and limit adverse events of drugs to healthy cell populations.

I. INTRODUCTION

“What is cancer?” is a question that must somehow be answered with clear ideas before proposing relevant mathematical models of cancer growth and phenotype evolution, together with methods to optimise its treatments. Recent contributions from philosophy of biology [4], immunology, and archeology of genomes, are reported here, stressing the importance of a default of cohesion within and between tissues of a same organism - as cancer is a disease of multicellular organisms only -, that manifests itself as an impairment of organismic control on both proliferation and differentiation.

Different mathematical models of tumour growth involving cancer cell population heterogeneity and plasticity, and drug treatments of cancer are presented here. They are all deterministic models, with a focus set on continuous phenotype-structured equations and adaptive dynamics of cell populations, healthy and cancer, and the effects on them of anticancer drugs. Constraints linked to avoiding the main pitfalls of therapeutics in oncology, namely therapy resistance in cancer cells and adverse toxic events in healthy cells, together with an objective function that is most often limited to maximising tumour cell kill, are considered, paving the way for the use of optimisation methods. Optimised clinical strategies for the continuous delivery of anticancer drugs may thus be designed by resorting to optimal control methods. Such optimised strategies should thwart some known pathological strategies used by cancer cell populations to dodge deadly insults induced by drugs, such as the development of drug-resistant cell phenotypes and the so-called ‘bet hedging’

by which tumours may simultaneously face different modes of insults or changing environments [5], [23], [44].

From a modelling point of view, the complete reversibility of the cell phenotypes under study when the environment changes, as represented by the reversible behaviour of the solutions of the equations in terms of cell-structuring traits (aka phenotypes), is consistent with the fact that we study in an isogenic context cell differentiations and de-differentiations, that rely on *epimutations*, i.e., epigenetic changes in the expression of genes, and not on actual mutations [24]. With or without interactions between cell populations, the equations are of the nonlocal Lotka-Volterra type, structured by adaptive cell phenotypes, with optional diffusion and advection terms. Adaptation is thus considered at a cell population status at which the situation of a cancer cell population is reversible (i.e., not after mutations), in particular for the occurrence of drug-induced drug resistance, which allows to theoretically justify the clinical notions of *drug holiday* and *re-challenge* in cancer [54].

II. WHERE BIOLOGY MEETS PHILOSOPHY

A. Heterogeneity and plasticity of cancer cell populations

Heterogeneity of cancer cell populations [35] has often been mentioned as a major obstacle to the efficacy of drug treatments, in particular of so-called ‘targeted therapies’ [22], as treatments that may be very effective on a dominant genetic clone of a tumour cease to be so when another clone emerges [15], be it in malignant haemopathies [15] or in solid tumours [21]. This certainly concerns genetic heterogeneity, i.e., situations in which spontaneous mutations have appeared due to the genetic instability of tumour cells, but isogenic heterogeneity resulting from nongenetic instabilities in gene expression due to the lability of epigenetic enzyme activity [51] is as much important [19], [20], [36], and it occurs before mutations, that remain rare events. The plasticity of cancer cells with respect to the expression of genes [12], [19], [36], [58], which, as all differentiations that make an isogenic multicellular organism so varied in cell types (20 for sponges [39], [40], about 200 for human beings), is due to epigenetic factors, strictly controlled at the organismic level in physiology, and out of control in cancer [19], [20]. It might be measured by the velocity of adaptation of cell phenotypes, such as fecundity, viability or motility, to changes in the cell environment.

B. Philosophy of cancer

A recently autonomised field of knowledge, the so-called *philosophy of cancer* [4], [12], [53] considers cancer as

*Jean Clairambault is with Laboratoire Jacques-Louis Lions, Sorbonne Université and Inria Paris, 4, place Jussieu, 75252 Paris cedex 05, France jean.clairambault@inria.fr

an impairment of organismic control on both proliferation and differentiations [4]. It should be stressed that cancer is a disease that can be met only in the animal kingdom, constituted of multicellular organisms, that, different from plants (plants may also develop tumours, but they remain localised in capsules, never threatening the existence of the organism), are endowed with other properties such as motility and heterotrophicity. Animals, also called Metazoa, emerge from a zygote (fecundated egg) and are thus completely isogenic. The same genome thus exists in all animal cells, and the extreme diversity of cell types that constitute an animal organism is only due to successive differentiations, by selective expression or repression of genes at each differentiation step. Such expressions or repressions are under strict epigenetic control, i.e., under the control of epigenetic enzymes that may be impaired in cancer. Furthermore, in cancer, these epigenetic mechanisms are easily manipulated in cancer cell populations in response to changes of the tumour microenvironment (such as due to the introduction of life-threatening drugs) to develop (reversible) resistant cell phenotypes. The question of understanding what makes a multicellular individual coherent, contrary to a tumour, may not always be summed up by its zygotic origin, and it is the object of the *philosophy of immunology* [48], [49], as one of the components that define a multicellular individual is the necessary existence of an immune system [39], [40], absent in tumours, which makes possible the success of immunotherapies.

C. The atavistic theory of cancer

In a completely independent way from the above mentioned works, the atavistic theory of cancer was speculated and popularised from 2011 on in a series of papers [13], [56], [30], relying or partly confirmed in previous [25], or following papers examining phylostratigraphic trees [17], [18], [31] or archeogenetic data [6], [10], [55]. It proposes that cancer is a reversal in evolution from a coherent multicellular organism towards an incoherent, nonviable state closer to unicellularity, but representing an autonomous, coarse and unsuccessful try towards multicellularity from partly organised collections of cells that escape organismic control, a state which these authors name Metazoa 1.0. The interest of considering this atavistic theory is that all cancers are constituted of cells in a same multicellular organism, that have kept in the memory of their common genome defence mechanisms against external aggressions from the environment, that can be unleashed and recruited due to the absence of organismic control, that otherwise would thwart them as introducing too loose cohesion of the ensemble, challenging the very existence of the organism [1]. Note however that such mechanisms, based on cell de-differentiations and re-differentiations, are at work, strictly controlled, in development and in adult physiological wound healing, some animals like the axolotl (a modern amphibian) being even able to safely and completely rebuild a cut limb.

III. TUMOUR GROWTH AND ITS CONTROL

A. Pitfalls in clinical oncology

In the clinic of cancers, drug treatments meet two main pitfalls: emergence of resistance in cancer cell populations and existence of adverse toxic effects in healthy cell populations. Although most mathematical optimisation works consider one or the other pitfall, it will be shown in the sequel that it is possible to manage them both simultaneously, and not only by following absolute limitations to both population densities, supposed to be taken from clinical considerations, but rather by following with time their respective parts and permanently controlling them.

B. Continuous models of tumour growth

Most models of tumour growth presented until recently were based on ODEs or PDEs structured in Cartesian space, not many of them are structured according to phenotypes [33], even fewer according to both phenotypes and space variables [34]. The privileged focus of this paper is on structuring tumour growth models in phenotypes [11].

C. Theoretical therapeutic control

Many models of control, including optimal control methods, mimicking therapeutics in the clinic have been proposed, for ODE or PDE models, and useful references on this topic are in particular [26], [29], [50]. A particular aspect is control in dormancy [16]. In the present article are presented examples of optimal control for phenotype-structured integro-differential or PDE models, usually of the nonlocal Lotka-Volterra type [2], [28], [47].

IV. MODELS OF ADAPTIVE DYNAMICS

A. Structured population models and adaptive dynamics

The general idea of structured models is to describe heterogeneity in a population of individuals represented by continuous densities by using continuous traits that are relevant to the question at stake, and that are susceptible to evolve with time by adaptation of such traits to changing environmental conditions or interactions with other populations of individuals. This way of representation originally comes mostly from the seminal paper of McKendrick [37] on public health problems (see in the original paper the equation p. 122) and from theoretical ecology [14], however it has been adapted to cell populations, see, e.g., [43], and a number of following papers. The structuring trait in the population of individuals may be age, size, or physiological traits determining adaptation to the environment such as fecundity, viability [7], [8], [32], motility or plasticity, as will be shown in the sequel. Indeed, as in general evolution, adaptation to stress induced by the environment [38], [41], [57] is a major determinant of trait differentiation.

B. Tumours and their micro-environment

Populations of cancer cells, in malignant haemopathies or in solid tumours, are amenable to such representations. Their micro-environment may be constituted of other cell

populations, in mutualistic [42] or competitive (in particular tumour-immune, for which many models have been published) interactions. It may also consist of mechanical or chemical constraints, as in particular by the introduction of anticancer drugs, that target different aspects of tumour growth, in particular cell death, intrinsic proliferation or action on the structuring traits. A trait of expression of resistance to a given drug has often been considered [7], [8], [33], [34], [45], [46].

C. Tumour strategies to dodge therapeutic control

As mentioned in the introduction, at least two of these strategies have been identified. In the first one [47], a phenotype of expression of drug resistance structures two cell populations, one tumoral, the other healthy, that interact only through a nonlocal logistic term. The model illustrates drug-induced drug resistance, present in cancer, not in healthy cell populations. It will be developed in the next subsection.

The other tumour strategy, that is explored in [3] (without control), is related to tumour bet hedging [5], [23], [44]. It hypothesises that in order to adapt to unpredictable changes in the tumour environment, tumour cell populations are amenable to display different cell phenotypes, with differentiated abilities to resist different threats. This is represented in the simplest possible way by phenotypic divergence, i.e., the division of the cell population into two phenotypically distinct subpopulations, each one of them maximising a a form of fitness related to two different phenotypes related by a flexible (i.e., not simply $y = 1 - x$) trade-off condition between them.

The main equation, where $z = (x, y, \theta)$ with for instance x =viability, y =fecundity, and θ =plasticity, runs as:

$$\partial_t n + \nabla \cdot \{Vn - A(\theta)\nabla n\} = (r(z) - d(z)\rho(t))n,$$

where

$$(Vn - A(\theta)\nabla n) \cdot \mathbf{n} = 0 \text{ for all } z \in \partial D$$

and

$$n(0, z) = n_0(z) \text{ for all } z \in D = \{C(x, y) \leq K\} \times [0, 1],$$

defining a trade-off between traits x and y , and the diffusion matrix

$$A(\theta) = \begin{pmatrix} a_{11}(\theta) & 0 & 0 \\ 0 & a_{22}(\theta) & 0 \\ 0 & 0 & a_{33} \end{pmatrix},$$

with a_{11} and a_{22} non decreasing functions of θ , gives the speed at which non-genetic epimutations occur, otherwise said, it is a representation of how the internal plasticity trait θ affects the non-genetic instability of traits x and y , by tuning the diffusion term $\nabla \cdot \{A(\theta)\nabla n\}$; the advection term

$$\nabla \cdot \{V(t, z)n\} = \nabla \cdot \{V_1(t, z), V_2(t, z), V_3(t, z)\}n,$$

where for instance $V(t, z) = 10^{-3}\theta(-y, -x, -(x + y))$ in simulations, represents the force of external evolutionary pressure on the population, i.e., changes in the environment;

and the integral $\rho(t) = \int_D n(t, z)dz$ stands for the total amount of individuals in the population at time t .

The reader is sent to [3] to check that phenotypic divergence around two different concentrated phenotypes (x_1, y_1) and (x_2, y_2) is shown to actually occur in the model, while the plasticity phenotype θ simultaneously decreases. In this setting where phenotype space is 3-dimensional, the plasticity trait θ tunes the Laplacian term, while the advection function represents the influence of the environment on the adaptable structure variable (3-d phenotype) $z = (x, y, \theta)$ present in the cell population with density $n(t, z)$.

D. A phenotype-structured model with optimal control

The model presented in [2] is a consistent illustration of a drug resistance phenotype-structured model in a cancer cell population with optimal control for the optimised delivery of an anticancer chemotherapy acting as an added death term in a reaction-diffusion setting. However, the somewhat richer integro-differential model presented in [47] takes simultaneously into account both emergence of drug-induced drug resistance (to be limited) and adverse toxic effects on a nonresistant healthy cell population, the two populations being linked in a mixed population term in the nonlocal logistic term of a Lotka-Volterra model, and it is described below.

The phenotype $x \in [0, 1]$ of expression of drug resistance ($x = 0$: total drug sensitivity; $x = 1$: total drug resistance) structures the two populations. However, the functions r_H, d_H, μ_H for proliferation, death and drug sensitivity, respectively, in healthy cells, are chosen much less sensitive to the drug levels and to the environment variables than their correspondent functions r_C, d_C, μ_C in cancer cells.

$$\begin{aligned} \text{(Healthy cell population H)} \quad & \frac{\partial}{\partial t} n_H(t, x) = \\ & \left[\frac{r_H(x)}{1 + k_H u_2(t)} - d_H(x)I_H(t) - u_1(t)\mu_H(x) \right] n_H(t, x) \\ \text{(Cancer cell population C)} \quad & \frac{\partial}{\partial t} n_C(t, x) = \\ & \left[\frac{r_C(x)}{1 + k_C u_2(t)} - d_C(x)I_C(t) - u_1(t)\mu_C(x) \right] n_C(t, x) \end{aligned}$$

Environment terms involving the two cell populations:

$$\begin{aligned} I_H(t) &= a_{HH} \cdot \rho_H(t) + a_{HC} \cdot \rho_C(t) \\ I_C(t) &= a_{CH} \cdot \rho_H(t) + a_{CC} \cdot \rho_C(t), \end{aligned}$$

$$\text{with } \rho_H(t) = \int_0^1 n_H(t, x) dx, \rho_C(t) = \int_0^1 n_C(t, x) dx,$$

and u_1 cytotoxic, u_2 cytostatic drug concentrations.

Optimal control problem **OCP**: find controls (u_1, u_2) minimising in fixed horizon T the objective function

$$C_T(u_1, u_2) = \rho_C(T) = \int_0^1 n_C(T, x) dx$$

under the constraints

$$\frac{\rho_H(t)}{\rho_H(t) + \rho_C(t)} \geq \theta_{HC}, \quad \rho_H(t) \geq \theta_H \cdot \rho_H(0)$$

(the last constraint, with, e.g., $\theta_H = 0.6$, to limit damage to healthy cells)

Then the following theorem may be shown:

Theorem: The optimal therapeutic trajectory (u_1, u_2) in large time $T > 0$ consists of 2 parts:

- a long-time part, with constant controls on $[0, T_1]$, at the end of which populations have almost concentrated in phenotype (for T_1 large);
- a short-time part on $[T_1, T]$ consisting of at most three arcs, for $T - T_1$ small:

- 1) a boundary arc, along the constraint $\frac{\rho_H(t)}{\rho_H(t) + \rho_C(t)} = \theta_{HC}$,
- 2) a free arc (no constraint saturating) with controls $u_1 = u_1^{\max}$ and $u_2 = u_2^{\max}$,
- 3) a boundary arc along the constraint $\rho_H(t) \geq \theta_H \cdot \rho_H(0)$ with $u_2 = u_2^{\max}$;

The reader is sent to [47] for proofs, numerical simulations and illustrations. The figure below shows the resulting optimal control (u_1, u_2) for $\theta_{HC} = 0.4$ and $\theta_H = 0.6$.




Fig. 1. Evolution with time t of the x resistance phenotype distribution curves $n_H(t, x)$ and $n_C(t, x)$ and solution (u_1, u_2) of the OCP problem for $T = 30$. In the lower two panels, one can see that the cytotoxic drug u_1 level is maintained positive, but as low as possible, close to zero, for a long time, before delivering its maximal tolerated dose on a short time interval, so as to keep the cell population as sensitive as possible until time T_1 (here $T_1 \approx 20$) - which is the *drug holiday* strategy used in the clinic of cancers [54] -, then delivered at a medium dose. The cytostatic drug u_2 level is maintained at a low level, but not close to zero, until some time before and close to T_1 , and then delivered at its maximum tolerated dose. One can see on the two rightmost panels that the cancer cell population ρ_C goes to extinction while the healthy cell population ρ_H is preserved over a prescribed level. The central panel shows an evaluation of the ratio of drug-sensitive cancer cells $\int_0^1 (1-x)n_C(t, x)dx$ over total cancer cells. Performed in modelling language AMPL with expert simulation routines IpOpt. Adapted from [47] with permission.

V. DISCUSSION ABOUT STRUCTURED MODELS

A. Including targets for control

In all mathematical models, be they phenotype-structured or not, that are designed in the perspective of therapies, physiologically based targets of control must be included in the equations. By this is mentioned the physiological nature of the control. If it primarily targets cell death (case of most classic chemotherapies), then it must be present in the equations as an added death term. If it targets proliferation, as for, e.g., tyrosine kinase inhibitors (TKIs) that are also antagonists of epidermal growth factor receptors (EGFR antagonists), then it must be present as tuning the intrinsic (that is, setting death terms to zero) instantaneous growth term $\frac{\partial n}{n \partial t}$. If it is an epigenetic drug that affects nongenetic instability, then it must address a term tuning a Laplacian $\frac{\partial^2 n}{\partial x^2}$ or other terms representing nongenetic instability in a reaction-diffusion equation. Controlling the tumour microenvironment might also be thought of as tuning an added advection term $\frac{\partial n}{\partial x}$, the structure variable x being here assumed to be one-dimensional for simplicity of presentation.

B. Taking into account heterogeneity

Various ODE models claim to take account of cell population heterogeneity by compartments representing discrete heterogeneity, such as (totally) drug-sensitive and (totally) drug-resistant cell subpopulations. This relies on a binary vision of structuring phenotypes, exchanges between compartments representing switching between discrete subpopulations, as in the *SIR* models for infectious diseases introduced by Kermack and McKendrick in 1927 [27]. The vision presented in this article is closer to the type of transport equation introduced by McKendrick (alone) in 1926 [37], to which nonlocal terms may be added, with or without diffusion and advection terms. Indeed, there is no reason why structuring traits in the population, such as fecundity, viability, motility and plasticity should be discrete. This is the reason why heterogeneity should *a priori* be represented by continuous traits. Of course, this depends on the underlying questions at stake, and on the ways of appreciating such heterogeneity. If the available measures and if the biological questions at stake may be summed up by discrete characters, why not design compartmental ODE models to take account of heterogeneity? However, in general, models structured in continuous traits, yielding integro-differential equations, or partial differential equations with diffusion and advection terms, should be the rule, as heterogeneity in biology is only coarsely represented by discrete variables.

C. Plasticity and tumour microenvironment

As mentioned earlier, plasticity understood as nongenetic instability may be represented by a Laplacian. This is the case in [3], [7], [8], where nongenetic instability or plasticity play a major role in the evolution of the solutions. The influence of evolutionary pressure (or stress) due to the tumour micro-environment is adequately represented in both cases by an advection (or drift) term, the specific effect

of which is studied in [9], and interpreted as displacing the expected concentration point of the one-dimensional structure variable of the nonlocal Lotka-Volterra reaction-diffusion under consideration. In [3], plasticity is endowed with the nature of an intrinsic trait tuning the Laplacian and evolving with time as the other (competing) traits. It is noteworthy that in order to represent competing phenotypes, such as fecundity and viability, a dimensionality of the space of phenotypes should be greater than one. If we add Cartesian space to phenotype space (as performed in [34], however with only one dimension in each case), this will make the model even more complex to study. Nevertheless, this is the setting of the development of actual solid tumours, in which local (in space) fixation of phenotypes may lead with genetic instability to the emergence of spatially isolated mutations according to a Darwinian-like phylogenetic tree as exemplified in [21]. In other words, should be considered firstly (reversible and costly in terms of energy to face micro-environmental stress) concentration of phenotypes in spatially isolated regions, and secondly, under the pressure of micro-environmental stress in these regions, emergence of mutations fixing on a less costly mode a favourable phenotype.

D. Interacting structured cell populations

Interactions of tumour cell populations with surrounding healthy ones may be mutualistic [42] or competitive. Structured models of tumour immune-interactions, for the latter, are in full development, since the beginning of modern immunotherapies, and they offer many opportunities, including for the development of control methods, to mathematicians.

VI. CONCLUSION AND PERSPECTIVES

In this review of mathematical models of tumour growth and its control, specifically focused on structured cell population models, principles of physiologically based modelling for cancer growth, with examples of models with and without control, have been presented. In the immediate future, modelling tumour-immune interactions with representation of modern immunotherapies involving tumour control by immune checkpoint inhibitors (that boost immune cells), and their adverse toxic events on healthy tissues, in particular muscular ones, is another goal which is presently pursued, with the same principle of structuring cell populations according to a continuous phenotype representing their heterogeneity. However, even immunotherapies act by tumour cell kill induced by drugs, and their adverse events, although rare, may be severe, even fatal, as in the case of cardiac or diaphragmatic myopathies. The grail of anticancer treatments might be, in particular by using redifferentiation therapies, to avoid cancer cell kill and its adverse toxic events on healthy tissues. Unfortunately, this has been achieved so far only in very special cases, such as acute promyelocytic leukaemia (APL) or chronic myelogenous leukaemia (CML) with drugs All-TransRetinoic Acid (ATRA) and Imatinib Mesylate, respectively. Progression in the understanding of mechanisms of tissue cohesion in multicellular organisms

and their recovery after local impairment in cancer might offer new opportunities for therapeutics, however most of them are still widely unknown.

REFERENCES

- [1] A. Aktipis, *et al.*, Cancer across the tree of life: cooperation and cheating in multicellularity, *Phil. Trans. R. Soc. B* 370:20140219 (2015)
- [2] L. Almeida, *et al.*, Evolution of cancer cell populations under cytotoxic therapy and treatment optimisation: insight from a phenotype-structured model, *ESAIM Math. Model. Numer. Anal.* 53:1157-1190 (2019)
- [3] F.E. Alvarez, *et al.*, Evolution of a structured cell population endowed with plasticity of traits under constraints on and between the traits, *J. Math. Biol.* 85:64, 49 pages (2022)
- [4] M. Bertolaso, *Philosophy of cancer. A dynamic and relational view*, Springer, Heidelberg, New York (2016)
- [5] B. Bratovsky & D. Horvath, Structure of Intratumor Heterogeneity: Is Cancer Hedging Its Bets? *arXiv,1307.0607* (2013)
- [6] K.J. Bussey, *et al.*, Ancestral gene regulatory networks drive cancer, *PNAS*, 114(24):6160-6162 (2017)
- [7] R.H. Chisholm, *et al.*, Emergence of drug tolerance in cancer cell populations: an evolutionary outcome of selection, non-genetic instability and stress-induced adaptation. *Cancer Research*, 75(6):930-939 (2015)
- [8] R.H. Chisholm, *et al.*, Cell population heterogeneity and evolution towards drug resistance in cancer: Biological and mathematical assessment, theoretical treatment optimisation, *Biochem. Biophys. Acta*, 1860:2627-2645 (2016)
- [9] R.H. Chisholm, *et al.*, Effects of an advection term in nonlocal Lotka-Volterra equations. *Comm. Math. Sciences*, 14(4):1181-1188 (2016)
- [10] L.H. Cisneros, *et al.*, Ancient genes establish stress-induced mutation as a hallmark of cancer. *PLoS One* 12(4):e0176258 (2017)
- [11] Clairambault, J. & Pouchol, C., A survey of adaptive cell population dynamics models of emergence of drug resistance in cancer, and open questions about evolution and cancer. *Biomath* 8:1905147, 23 pages, (2019)
- [12] Clairambault, J., Plasticity in cancer cell populations: biology, mathematics and philosophy of cancer. *Springer LNBI* 12508, pp. 3-9 (2020)
- [13] Davies, P.C.W. & Lineweaver, C.H., Cancer tumors as Metazoa 1.0: tapping genes of ancient ancestors, *Phys. Biol.* 8(1):015001, 7 pages (2011)
- [14] O. Diekmann, A beginner's guide to adaptive dynamics, *Mathematical Modelling of Population Dynamics*, vol. 63, Banach Center Publications, Institute of Mathematics, Polish Academy of Sciences, Warszawa, pp. 47-86 (2004)
- [15] L. Ding, *et al.*, Clonal evolution in relapsed acute myeloid leukaemia revealed by whole genome sequencing, *Nature* 481:506-510 (2012)
- [16] W. Djema, *et al.*, Control in Dormancy or Eradication of Cancer Stem Cells: Mathematical Modeling and Stability Issues. *Journal of Theoretical Biology*, 449:103-123 (2018)
- [17] T. Domazet-Lošo & D. Tautz, An ancient evolutionary origin of genes associated with human genetic diseases, *Mol. Biol. Evol.*, 25(12):2699-2707 (2008)
- [18] T. Domazet-Lošo & D. Tautz, Phylostratigraphic tracking of cancer genes suggests a link to the emergence of multicellularity in Metazoa, *BMC Biol.*, 8(1):66 (2010)
- [19] H. Easwaran, *et al.*, Cancer Epigenetics: Tumor Heterogeneity, Plasticity of Stem-like States, and Drug Resistance, *Molecular Cell*, 54:716-727 (2014)
- [20] A.P. Feinberg, *et al.*, Epigenetic modulators, modifiers and mediators in cancer aetiology and progression, *Nature Rev. Genet.*, 17(5):284-99 (2016)
- [21] M. Gerlinger, *et al.*, Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med* 366(10):883-892 (2012)
- [22] R.J. Gillies, D. Verduzco, R.A. Gatenby, Evolutionary dynamics of carcinogenesis and why targeted therapy does not work, *Nature Rev Cancer*, 12:487-493 (2012)
- [23] C.A. Gravenmier, *et al.*, Adaptation to Stochastic Temporal Variations in Intratumoral Blood Flow: The Warburg Effect as a Bet Hedging Strategy, *Bull Math Biol*, 80(5):954-970 (2017)
- [24] S. Huang, Genetic and non-genetic instability in tumor progression: link between the fitness landscape and the epigenetic landscape of cancer cells, *Canc. Metastasis Rev.*, 32:423-448 (2013)

- [25] L. Israel, Tumour progression: random mutations or an integrated survival response to cellular stress conserved from unicellular organisms? *J. Theor. Biol.*, 178(4):375-380 (1996)
- [26] A.M. Jarrett, *et al.*, Optimal Control Theory for Personalized Therapeutic Regimens in Oncology: Background, History, Challenges, and Opportunities, *J Clin Med* 9:1314 (2020)
- [27] W.O. Kermack, A.G. McKendrick, Contribution to the mathematical theory of epidemics, *Proc Roy Soc. A*, 115:700-721 (1927)
- [28] M. Kuznetsov, J. Clairambault, V. Volpert, Improving cancer treatments via dynamical biophysical models, *Physics of Life Reviews* 39:1-48 (2021)
- [29] U. Ledzewicz, *et al.*, On the role of tumor heterogeneity for optimal cancer chemotherapy, *Networks & Heterogeneous Media*, 14(1):131-147 (2019)
- [30] Lineweaver, C.H., Davies, P.C.W. & Vincent, M., Targeting cancer's weaknesses (not its strengths): Therapeutic strategies suggested by the atavistic model. *BioEssays* (2014)
- [31] Lineweaver, C.H., Davies, P.C.W. Comparison of the atavistic model of cancer to somatic mutation theory: Phylostratigraphic analyses support the atavistic model, Chap. 12. In: Gerstman BS (ed) *The Physics of Cancer. Research Advances*. World Scientific, Singapore, pp 243-261 (2020)
- [32] T. Lorenzi, *et al.*, Tracking the evolution of cancer cell populations through the mathematical lens of phenotype-structured equations. *Biology Direct*, 11:43 (2016)
- [33] A. Lorz, *et al.*, Populational adaptive evolution, chemotherapeutic resistance and multiple anti-cancer therapies, *ESAIM Math. Model. Numer. Anal.*, 47(2):377-399 (2013)
- [34] A. Lorz, *et al.*, Modeling the effects of space structure and combination therapies on phenotypic heterogeneity and drug resistance in solid tumors, *Bull. Math. Biol.*, 77(1):1-22 (2015)
- [35] A. Marusyk, *et al.*, Intra-tumour heterogeneity: a looking glass for cancer?, *Nature Rev. Cancer*, 12: 323-33 (2012)
- [36] K.D. McCullough, *et al.*, Plasticity of the neoplastic phenotype in vivo is regulated by epigenetic factors. *Proc Natl Acad Sci USA*, 95(26):15333-15338 (1998)
- [37] A.G. McKendrick, Applications of mathematics to medical problems, *Proc. Edinburgh Math. Soc.* 54, 98-13 (1926)
- [38] R.E. Michod, *et al.*, Life-history evolution and the origin of multicellularity, *J. Theor. Biol.*, 239:257-272 (2006)
- [39] W.E.G. Müller Review: How was metazoan threshold crossed? The hypothetical Urmetazoa. *Comp Biochem Physiol Part A Molecular Integr Physiol* 129(2):433-460 (2001)
- [40] W.E.G. Müller & I.M. Müller, Origin of the Metazoan Immune System: Identification of the Molecules and their Functions in Sponges, *Integrative Comparative Biology*, 43:281-292 (2003)
- [41] Nedelcu A.M., Michod R.E. Stress responses co-opted for specialized cell types during the early evolution of multicellularity. *BioEssays* 42(5):2000029 (2020)
- [42] T.N. Nguyen, *et al.*, Adaptive dynamics of hematopoietic stem cells and their supporting stroma: A model and mathematical analysis, *Math Biosci Eng*, 16(5): 4818-4845 (2019)
- [43] B. Perthame, *Transport equations in biology*, Birkhäuser, Springer (2007)
- [44] T. Philippi, J. Seger, Hedging one's evolutionary bets, revisited. *Trends Ecol Evolut* 4(2):41-44 (1989)
- [45] A.O. Pisco, *et al.*, Non-darwinian dynamics in therapy-induced cancer drug resistance, *Nat. Commun.* 4:2467 (2013)
- [46] A.O. Pisco & S. Huang, Non-genetic cancer cell plasticity and therapy-induced stemness in tumour relapse: 'What does not kill me strengthens me', *Br J Cancer*, 112(11):1725-1732 (2015)
- [47] Pouchol, C., Clairambault, J., Lorz, A. & Trélat, E., Asymptotic analysis and optimal control of an integro-differential system modelling healthy and cancer cells exposed to chemotherapy. *J. Math. Pures Appl.* 116, 268-308 (2018)
- [48] Pradeu, T., *The limits of the Self*, Oxford University Press (2012)
- [49] Pradeu, T., *Philosophy of immunology*, Cambridge University Press (2019)
- [50] H. Schättler & U. Ledzewicz, *Optimal Control for Mathematical Models of Cancer Therapies*, Springer (2015)
- [51] S.V. Sharma, *et al.*, A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations, *Cell*, 141(1):69-80 (2010)
- [52] Shen, S. & Clairambault, J.: Cell plasticity in cancer cell populations (review) [version 1; peer review: 2 approved]. *F1000Research* 2020, 9 (F1000 Faculty Rev) 635, 16 pages (2020)
- [53] Strauss B, Bertolaso M, Ernberg I, Bissell MJ. *Rethinking cancer: a new paradigm for the postgenomics era*. MIT Press (2021)
- [54] G. Tonini, *et al.*, Rechallenge therapy and treatment holiday: different strategies in management of metastatic colorectal cancer, *Journal of experimental & clinical cancer research*, 32(1):1 (2013)
- [55] A.S. Trigos, *et al.*, Altered interactions between unicellular and multicellular genes drive hallmarks of transformation in a diverse range of solid tumors, *PNAS* 114 (24):6406-6411 (2017)
- [56] M.D. Vincent, Cancer: a de-repression of a default survival program common to all cells?: a life-history perspective on the nature of cancer, *Bioessays*, 34(1):72-82 (2011)
- [57] G.P. Wagner, E.M. Erkenbrack, A.C. Love, Stress-induced evolutionary innovation: a mechanism for the origin of cell types. *BioEssays* 41(4):180-188 (2019)
- [58] S. Yuan, R.J. Norgard, B.Z. Stanger, Cellular plasticity in cancer. *Cancer Discov.* 9(7):837-851 (2019)