



HAL
open science

Strong Shift Equivalence and Bialgebras in traced monoidal categories

Emmanuel Jeandel

► **To cite this version:**

Emmanuel Jeandel. Strong Shift Equivalence and Bialgebras in traced monoidal categories. The 5th International Conference on Applied Category Theory 2022, 2022, Strathclyde, United Kingdom. hal-04318123

HAL Id: hal-04318123

<https://inria.hal.science/hal-04318123v1>

Submitted on 1 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Strong Shift Equivalence and Bialgebras in traced monoidal categories

arXiv:2107.10734

E. Jeandel


Université de Lorraine, France

Plan

- 1 Symbolic Dynamics
- 2 Categories for symbolic dynamics
- 3 Applications
- 4 Conclusion


Edge shift

An edge shift is the set of all biinfinite walks (on edges) in a finite graph




Edge shift

An edge shift is the set of all biinfinite walks (on edges) in a finite graph



Edge shift

An edge shift is the set of all biinfinite walks (on edges) in a finite graph



... *actactactact.actactactact* ...

... *teamteam.teamteam* ...


... *eeeeeeee.eeeeeeee* ...

... *actactactact.eeeeeeee* ...

... *actamteacte.amtamteact* ...


Conjugacy

Edge shifts are conjugate if they're isomorphic via local transformations



Conjugacy

Edge shifts are conjugate if they're isomorphic via local transformations




use different symbols for a
depending on the next symbol.

$ac \rightarrow ac$

$am \rightarrow bm$

Conjugacy

Edge shifts are conjugate if they're isomorphic via local transformations




use different symbols for a
depending on the next symbol.

$ac \rightarrow ac$

$am \rightarrow bm$

Conjugacy

Edge shifts are conjugate if they're isomorphic via local transformations




use same symbols for c and m

$ac \rightarrow ac$

$bm \rightarrow bc$

Conjugacy

Edge shifts are conjugate if they're isomorphic via local transformations




use same symbols for c and m

$ac \rightarrow ac$

$bm \rightarrow bc$

Conjugacy

Edge shifts are conjugate if they're isomorphic via local transformations



More generally, choose a vertex to duplicate

- Duplicate the inputs
- Redistribute the outputs

or vice-versa

SSE

This can be defined directly on adjacency matrices:

Definition

Two matrices M and N are 1-step equivalent if $M = RS$ and $N = SR$ for (nonnecessarily square) nonnegative integral matrices R, S

This can be defined directly on adjacency matrices:

Definition

Two matrices M and N are 1-step equivalent if $M = RS$ and $N = SR$ for (nonnecessarily square) nonnegative integral matrices R, S

Example:

$$M = \begin{pmatrix} 0 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix} \quad N = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

$$R = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad S = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$$

SSE

This can be defined directly on adjacency matrices:

Definition

Two matrices M and N are 1-step equivalent if $M = RS$ and $N = SR$ for (nonnecessarily square) nonnegative integral matrices R, S

Definition

Strong shift equivalence (SSE) is the transitive closure of 1-step equivalence.

Graphs G_1 and G_2 represent conjugate edge shifts iff their adjacency matrices are SSE.

Main open problem of symbolic dynamics: decide conjugacy/SSE

- Open since the 70s
- Decidable for matrices in \mathbb{Z} (Krieger, 1980)
 - (almost the) same as conjugacy in $GL_n(\mathbb{Z})$
- Decidable for one-sided edge-shifts (Williams, 1973)
 - The rewriting system on graphs is confluent.

Invariants

Definition

An invariant is a function ψ s.t. $\psi(M) = \psi(N)$ if M and N are SSE.

- To prove that M and N are NOT SSE, just find an invariant that distinguishes them

Examples:

- Trace of the matrix
- More generally, number of cycles of size k
- The entropy (exponential growth of the number of paths of size n)
- ...

In this talk: A systematic way to find invariants

Plan

- 1 Symbolic Dynamics
- 2 Categories for symbolic dynamics
- 3 Applications
- 4 Conclusion

Goal

Build a category where arrows can represent matrices and equivalence classes of SSE.

Starting point



Folklore

The bialgebra PROP is exactly the PROP of nonnegative integer matrices.

Starting point

Folklore

The bialgebra PROP is exactly the PROP of nonnegative integer matrices.


A commutative monoid $(\eta : 0 \rightarrow 1, \mu : 2 \rightarrow 1)$  

Starting point

Folklore

The bialgebra PROP is exactly the PROP of nonnegative integer matrices.

A commutative monoid $(\eta : 0 \rightarrow 1, \mu : 2 \rightarrow 1)$



A cocommutative comonoid $(\epsilon : 1 \rightarrow 0, \Delta : 1 \rightarrow 2)$




Starting point

Folklore

The bialgebra PROP is exactly the PROP of nonnegative integer matrices.



A commutative monoid ($\eta : 0 \rightarrow 1, \mu : 2 \rightarrow 1$)



A cocommutative comonoid ($\epsilon : 1 \rightarrow 0, \Delta : 1 \rightarrow 2$)



Bialgebra rules




Starting point

Folklore

The bialgebra PROP is exactly the PROP of nonnegative integer matrices.

arrows $n \rightarrow m$ are exactly matrices of size $m \times n$.

$$M = \begin{pmatrix} 0 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$$



Technically, our matrices represent graph, so with inputs = outputs

SSE states that $RS \equiv SR$, how can we simulate that ?

Technically, our matrices represent graph, so with inputs = outputs

SSE states that $RS \equiv SR$, how can we simulate that ?


With a trace!

Trace

Consider the traced bialgebra prop (the previous prop + a trace).

Trace



Consider the traced bialgebra prop (the previous prop + a trace).



Trace

In this prop, we can associate a scalar $t(M)$ to each matrix M .



$$M = \begin{pmatrix} 0 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$$



Trace


In this prop, we can associate a scalar $t(M)$ to each matrix M .
Is it true that $t(M) = t(N)$ iff M and N are SSE ?

$$M = \begin{pmatrix} 0 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$$



Flow equivalence

$M = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $N = (1)$ satisfy $t(M) = t(N)$ but are not SSE.



Flow equivalence

$t(M) = t(N)$ iff M and N are flow-equivalent, another equivalence notion coming from symbolic dynamics

(First observed by David Hillman, 1995)

We have lost a notion of time, that we need to recover

Solution


We add a generator that is a morphism for both the monoid and the comonoid.



Solution

We add a generator that is a morphism for both the monoid and the comonoid.

$$M = \begin{pmatrix} 0 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$$



Note: this new prop represents matrices in $\mathbb{Z}_+[t]$ rather than \mathbb{Z}_+

Solution

We add a generator that is a morphism for both the monoid and the comonoid.

Theorem

In this prop, $t(M) = t(N)$ iff M and N are SSE.

Needs some new (easy) results in symbolic dynamics for the proof.

To recap: we need one bialgebra, a bialgebra morphism and a trace.

Plan

- 1 Symbolic Dynamics
- 2 Categories for symbolic dynamics
- 3 Applications**
- 4 Conclusion

What do we gain from it ?

- If one knows a concrete representation of this prop, one can decide conjugacy
- One can find invariants !

- Let \mathcal{C} be our category.
- Let \mathcal{D} be any traced category that contains a bialgebra
- By the universal property, there is a functor from \mathcal{C} to \mathcal{D}
- Using this functor, we can associate to each matrix M an arrow $\psi(M)$ in \mathcal{D} s.t. $\psi(M) = \psi(N)$ if M and N are SSE.

Good news: There are a lot of bialgebras in the wild

Bad news:

- Some of them are in categories that are not traced
- Some of them are Hopf algebras
 - Hopf algebras represent matrices with coefficients in \mathbb{Z} , not in \mathbb{Z}_+

Example from algebra 1/2

- If M is a finite abelian monoid, $\mathbb{K}[M]$ has a structure of a bialgebra
 - Product is the monoid product extended linearly
 - Coproduct is the copy extended linearly

Theorem

Let \mathcal{M} be an additive monoid, and h an homomorphism. Then the number of solutions in \mathcal{M} of the equation $h(Mx) = x$ is an invariant for SSE.

Example from algebra 2/2

- $\mathbb{K}[X]$ has a structure of a bialgebra
 - Product is the product of polynomials
 - Coproduct is defined by $\Delta(X) = 1 \otimes X + X \otimes 1$

Problem: no trace!

Example from algebra 2/2

- $\mathbb{K}[X]$ has a structure of a bialgebra
 - Product is the product of polynomials
 - Coproduct is defined by $\Delta(X) = 1 \otimes X + X \otimes 1$

Solution: Replace $\mathbb{K}[X]$ by formal series in a complete semiring (like $\mathbb{R}_+ \cup \{+\infty\}$).

If we take the morphism $h(X) = tX$ we get:

Theorem

The quantity $f_M(t) = \frac{1}{\det(I-tM)}$ is an invariant for SSE

This is the well-known Zeta function of a subshift.

Example from Category Theory

Let \mathcal{D} be a category with finite limits and colimits.

- Suppose one has a monoid X in a \mathcal{D} with a morphism h .
- It extends to a bialgebra using the diagonal map as a comonoid.

Problem: What if there is no trace in \mathcal{D} ?

Example from Category Theory

Let \mathcal{D} be a category with finite limits and colimits.

- Suppose one has a monoid X in a \mathcal{D} with a morphism h .
- It extends to a bialgebra using the diagonal map as a comonoid.

Solution: Replace \mathcal{D} with $\text{cospan}(\mathcal{D})$ which is a compact category.

Example from Category Theory

Let \mathcal{D} be a category with finite limits and colimits.

- Suppose one has a monoid X in a \mathcal{D} with a morphism h .
- It extends to a bialgebra using the diagonal map as a comonoid.

Theorem

Let M be a $n \times n$ matrix.

The object that is the coequalizer of $X^n \begin{array}{c} \xrightarrow{id^n} \\ \xrightarrow{h^n \circ M} \end{array} X^n$ is an invariant for SSE.

Example from Category Theory

Let \mathcal{D} be a category with finite limits and colimits.

- Suppose one has a monoid X in a \mathcal{D} with a morphism h .
- It extends to a bialgebra using the diagonal map as a comonoid.

Theorem

Let M be a $n \times n$ matrix.

The object that is the coequalizer of $X^n \begin{array}{c} \xrightarrow{id^n} \\ \xrightarrow{h^n \circ M} \end{array} X^n$ is an invariant for SSE.

- Starting with the monoid $(\mathbb{Z}, +)$ in the category of abelian groups, we obtain the *Bowen-Franks* group (1977)
- Starting with the monoid $(\mathbb{Z}[t], +)$ in the category of $\mathbb{Z}[t]$ -modules, we obtain the dimension group of Krieger (1977)

Plan

- 1 Symbolic Dynamics
- 2 Categories for symbolic dynamics
- 3 Applications
- 4 Conclusion

Where to go from here ?

- We have a systematic way to find invariants
- Investigate new bialgebras to find new invariants!