

# High dimensional clustering of coupound precipitation and wind extremes over Europe

Gwladys Toulemonde, Alexis Boulin, Elena Di Bernadino, Thomas Laloë

## ▶ To cite this version:

Gwladys Toulemonde, Alexis Boulin, Elena Di Bernadino, Thomas Laloë. High dimensional clustering of coupound precipitation and wind extremes over Europe. ICSDS 2023 - IMS International Conference on Statistics and Data Science, IMS, Dec 2023, Lisbon, Portugal. hal-04373309

## HAL Id: hal-04373309 https://inria.hal.science/hal-04373309

Submitted on 15 Jan2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

### High-dimensional clustering of compound precipitation and wind extremes over Europe

# $\frac{\text{Gwladys Toulemonde}^1, \text{ Alexis Boulin}^2, \text{ Elena Di}}{\text{Bernardino}^3, \text{ Thomas Laloë}^3}$

 <sup>1</sup> Univ Montpellier, CNRS, IMAG and Inria Lemon, France, gwladys.toulemonde@umontpellier.fr
<sup>2</sup> Université Côte d'Azur, CNRS, LJAD and Inria Lemon, France
<sup>3</sup> Université Côte d'Azur, CNRS, LJAD, France

#### Abstract

Disastrous climate events such as floods, wildfires, and heatwaves often occur due to the simultaneous extreme behaviour of several interacting processes. Since in these compound events several spatiotemporal factors are jointly extreme and by their very nature are of high dimension, it is for a proper understanding of them to develop dependence summary measures that are appropriate for extreme value random vectors. These latter is a key ingredient to propose spatial clustering of these temporal processes. Based on the recent development of an algorithm specifically tailored for AI-block models (see Boulin et al., 2023) we propose in this talk a clustering method adapted to compound extreme events. We exemplify this method proposing a regionalization task. More precisely we identify regions based on gridded data from observations and climate model ensembles over Europe. This approach uses daily precipitation sums and daily maximum wind speed data from the ERA5 reanalysis dataset from 1979 to 2022.

### Keywords

Spatial clustering, compound extremes, extremal dependence.

### References

Boulin, A., Di Bernardino, E., Laloë, T., Toulemonde G. (2023). Highdimensional variable clustering based on sub-asymptotic maxima of a weakly dependent random process. arXiv:2302.00934.

108