

Asymptotic estimations of a perturbed symmetric eigenproblem

Armand Gissler, Anne Auger, Nikolaus Hansen

To cite this version:

Armand Gissler, Anne Auger, Nikolaus Hansen. Asymptotic estimations of a perturbed symmetric eigenproblem. Applied Mathematics Letters, 2024, 150, pp.108951. $10.1016/j.$ aml.2023.108951. hal-04386103

HAL Id: hal-04386103 <https://inria.hal.science/hal-04386103>

Submitted on 10 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International](http://creativecommons.org/licenses/by-nc-sa/4.0/) [License](http://creativecommons.org/licenses/by-nc-sa/4.0/)

Asymptotic estimations of a perturbed symmetric eigenproblem

Armand Gissler^{a,∗}, Anne Auger^a, Nikolaus Hansen^a

a Inria, CMAP, CNRS, Ecole polytechnique, Institut Polytechnique de Paris, 91120 Palaiseau, France ´

Abstract

We study ill-conditioned positive definite matrices that are disturbed by the sum of *m* rank-one matrices of a specific form. We provide estimates for the eigenvalues and eigenvectors. When the condition number of the initial matrix tends to infinity, we bound the values of the coordinates of the eigenvectors of the perturbed matrix. Equivalently, in the coordinate system where the initial matrix is diagonal, we bound the rate of convergence of coordinates that tend to zero.

Keywords: Perturbation of symmetric matrices, estimation of eigenvectors *2020 MSC:* 15A42, 15B57

1. Introduction

Given a $d \times d$ symmetric matrix with known eigenvectors and eigenvalues denoted *B*, and a rank-one matrix vv^T where $v \in \mathbb{R}^d$, eigenvalues and eigenvectors of matrices of the form

$$
A = B + v v^T \tag{P_1}
$$

have been widely studied, notably in the context of perturbation theory. For instance, the eigenvalues of (*P*1) can be estimated and a formula for the eigenvectors is known [1, 2, 3]. Specifically, if *B* is a diagonal matrix $diag(\lambda_1, \lambda_2, \ldots, \lambda_d)$ with distinct eigenvalues and *v* has only nonzero entries, then the component *j* of the unit eigenvector associated to eigenvalue ^ν*ⁱ* of the updated matrix *^A* satisfies the so-called Bunch-Nielsen-Sorensen formula

$$
C_i \times \frac{[v]_j}{\lambda_j - v_i} \quad \text{for } i, j = 1, ..., d
$$
 (1)

where C_i is a nonzero normalization constant and $[.]_j$ denotes the *j*-th coordinate, a notation we will continue to use in the sequel. Several results have been established for additive perturbations of rank 1 [4, 5] use in the sequel. Several results have been established for additive perturbations of rank 1 [4, 5] and of higher rank [6, 7]. Symmetric and nonsymmetric perturbation eigenvalue problems have been studied [8] as well as perturbation results for invariant subspaces [9]. In this paper, we provide relative perturbation bounds for the eigenvectors of positive definite matrices. In contrast to previous relative perturbation results for eigenvalues [10] and invariant subspaces [11, 12], the bounds in our result depend on the eigenvalues of the initial matrix *B* rather than the norm of the perturbation, see Eq. (2) below.

Specifically, we consider the perturbation with a sum of *m* rank-one matrices of the form

$$
A^{(m)} = B + \sqrt{B} \sum_{i=1}^{m} [v^{(i)}][v^{(i)}]^T \sqrt{B}
$$
 (P_m)

with $B = P \text{diag}(\lambda_1, ..., \lambda_d) P^T$ where *P* is an orthogonal matrix, $\lambda_1 \geq ... \geq \lambda_d > 0$ are the eigenvalues of *B*, and $v^{(1)} = v^{(m)} \in \mathbb{R}^d$. The square root $\sqrt{B} = P \text{diag}(\sqrt{\lambda_1}, ..., \sqrt{\lambda_d}) P^T$ is defined as the unique symmetr with $B = P$ diag($\lambda_1, ..., \lambda_d$) P ² where *P* is an orthogonal matrix, $\lambda_1 \geq \cdots \geq \lambda_d > 0$ are the eigenvalues of *B*, and $v^{(1)}, ..., v^{(m)} \in \mathbb{R}^d$. The square root $\sqrt{B} := P$ diag($\sqrt{\lambda_1}, ..., \sqrt{\lambda_d}$) P^T is defined as the un

[∗]Corresponding author

Email addresses: firstname.lastname@polytechnique.edu (Armand Gissler), firstname.lastname@inria.fr (Anne Auger), firstname.lastname@inria.fr (Nikolaus Hansen)

Preprint submitted to Applied Mathematics Letters January 10, 2024

definite matrix such that $\sqrt{B} \times$ √ $B = B$, see e.g. [13, Theorem 7.2.6]. Matrices of the form (P_m) are used in various applications in different domains. For instance, low rank updates of covariance matrices are used in stochastic optimization [14, 15], system identification [16, p. 369], and adaptive Markov Chain Monte Carlo methods [17]. Our motivation is to study the eigenvectors of *A* in (P_m) , denoted as $e_i^{(m)}$ in the sequel, *when the matrix B is highly illconditioned*. When $d = 2$ and $m = 1$ we can compute the eigenvectors of $A^{(1)}$ explicitly. As an example, consider $B = \text{diag}(\lambda_1, 1)$ where $\lambda_1 > 1$ and $v^{(1)} = [1, 1]^T$. Then, the unit eigenvector associated to the largest eigenvalue of *A*⁽¹⁾ obeys $\sqrt{1 + s^2} \times e_1^{(1)}$ $\lambda_1^{(1)}$ = [1, *s*]^{*T*} with *s* = $\lambda_1^{1/2}$ × (1 − λ_1^{-1} − $\sqrt{1 - \lambda_1^{-1} + \lambda_1^{-2}}$) = $-\lambda_1^{-1/2}/2 + O(\lambda_1^{-3/2})$ and hence $[e_1^{(1)}]$ ⁽¹⁾₁₂ = $\lambda_1^{-1/2} + o(\lambda_1^{-1/2})$ when $\lambda_1 \to \infty$. Hence, the (second) coordinate of the (first) unit eigenvector of *A*⁽¹⁾ vanishes like $1/\sqrt{\lambda_1}$ when $\lambda_1 \to \infty$. In this paper, we generalize this result to the case where $d \ge 2$ and $m \ge 1$, as summarized in the following theorem which directly follows from Theorem 4 below. in the following theorem which directly follows from Theorem 4 below.

Theorem 1. If $e_i^{(m)}$ is a unit eigenvector corresponding to the *i*-th largest eigenvalue (counted with multiplicity) of *i* $A^{(m)}$ in (P_m) and $e_j^{(0)}$ is a unit eigenvector corresponding to the j-th largest eigenvalue λ_j of B, then

$$
\left| \left\langle e_i^{(m)}, e_j^{(0)} \right\rangle \right| \leq C_m \times \sqrt{\frac{\min\{\lambda_i, \lambda_j\}}{\max\{\lambda_i, \lambda_j\}}}
$$
\n(2)

where $C_m > 0$ *is a constant which depends polynomially on d and* $\max_{k=1,\dots,m} ||v^{(k)}||$.

When *B* is diagonal, $e_j^{(0)}$ is the *j*-th canonical unit vector. Hence $|\langle e_i^{(m)}, e_j^{(0)} \rangle| = [e_i^{(m)}]_j$ and the theorem implies in particular that the *j*-th coordinate of $e_i^{(m)}$ converges to zero at least as fast as $\sqrt{\min\{\lambda_i, \lambda_j\}} \max\{\lambda_i, \lambda_j\}$ when the latter tends to 0 (which is tight in the above example when $d = 2$ and $m = 1$), thereby limiting the change of the angle between these eigenvectors. Considerations on the angle between eigenspaces have been made previously [18], however matrices on the form of (P_m) have not been studied in this context. In the remainder, we always choose w.l.o.g. the coordinate system where the matrix *B* of (P_m) is diagonal and has decreasingly ordered diagonal values.

This inequality is crucial to study the stability of a Markov chain underlying the CMA-ES algorithm [19, 15]. Proofs of linear convergence for Evolutionary Strategies (ES) rely on a drift condition [20, Theorem 17.0.1] to prove the ergodicity of an underlying Markov chain, see e.g. [21, 22]. To apply this approach to CMA-ES, a potential function is defined on the state-space of this Markov chain and its expected decrease is proven outside a compact set. The state space includes a covariance matrix, updated as

$$
C_{t+1} = (1 - c)C_t + c\sqrt{C_t} \sum_{i=1}^{m} w_i U_i U_i^T \sqrt{C_t} ,
$$
\n(3)

where $c \in [0, 1], w_1, \ldots, w_m$ are positive weights that sum to 1, and the vectors U_i , $i = 1, \ldots, m$, are Gaussian vectors ranked according to a fitness function [15] Eq. (11)]. Hence (P) accompasses the undate of this cov ranked according to a fitness function [15, Eq. (11)]. Hence, (*Pm*) encompasses the update of this covariance matrix. Eq. (2) is needed to bound the expected condition number of the updated covariance matrix, since it controls the influence of small eigenvalues on the growth of the largest eigenvalues.

This paper is organized as follows. In Section 2, we study the eigenvalues of (P_m) . In Section 3, we provide bounds for the coordinates of the eigenvectors using Eq. (1), and provide an empirical result suggesting that these bounds are tight.

2. Bounds on the eigenvalues of (P_m)

The Bunch-Nielsen-Sorensen formula (1) which we will use in Section 3 requires the eigenvalues of the *updated* matrix. Thus, we first derive bounds on the (decreasingly ordered) eigenvalues

$$
\lambda_i(A^{(m)}) = \max_{V \subset \mathbb{R}^d, \dim V = i} \min_{v \in V, v \neq 0} \frac{v^T A^{(m)} v}{v^T v} = \min_{V \subset \mathbb{R}^d, \dim V = d - i + 1} \max_{v \in V, v \neq 0} \frac{v^T A^{(m)} v}{v^T v} \quad \text{for } i = 1, ..., d
$$
 (4)

where the equalities ensue from the min-max principle and from Gersgorin's circle theorem [13, Theorems 4.2.6 and 6.1.1].

Theorem 2.7 in [10] and Theorem 2.1 in [23, p. 175] provide an estimation for the eigenvalues of (*Pm*):

$$
\lambda_i \leqslant \nu_i \leqslant \lambda_i \times \left(1 + md \times \max_{k=1,\dots,m} ||\nu^{(k)}||_{\infty}^2\right) \quad \text{for } i \in \{1,\dots,d\} \quad . \tag{5}
$$

The next lemma provides a slightly tighter upper bound on these eigenvalues after a single rank-one pertubation $(m = 1)$ and is used in Proposition 3.

Lemma 2. Let $D = \text{diag}(\lambda_1, \dots, \lambda_d)$ be a diagonal matrix with $\lambda_1 > \dots > \lambda_d > 0$. Let $v \in \mathbb{R}^d_+$ be a vector with only *nonzero entries.* Let $A = D + \sqrt{D}vv^T \sqrt{D}$ and $v_1 \ge v_2 \ge \cdots \ge v_d$ denote the eigenvalues of A. Then,

$$
\nu_i \leq \lambda_i \times \left(1 + (d - i + 1) \|\nu\|_{\infty} \|\nu\|_{j_i}\right) \quad \text{for all } i \in \{1, \dots, d\}
$$

where $j_i \in \text{Arg max}_{j=i,\dots,d} \left\{ |[v]_j| \mid \lambda_j \geq \lambda_i \times \left(1 - \sqrt{\frac{\lambda_j}{\lambda_i}}\right) \right\}$ $\frac{\overline{\lambda_j}}{\overline{\lambda_i}}(d-i+1) \|\nu\|_{\infty} |[\nu]_j| \bigg)$

Proof. Fix $i \in \{1, ..., d\}$ and remark that, by Eq. (4), we have $v_i \le \max_{v \in \bar{V}_i, ||v|| = 1} v^T A v = \lambda_1 ([A]_{i:d,i:d})$, where $\bar{V}_i =$
Vect(e, e) with e being the ith vector of the standard basis of \mathbb{R}^d and with [4], ... Vect (e_i, \ldots, e_d) with e_i being the *i*th vector of the standard basis of \mathbb{R}^d , and with $[A]_{i,d,i,d}$ denoting the submatrix of A from rows and columns with indices between *i* and *d* included. But, by [13] Theore from rows and columns with indices between *i* and *d* included. But, by [13, Theorem 6.1.1], we also have that

$$
\lambda_1\left([A]_{i:d,i:d}\right) \leqslant \max_{j=i,\dots,d} \left(\sum_{k=i}^d |[A]_{j,k}|\right) =: \max_{j\geqslant i} B_j.
$$

Since $A = D +$ $\sqrt{D}vv^T \sqrt{D}$, then $[[A]_{j,k}] \le \sqrt{\lambda_j \lambda_k} (\mathbb{1}\{j = k\} + ||v||_{\infty}[[v]_j])$. If $j \ge i$ is such that $\lambda_j \ge \lambda_i \times (1 - \sqrt{\lambda_j/\lambda_i} (d - \lambda_j))$
then by definition of it we have then $[[v]_j] \le ||v||_{\infty}$ it vielding to $R_i \le \lambda_i \times (1 + (d - i + 1)||v||$ $i + 1$ ||v||_{|∞}|[v]_j|), then by definition of j_i we have then $[[v]_j] \leq [[v]_{j_i}]$, yielding to $B_j \leq \lambda_i \times (1 + (d - i + 1) ||v||_{\infty}[[v]_{j_i}])$.
Any other $i > i$ satisfies $\lambda_i < \lambda_j = \sqrt{\lambda_j \lambda_j} (d - i + 1) ||v||$. [[v], hence by sum $B_i < \$ Any other $j \ge i$ satisfies $\lambda_j < \lambda_i - \sqrt{\lambda_j \lambda_i} (d - i + 1) ||\nu||_{\infty} |[\nu]_j|$, hence by sum $B_j \le \lambda_i$. All in all, $\max_{j \ge i} B_j \le \lambda_i \le (1 + (d - i + 1) ||\nu|| \cdot ||\nu||)$ proving Eq. (6) $\lambda_i \times (1 + (d - i + 1) ||v||_{\infty} |[v]_{j_i}|)$, proving Eq. (6).

3. Estimating the eigenvectors of (P_m)

We use the bounds from Lemma 2 and Eq. (5) to estimate the eigenvectors of (P_m) by applying Eq. (1), for $m = 1$ in the next section and $m \geq 1$ in Section 3.2.

3.1. Rank-one perturbation

In Proposition 3, we obtain bounds on the coordinates of the eigenvectors of (P_m) (B is assumed to be diagonal) when $m = 1$, which comes as a consequence of Eq. (1).

Proposition 3. Let $D = \text{diag}(\lambda_1, ..., \lambda_d)$ be a diagonal matrix with $\lambda_1 \geq ... \geq \lambda_d > 0$. Let $v \in \mathbb{R}^d$ and $V := \text{max} \{d^{-1/2} \text{ [to]} \}$ Consider the matrix $A = D + \sqrt{D}w^T \sqrt{D}$ and $v_i \geq ... \geq v_i$ its eigenvalues and $(e^{(1)} \max\{d^{-1/2}, ||v||_{\infty}\}\$. Consider the matrix $A = D + c$ corresponding orthonormal basis of eigenvector *values and matrix with* $\lambda_1 \geq \cdots \geq \lambda_d > 0$. Let $v \in \mathbb{R}^d$.
 $\sqrt{D}vv^T \sqrt{D}$ and $v_1 \geq \cdots \geq v_d$ its eigenvalues and $(e_1^{(1)})$.

Then $e_1^{(1)}, \ldots, e_d^{(1)}$ *d*) *a corresponding orthonormal basis of eigenvectors. Then,*

$$
\left| [e_i^{(1)}]_j \right| \leqslant 5d^2 V^4 \sqrt{\frac{\min\{\lambda_i, \lambda_j\}}{\max\{\lambda_i, \lambda_j\}}} \quad \text{for all } i, j \in \{1, \dots, d\} \tag{7}
$$

Proof. We prove first that, if $\max\{\lambda_i, \lambda_j\} > (1 + dV^2) \times \min\{\lambda_i, \lambda_j\}$, then

$$
\left| [e_i^{(1)}]_j \right| \leq (d-i+1)V^2 \times \frac{\inf_{\rho \in (0,1)} \psi(\rho, (d-i+1)V^2)}{1 - (1 + (d-i+1)V^2) \frac{\min\{\lambda_i, \lambda_j\}}{\max\{\lambda_i, \lambda_j\}}} \sqrt{\frac{\min\{\lambda_i, \lambda_j\}}{\max\{\lambda_i, \lambda_j\}}} \tag{8}
$$

with $\psi(\rho, W) = \max\{2(1 - \rho)^{-1/2}, 2\rho^{-1}W\}$, from which we deduce Eq. (7).

First suppose that the eigenvalues λ_i of *D* are distinct, and that all entries of *v* are nonzero. Then, by Eq. (1), we have for *i*, *j* ∈ {1, ..., *d*} that $[e_i^{(1)}]_j = C_i \frac{[\sqrt{D}v]_j}{\lambda_j - v_i}$ $\frac{\Delta V}{\lambda_j - \nu_i} = C_i$ $\frac{\sqrt{\lambda_i}[\nu]}{i}$ $\frac{\partial \lambda_i[v]}{\partial_j - v_i}$, where $C_i \in \mathbb{R}$ is chosen such that $||e_i^{(1)}|| = 1$, hence

$$
|C_i| = \left\| \left(\frac{\sqrt{\lambda_j} [v]_j}{\lambda_j - v_i} \right)_{j=1,\dots,d} \right\|^{-1} = \left(\sum_{j=1}^d \left| \frac{\sqrt{\lambda_j} [v]_j}{\lambda_j - v_i} \right|^2 \right)^{-1/2} \leq \min_{1 \leq j \leq d} \frac{|\lambda_j - v_i|}{\sqrt{\lambda_j} |[v]_j|}.
$$
 (9)

Combining [23, Theorem 2.1, p. 175] with Eq. (6), we have $\lambda_i < \nu_i \leq \lambda_i \times (1 + (d - i + 1)V[[\nu]_i])$, where j_i is defined
in Lemma 2. By definition of *i*, we have $0 < \lambda_i - \lambda_i < \lambda_i (d - i + 1)V[[\nu]_i]$. By sum, we obtain $0 < \nu_i - \lambda_i <$ in Lemma 2. By definition of *j_i* we have $0 \le \lambda_i - \lambda_j$, $\le \lambda_i(d - i + 1)V[[v]_{j_i}]$. By sum, we obtain $0 \le v_i - \lambda_{j_i} \le 2\lambda(d - i + 1)V[[v]_{j_i}]$. $2\lambda_i(d - i + 1)V[[v]_{j_i}]$. We apply this to Eq. (9) to get

$$
|C_i| \leqslant \frac{\nu_i - \lambda_{j_i}}{\sqrt{\lambda_{j_i}} |[\nu]_{j_i}|} \leqslant \frac{2\lambda_i (d - i + 1)V|[\nu]_{j_i}|}{\sqrt{\lambda_{j_i}} |\nu|_{j_i}|} = \frac{\lambda_i}{\sqrt{\lambda_{j_i}}} 2(d - i + 1)V \tag{10}
$$

Let $\rho \in (0, 1)$. If $(d - i + 1)V^2 \sqrt{\lambda_{j_i}} \leq \rho \sqrt{\lambda_j}$
 $(1 - e)^{-1/2} \times 2(d - i + 1)V^2 \sqrt{\lambda_j}$ Otherwise λ*i* , then by definition of j_i , λ_j $\geq \lambda_i \times (1 - \rho)$, and by Eq. (10), $|C_i| \leq$ $(1 - \rho)^{-1/2} \times 2(d - i + 1)V \sqrt{}$ $\sqrt[n]{\lambda_j}$ $\leq P \sqrt{\lambda_i}$, then by definition of *j_i*, $\sqrt[n]{\lambda_j}$
 $\sqrt[n]{\lambda_i}$. Otherwise, $|C_i| \leq P^{-1} \times 2(d - i + 1)^2 V^3 \sqrt{2d - 1}$ $\overline{\lambda_i}$. All in all, for $\rho \in (0, 1)$,

$$
|C_i| \leq (d-i+1)V\sqrt{\lambda_i} \times \max\left\{2(1-\rho)^{-1/2}, 2\rho^{-1}(d-i+1)V^2\right\} =: C_\rho \sqrt{\lambda_i} \tag{11}
$$

Then, $| [e_i^{(1)}]_j | = |C_i| \sqrt{\lambda_j} | [v]_j | / |\lambda_j - v_i | \le \sqrt{\lambda_i \lambda_j} / |\lambda_j - v_i| \times \inf_{\rho \in (0,1)} C_\rho$. By Eq. (6), when $\lambda_j < \lambda_i$, $| [e_i^{(1)}]_j | \le$ | λ*j* $\min_{\rho \in [0,1]} C_{\rho} \times (1 - \lambda_j/\lambda_i)^{-1} \sqrt{\lambda_j/\lambda_i}$. By Eq. (5), when $\lambda_j > (1 + dV^2)\lambda_i$, $|[e_i^{(1)}]_j| \leq \min_{\rho \in [0,1]} C_{\rho} \times (1 - (1 + dV^2)\lambda_i)$ $dV^2 \frac{\lambda_i}{\lambda_j}$ $\frac{1}{\lambda_j} \frac{\sqrt{\lambda_i}{\lambda_j}}{V^2}$

. If the eigenvalues of *D* are not distinct or not all entries of *v* are nonzero, we consider a sequence of diagonal matrices $\{D_k = \text{diag}(\lambda_1^k, \dots, \lambda_d^k)\}_{k \in \mathbb{N}}$ such that the diagonal elements $\lambda_1^k > \lambda_2^k > \dots > \lambda_d^k > 0$ are distinct and $D_k \to D$
when $k \to \infty$ and a sequence of vectors $\{y_k \in \mathbb{R}^d\}_{k \in \mathbb{N}}$ with only nonzero en when $k \to \infty$, and a sequence of vectors $\{v_k \in \mathbb{R}^d\}$ *k*∈N with only nonzero entries where $||v_k||_{\infty} \le N$ and $v_k \to v$ when $k \to \infty$. Denote then $A_k = D_k + \sqrt{D_k} v_k v_k^T \sqrt{D_k}$ and $v_1^k \geq \cdots \geq v_d^k$ its eigenvalues. Note that $A_k \to A$ when $k \to \infty$ so by continuity of the eigenvalues $v_1^k \to v$, when $k \to \infty$. Eurthermore, we just proved th $k \to \infty$, so by continuity of the eigenvalues, $v_i^k \to v_i$ when $k \to \infty$. Furthermore, we just proved that if e_1^k, \ldots, e_d^k are unit eigenvalues of 4, corresponding respectively to the eigenvalues v^k at then e^k an unit eigenvectors of A_k corresponding respectively to the eigenvalues v_1^k, \ldots, v_d^k , then e_1^k, \ldots, e_d^k and $\lambda_1^k, \ldots, \lambda_d^k$ satisfy
Eq. (8) Moreover, the vectors e^k all belong to the unit sphere of \mathbb{R}^d Eq. (8). Moreover, the vectors e_i^k all belong to the unit sphere of \mathbb{R}^d , so up to considering a subsequence of $\{A_k\}_{k\in\mathbb{N}}$, we can assume w.l.o.g. that each e_i^k tends to a vector $e_i^{(1)} \in \mathbb{R}^d$ when $k \to \infty$. As (e_1^k, \ldots, e_d^k) is an orthonormal system of e_i^k and $e_i^{(1)}$ is an $e_i^{(1)}$ is an $e_i^{(1)}$ is an $e_i^{(1)}$ is an $e_i^{(1)}$ of \mathbb{R}^d , so is its limit ($e_1^{(1)}$) $e_d^{(1)}, \ldots, e_d^{(1)}$
nit $k \to \infty$ ⁽¹⁾) and $e_i^{(1)}$ is an eigenvector of *A* corresponding to the eigenvalue v_i . Therefore, Eq. (8) ∞ in the equation satisfied by e^k and e^{k} and e^{k} holds by taking the limit $k \to \infty$ in the equation satisfied by e_1^k, \ldots, e_k^k and $\lambda_1^k, \ldots, \lambda_d^k$.
To obtain Eq. (7), note that when $q = 1/2$, we have $2(1 - q)^{-1/2} < 4 < 4dV^2$ and

To obtain Eq. (7), note that when $\rho = 1/2$, we have $2(1 - \rho)^{-1/2} \leq 4 \leq 4dV^2$, and $2\rho^{-1}(d - i + 1)V^2 \leq 4dV^2$, and $2\rho^{-1}(d - i + 1)V^2 \leq 4dV^2$, and $2\rho^{-1}(d - i + 1)V^2 \leq 4dV^2$. when $\max\{\lambda_i, \lambda_j\} / \min\{\lambda_i, \lambda_j\} > 1 + 4dV^2$, by Eq. (8), then,

$$
(1 - (1 + dV^2) \min\{\lambda_i, \lambda_j\} / \max\{\lambda_i, \lambda_j\})^{-1} \leq (1 + 4dV^2) / (4dV^2) \leq 5/4,
$$

as $V \ge d^{-1/2}$, and thus Eq. (7) holds. If otherwise $\max\{\lambda_i, \lambda_j\} \le (1+4dV^2) \min\{\lambda_i, \lambda_j\}$, as $\max\{\lambda_i, \lambda_j\} / \min\{\lambda_i, \lambda_j\} \ge 1$, we find $|[e_i]_j| \le 1 \le (1+4dV^2)\sqrt{\max\{\lambda_i, \lambda_j\} / \min\{\lambda_i, \lambda_j\}}$. Since $1 \le dV^2$, then $(1+4dV^2) \$ holds. \Box

3.2. Sum of m rank-one matrices pertubation

Our final Theorem 4 generalizes Proposition 3 to any value $m \geq 1$ and is obtained by induction using Eq. (8). Theorem 4 implies in particular Theorem 1 via the spectral theorem.

Theorem 4. Let $D = \text{diag}(\lambda_1, ..., \lambda_d)$ be a diagonal matrix with $\lambda_1 \geq \cdots \geq \lambda_d > 0$. Let $V \geq 1/\sqrt{N}$
sequence of vectors $v^{(i)} \in \mathbb{R}^d$ such that $||v^{(i)}|| \leq V$ for all $i \in \mathbb{N}$. For $m \in \mathbb{N}$ let $A^{(m)} - D + \sqrt{D}$ $\geq 1/\sqrt{d}$ and consider a sequence of vectors $v^{(i)} \in \mathbb{R}^d$ such that $||v^{(i)}||_{\infty} \leqslant V$ for all $i \in \mathbb{N}$. For $m \in \mathbb{N}$, let $A^{(m)} = D + \sqrt{D} \sum_{i=1}^m [v^{(i)}][v^{(i)}]^T \sqrt{D}$ *and* $v_1^{(m)} \geqslant \cdots \geqslant v_d^{(m)}$
Then $\sigma_d^{(m)}$ the eigenvalues of $A^{(m)}$ and $(e_1^{(m)})$ $e_1^{(m)}, \ldots, e_d^{(m)}$ *d*) *a corresponding orthonormal system of eigenvectors. Then*

$$
|[e_i^{(m)}]_j| \leq C_m \sqrt{\frac{\min\{\lambda_i, \lambda_j\}}{\max\{\lambda_i, \lambda_j\}}}
$$
 for all $i, j \in \{1, ..., d\}$ and $m \in \mathbb{N}$ (12)

with $C_0 = 1$ *and* $C_{m+1} = 5d^7V^4C_m^5$ $1 + dmV^2$.

Figure 1: Value of $| [e_1^{(m)}]_j |$ as a function of λ_1/λ_j where $e_1^{(m)}$ is an eigenvector associated to the largest eigenvalue of $A^{(m)}$ from (P_m) for different dimensions and values of *m* as given in the legend. The $v^{(i)}$ are independent standard Gaussian vectors (with the same realization for all values of λ_1) and the eigenvalues of the diagional matrix *B* are chosen uniformly on a log scale between $\lambda_d = 1$ and λ_1 . The value $\left| [e_1^{(m)}]_j \right|$ behaves consistent with $\Theta(\sqrt{\lambda_j/\lambda_1})$.

 \Box

Proof. For $a, b > 0$, denote $\alpha(a, b) = \sqrt{\min\{a, b\}} / \max\{a, b\}$. Let $m \in \mathbb{N}$ and assume that Eq. (12) holds which is true if $m = 0$ since $C_0 = 1$. Observe now that $A^{(m+1)} = A^{(m)} + \sqrt{D}[\nu^{(m+1)}][\nu^{(m+1)}]^T \sqrt{D}$. In the s $\sqrt{\min\{a, b\}}/\max\{a, b\}$. Let $m \in \mathbb{N}$ and assume that Eq. (12) holds which is we that $A^{(m+1)} - A^{(m)} + \sqrt{D}[\mathbf{v}^{(m+1)}][\mathbf{v}^{(m+1)}]^T \sqrt{D}$ In the system of coordinates $e_1^{(m)}, \ldots, e_d^{(m)}$
O fan i 5 (1) $d^{(m)}$, *A*^(*m*) writes as *D*^(*m*) := diag($v_1^{(m)}$) $_1$, $,\ldots, r$ (*m*) $\binom{m}{d}$. Since $\lambda_1, \ldots, \lambda_d > 0$, and as $A^{(m)} \ge D$, then $\binom{m}{i} \geq \lambda_i > 0$ for $i \in \{1, ..., d\}$, and

$$
\left\langle \sqrt{D} \nu^{(m+1)}, e_i^{(m)} \right\rangle = \sum_{j=1}^d [e_i^{(m)}]_j \sqrt{\lambda_j} [\nu^{(m+1)}]_j = \sqrt{D_{ii}^{(m)}} \times \sum_{j=1}^d \sqrt{\lambda_j / \nu_i^{(m)}} [e_i^{(m)}]_j [\nu^{(m+1)}]_j =: \left[\sqrt{D^{(m)}} \nu^{(m+1)} \right]_i.
$$

Hence $[A^{(m+1)}]_{\mathcal{B}^{(m)}} = D^{(m)} +$ √ $\overline{D^{(m)}}[w^{(m+1)}]$ $[w^{(m+1)}]^T$ \sqrt{N} *D*(*m*) with

$$
\|w^{(m+1)}\|_{\infty} \leq \sum_{j=1}^d \sqrt{\lambda_j/\nu_i^{(m)}} |[e_i^{(m)}]_j| \times V \leq \sum_{j=1}^d \sqrt{\lambda_j/\lambda_i} |[e_i^{(m)}]_j| \times V \leq dV \times C_m.
$$

We apply Proposition 3 to $\left[A^{(m+1)}\right]_{\mathcal{B}^{(m)}}$ so that, for $i, k \in \{1, ..., d\}$,

$$
|\langle e_i^{(m+1)}, e_k^{(m)} \rangle| = |[[e_i^{(m+1)}]_{\mathcal{B}^{(m)}}]_k| \leqslant 5d^6V^4C_m^4 \alpha(v_i^{(m)}, v_k^{(m)})
$$
.

By Eq. (5), $|\langle e_i^{(m+1)}, e_k^{(m)} \rangle|$ $|\langle k \rangle| \le (5d^6V^4C_m^4)(1+dmV^2)^{1/2} \alpha(\lambda_i, \lambda_k)$. Since $|[e_k^{(m)}\rangle]$ ${k^{(m)} \choose k}$ _{*j*} $| \leq C_m \alpha(\lambda_i, \lambda_k)$, then,

$$
|[e_i^{(m+1)}]_j| \leq \sum_{k=1}^d |[e_k^{(m)}]_j| \times |\langle e_i^{(m+1)}, e_k^{(m)} \rangle| \leq dC_m \times 5d^6V^4C_m^4(1+dmV^2)^{1/2}\alpha(\lambda_i, \lambda_j).
$$

This proves by induction that Eq. (12) holds for all $m \in \mathbb{N}$.

3.3. Thightness

Figure 1 shows numerical computations of coordinates of the first eigenvector of $A^{(m)}$ in dimension 2, 5, 10. The coordinates seem to obey $\Theta(\min\{\lambda_i, \lambda_j\})$ max $\{\lambda_i, \lambda_j\})$ in all cases which suggests that this rate in our upper bounds is tight. However we do not expect the constant C , given in Theorem 4 to be tight. tight. However we do not expect the constant C_m given in Theorem 4 to be tight.

Acknowledgments

The authors would like to thank Stéphane Gaubert for his constructive remarks and feedback on a prior version of the manuscript as well as the anonymous referee for their valuable review and comments.

References

- [1] G. H. Golub, Some Modified Matrix Eigenvalue Problems, SIAM Review 15 (2) (1973) 318–334.
- [2] J. R. Bunch, C. P. Nielsen, D. C. Sorensen, Rank-one modification of the symmetric eigenproblem, Numerische Mathematik 31 (1) (1978) 31–48.
- [3] I. C. F. Ipsen, B. Nadler, Refined Perturbation Bounds for Eigenvalues of Hermitian and Non-Hermitian Matrices, SIAM Journal on Matrix Analysis and Applications 31 (1) (2009) 40–53.
- [4] J. Ding, A. Zhou, Eigenvalues of rank-one updated matrices with some applications, Applied Mathematics Letters 20 (12) (2007) 1223–1226. [5] J. Bénasséni, A. Mom, Inequalities for the eigenvectors associated to extremal eigenvalues in rank one perturbations of symmetric matrices,
- Linear Algebra and its Applications 570 (2019) 123–137. [6] R. C. Thompson, The behavior of eigenvalues and singular values under perturbations of restricted rank, Linear Algebra and its Applications
- 13 (1) (1976) 69–78. [7] R. Mathias, Spectral Perturbation Bounds for Positive Definite Matrices, SIAM Journal on Matrix Analysis and Applications 18 (4) (1997) 959–980.
- [8] R. Bhatia, Perturbation Bounds for Matrix Eigenvalues, no. 53 in Classics in Applied Mathematics, Society for Industrial and Applied Mathematics, Philadelphia, PA, 2007.
- [9] M. Karow, D. Kressner, On a Perturbation Bound for Invariant Subspaces of Matrices, SIAM Journal on Matrix Analysis and Applications 35 (2014) 599–618.
- [10] I. C. F. Ipsen, Relative perturbation results for matrix eigenvalues and singular values, Acta Numerica 7 (1998) 151–201.
- [11] N. Truhar, I. Slapničar, Relative perturbation bound for invariant subspaces of graded indefinite Hermitian matrices, Linear Algebra and its Applications 301 (1) (1999) 171–185.
- [12] I. C. F. Ipsen, An overview of relative sinΘ theorems for invariant subspaces of complex matrices, Journal of Computational and Applied Mathematics 123 (1) (2000) 131–153.
- [13] R. A. Horn, C. R. Johnson, Matrix analysis, Cambridge university press, 2012.
- [14] G. Kjellstrom, L. Taxen, Stochastic optimization in system design, IEEE Transactions on Circuits and Systems 28 (7) (1981) 702–715.
- [15] N. Hansen, S. D. Muller, P. Koumoutsakos, Reducing the Time Complexity of the Derandomized Evolution Strategy with Covariance Matrix ¨ Adaptation (CMA-ES), Evolutionary Computation 11 (1) (2003) 1–18.
- [16] L. Ljung, System Identification. Theory for the User., Information and System Sciences Series, Prentice Hall PTR, Upper Saddle River, NJ 07458, 1999.
- [17] H. Haario, E. Saksman, J. Tamminen, An Adaptive Metropolis Algorithm, Bernoulli 7 (2) (2001) 223–242.
- [18] C. Davis, W. M. Kahan, The Rotation of Eigenvectors by a Perturbation III, SIAM Journal of Numerical Analysis 7 (Mar. 1970).
- [19] N. Hansen, A. Ostermeier, Completely Derandomized Self-Adaptation in Evolution Strategies, Evolutionary Computation 9 (2) (2001) 159– 195.
- [20] S. P. Meyn, R. L. Tweedie, Markov Chains and Stochastic Stability, Springer Science & Business Media, 2012.
- [21] A. Auger, N. Hansen, Linear Convergence of Comparison-based Step-size Adaptive Randomized Search via Stability of Markov Chains, SIAM Journal on Optimization 26 (3) (2016) 1589–1624.
- [22] C. Toure, A. Auger, N. Hansen, Global linear convergence of evolution strategies with recombination on scaling-invariant functions, Journal of Global Optimization 86 (1) (2023) 163–203.
- [23] G. W. Stewart, Matrix Algorithms: Volume II: Eigensystems, SIAM, 2001.