



HAL
open science

The 3-dicritical semi-complete digraphs

Frédéric Havet, Florian Hörsch, Lucas Picasarri-Arrieta

► **To cite this version:**

Frédéric Havet, Florian Hörsch, Lucas Picasarri-Arrieta. The 3-dicritical semi-complete digraphs. 2024. hal-04471658

HAL Id: hal-04471658

<https://inria.hal.science/hal-04471658>

Preprint submitted on 21 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

The 3-dicritical semi-complete digraphs

Frédéric Havet^{*,1}, Florian Hörsch^{†,2}, and Lucas Picasarri-Arrieta^{‡,1}

¹ Université Côte d'Azur, CNRS, Inria, I3S, Sophia-Antipolis, France
²CISPA Saarbrücken, Germany

{frederic.havet, lucas.picasarri-arrieta}@inria.fr
 florian.hoersch@cispa.de

Abstract

A digraph is 3-dicritical if it cannot be vertex-partitioned into two sets inducing acyclic digraphs, but each of its proper subdigraphs can. We give a human-readable proof that the number of 3-dicritical semi-complete digraphs is finite. Further, we give a computer-assisted proof of a full characterization of 3-dicritical semi-complete digraphs. There are eight such digraphs, two of which are tournaments. We finally give a general upper bound on the maximum number of arcs in a 3-dicritical digraph.

1 Introduction

The purpose of this article is to completely characterize all semi-complete 3-dicritical digraphs, hence aiming to approach an analogue of some results on the maximum density of critical graphs. We first recall a few classical definitions and then give an overview of the previous work and motivation for our work.

1.1 Definitions

Our notation follows [4]. In this article, graphs and digraphs contain no parallel edges or arcs, respectively, and no loops. For some positive integer k , we use $[k]$ to denote the set $\{1, \dots, k\}$. The order of a graph G (resp. digraph D) is denoted by $n(G)$ (resp. $n(D)$) and its number of arcs is denoted by $m(G)$ (resp. $m(D)$).

The *path* v_1, \dots, v_n is the graph with vertex set $\{v_i \mid i \in [n]\}$ and edge set $\{v_i v_{i+1} \mid i \in [n-1]\}$. The *length* of a path is its number of edges. A *matching* is a set of pairwise disjoint edges. A graph H is a *subgraph* of a graph G if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$ hold. If additionally $V(H) \neq V(G)$ or $E(H) \neq E(G)$, then H is called a *proper* subgraph of G . For some $S \subseteq V(G)$, we define $G[S]$ to be the subgraph of G *induced by* S , that is, the graph whose vertex set is S and whose edge set contains all the edges of $E(G)$ with both endvertices in S .

Similarly, a digraph D' is a *subdigraph* of a digraph D if $V(D') \subseteq V(D)$ and $A(D') \subseteq A(D)$ hold. If additionally $V(D') \neq V(D)$ or $A(D') \neq A(D)$ holds, then D' is called a *proper* subdigraph of D . If $V(D') = V(D)$, then D is called a *spanning* subdigraph of D . For some $S \subseteq V(D)$, we define $D[S]$ to be the subdigraph of D *induced by* S , that is, the digraph whose vertex set is S and whose arc set contains all the arcs of $A(D)$ with both endvertices in S . When we say that a digraph D contains another digraph D' as an (induced) subdigraph, we mean that D has an (induced) subdigraph which

*Research supported by the research grant DIGRAPHS ANR-19-CE48-0013

†Research supported by the European Research Council (ERC) consolidator grant No. 725978 SYSTEMATIC-GRAPH.

‡Research supported by the research grants DIGRAPHS ANR-19-CE48-0013 and ANR-17-EURE-0004.

is isomorphic to D' , hence not necessarily maintaining vertex labels. In case we want vertex labels to be maintained, we speak of an (induced) *labelled* subgraph. For two disjoint sets $X, Y \subseteq V(D)$, we say that X *dominates* Y (and Y is dominated by X) if $xy \in A(D)$ holds for all $x \in X$ and $y \in Y$.

The *underlying graph* of a digraph D , denoted by $UG(D)$, is the graph on the same vertex set which contains an edge linking two vertices if the digraph contains at least one arc linking these two vertices. Given an arc uv in a digraph, we say that v is an *out-neighbour* of u and that u is an *in-neighbour* of v . The set of out-neighbours of a vertex u in a digraph D is denoted by $N_D^+(u)$, and its set of in-neighbours in D is denoted by $N_D^-(u)$. The out-degree of u in D is its number of out-neighbours, and its in-degree is its number of in-neighbours.

A *digon* is a pair of arcs in opposite directions linking the same vertices. An arc is called *simple* if it is not contained in a digon. A digraph in which every arc is contained in a digon is called *bidirected*. An *oriented graph* is a digraph with no digon. A digraph D is called *semi-complete* if for all distinct $u, v \in V(D)$, at least one of the arcs uv and vu exists. A semi-complete digraph with no digon is called a *tournament*. The *directed cycle* on $n \geq 2$ vertices is the digraph with vertex set $\{v_1, \dots, v_n\}$ and arc set $\{v_i v_{i+1} \mid i \in [n-1]\} \cup \{v_n v_1\}$. The directed cycle on three vertices is called the *directed triangle*. A digraph is *acyclic* if it does not contain any directed cycle. For an acyclic digraph D , there is an ordering v_1, \dots, v_n of $V(D)$ such that for every arc $v_i v_j \in A(D)$, we have $i < j$. Such an ordering is called *acyclic*. An acyclic tournament is called *transitive*. For some positive integer n , we use TT_n to denote the unique transitive tournament on n vertices.

1.2 Context

A k -colouring of a graph G is a function $\phi : V(G) \rightarrow [k]$. It is *proper* if $\phi(u) \neq \phi(v)$ for every edge $uv \in E(G)$. We say that G is k -colourable if it admits a proper k -colouring. Colourability of graphs is one of the most deeply studied subjects in graph theory and countless aspects and variations of this parameter have been considered. One easy observation is that if a graph is k -colourable, then so is each of its subgraphs. This leads to the study of graphs that are in some way minimal obstructions to $(k-1)$ -colourability, which are exactly k -critical graphs: a graph G is k -critical if G is not $(k-1)$ -colourable, but all of its proper subgraphs are. Observe that every k -critical graph is k -colourable. Dirac [8–11] established the basic properties of critical graphs. For $k = 1, 2$, the only k -critical graph is K_k , the complete graph on k vertices, and a graph is 3-critical if and only if it is an odd cycle. For every $k \geq 4$, there is no k -critical graph on $k+1$ vertices and for every $n \geq k+2$, there exists a k -critical graph on n vertices. Further, for every $k \geq 4$, the structure of k -critical graphs is immensely rich and many aspects of k -critical graphs have been studied, concerning both their importance in their own respect and their role when approaching colourability questions.

One feature of particular interest is the density of k -critical graphs. For some $n \geq k$, let $g_k(n)$ be the minimum number of edges of a k -critical graph of order n with the convention $g_k(k+1) = +\infty$. The sequence g_k is rather well understood due to the following result of Kostochka and Yancey [21].

Theorem 1 (Kostochka and Yancey [21]). *Let n and k be two integers such that $n > k \geq 4$.*

If G is a k -critical graph on n vertices, then $m(G) \geq \left\lceil \frac{(k+1)(k-2)n - k(k-3)}{2(k-1)} \right\rceil$. In other words, $g_k(n) \geq \left\lceil \frac{(k+1)(k-2)n - k(k-3)}{2(k-1)} \right\rceil$.

This lower bound is sharp for a significant amount of values of n and k and close to sharp for all remaining ones, which is certified by k -critical graphs that can be obtained from complete graphs using a construction due to Hajós [15]. Theorem 1 confirms a conjecture of Gallai [13] and improves on a collection of earlier results [10, 13, 14, 19, 22]. A more detailed overview can be found in [21].

The *girth* of a graph G is the minimum length of a cycle in G or $+\infty$ if G is a forest. Kostochka and Stiebitz [18] showed that if a k -critical G has girth at least 4, then $m(G) \geq (k - o(k))n(G)$. Informally, this means that the minimum average degree of a k -critical triangle-free graph is (asymptotically) twice

the minimum average degree of a k -critical graph. Liu and Postle [23] showed that the minimum number of edges of a 4-critical graph is significantly larger than $g_4(n)$ if we require the graph to have girth 5. More concretely, they show the existence of constants c and $\varepsilon > 0$ such that every 4-critical graph G of girth at least 5 contains at least $(\frac{5}{3} + \varepsilon)n(G) + c$ edges. Likewise, Postle [27] showed that every 5-critical graph with girth at least 4 satisfies $m(G) \geq (\frac{9}{4} + \varepsilon)n(G) - \frac{5}{4}$ for $\varepsilon = \frac{1}{84}$.

It is also interesting to determine the maximum number of edges in a k -critical graph. Erdős [12] asked, for every fixed $k \geq 4$, whether there exists a constant $c_k > 0$ such that there exists arbitrarily large k -critical graphs G with at least $c_k \cdot n(G)^2$ edges. This was proved by Dirac [8] when $k \geq 6$ and then by Toft [29] when $k \in \{4, 5\}$. The best current upper bound on such a constant c_k has recently been given by Luo, Ma, and Yang [24], improving on an earlier result of Stiebitz [28].

Several analogues of colouring have been introduced for digraphs. In 1982, Neumann-Lara [25] introduced the one of dicolouring. A k -dicolouring of a digraph D is a function $\phi : V(D) \rightarrow [k]$ such that $D[\phi^{-1}(i)]$ is acyclic for every $i \in [k]$. We say that D is k -dicolourable if it admits a k -dicolouring. Dicolourability is a generalization of colouring to digraphs: indeed there is a trivial one-to-one correspondence between the proper k -colourings of a graph G and the k -dicolourings of the associated bidirected graph \overleftrightarrow{G} obtained from G by replacing every edge by a digon.

We say that D is k -dicritical if D is not $(k - 1)$ -dicolourable, but all of its proper subdigraphs are. Observe that every k -dicritical digraph is k -dicolourable. The interest in k -dicritical graphs arises in a similar way as the interest in k -critical graphs. While the only 1-dicritical digraph is the digraph on one vertex and a graph is 2-dicritical if and only if it is a directed cycle, already 3-dicritical digraphs have a very diverse structure. Analogues of Hajós' construction have been found by Bang-Jensen et al. [3]. Again, for some $n \geq k$, it is natural to consider $d_k(n)$, the minimum number of arcs of a k -dicritical digraph of order n , with the convention $d_k(n) = +\infty$ if no such digraph exists.

Observe that, for every $k \geq 2$, the digraph obtained from a bidirected complete graph on $k + 1$ vertices by removing the arcs of a directed triangle is a k -dicritical digraph on $k + 1$ vertices. Note also that a graph G is k -critical if and only if its associated bidirected graph \overleftrightarrow{G} is k -dicritical. Together with the fact that there exist k -critical graphs of every order at least $k + 2$, this implies that there exists a k -dicritical digraph of any order at least k . In other words, we have $d_k(n) < +\infty$ for all $n \geq k$. Moreover, it holds $d_k(n) \leq 2g_k(n)$ for all $n \geq k$.

Kostochka and Stiebitz [20] conjectured that the bidirected k -dicritical digraphs obtained from k -critical graphs are the sparsest k -dicritical digraphs.

Conjecture 2 (Kostochka and Stiebitz [20]). Let n and k be two integers such that $n - 2 \geq k \geq 4$. Then $d_k(n) = 2g_k(n)$ and the k -dicritical digraphs of order n with $d_k(n)$ arcs are the bidirected graphs associated to k -critical graphs with $g_k(n)$ edges.

With Theorem 1 this conjecture implies the following slightly weaker one.

Conjecture 3 (Kostochka and Stiebitz [20]). Let n and k be two integers such that $n > k \geq 4$. If D is a k -dicritical digraph on n vertices, then $m(D) \geq \left\lceil \frac{(k+1)(k-2)n - k(k-3)}{k-1} \right\rceil$ and all digraphs attaining this bound are bidirected.

In [20], Kostochka and Stiebitz confirmed Conjecture 3 for $k \leq 4$.

Similarly to the undirected case, it is expected that the minimum number of arcs in a k -dicritical digraph of order n is larger than $d_k(n)$ if we impose this digraph to have no short directed cycles, and in particular if the digraph is an oriented graph. Let $o_k(n)$ denote the minimum number of arcs in a k -dicritical oriented graph of order n with the convention $o_k(n) = +\infty$ if there is no k -dicritical oriented graph of order n . Clearly, we have $o_k(n) \geq d_k(n)$. Kostochka and Stiebitz [20] posed a conjecture suggesting that there is a significant gap between $d_k(n)$ and $o_k(n)$.

Conjecture 4 (Kostochka and Stiebitz [20]). There exists $\varepsilon > 0$ such that $o_k(n) \geq (1 + \varepsilon) \cdot d_k(n)$ for every $k \geq 4$ and sufficiently large n .

Observe that this conjecture trivially holds when $o_k(n) = +\infty$. However we do not know the set N_k of integers n for which $o_k(n) < +\infty$, that is, for which there exists a k -dicritical oriented graph on n vertices. Already the minimum number n_k of vertices of a k -dicritical oriented graph is unknown except for small values of k : clearly $n_2 = 3$; Neumann Lara [26] proved $n_3 = 7$ and $n_4 = 11$; Bellitto et al. [5] recently established $n_5 = 19$. As observed by Aboulker et al. [2] using a lemma of Hoshino and Kawarabayashi [17], there exists a smallest integer p_k such that there exists a k -dicritical oriented graph on n vertices for any $n \geq p_k$. Moreover, while $p_3 = n_3 = 7$, they showed that $p_4 \neq n_4$ because there is no 4-dicritical oriented graph on 12 vertices.

Conjecture 4 has been confirmed for $k = 3$ by Aboulker et al. [2] and for $k = 4$ by the first and third authors, and Rambaud [16].

We now turn our attention to the maximum density of k -dicritical digraphs which is the main subject of the present article. For every $k \geq 3$, Hoshino and Kawarabayashi [17] constructed an infinite family of k -dicritical oriented graphs D on n vertices which satisfy $m(D) \geq (\frac{1}{2} - \frac{1}{2^{k-1}})n^2$, and they conjectured that this bound is tight.

Conjecture 5 (Hoshino and Kawarabayashi [17]). Let $k \geq 3$ be an integer. If D is a k -dicritical oriented graph, then $m(D) \leq (\frac{1}{2} - \frac{1}{2^{k-1}})n^2$.

Aboulker [1] observed that, since a tournament has $\frac{1}{2}n(n-1)$ arcs, this conjecture implies that the number of k -dicritical tournaments is finite, and he asked whether this latter statement holds. It trivially does for the case $k = 2$.

1.3 Our results

In this paper, we positively answer Aboulker's question in the case $k = 3$ by showing that the collection of 3-dicritical semi-complete digraphs is finite, and hence so is the subcollection of 3-dicritical tournaments.

Theorem 6. *There is a finite number of 3-dicritical semi-complete digraphs.*

While the proof of Theorem 6 is fully human readable, the result is obtained by showing that the number of vertices of any 3-dicritical semi-complete digraph does not exceed a pretty large number which originates from a Ramsey-type argument.

We after use a computer-assisted proof to provide the following characterization of all 3-dicritical semi-complete digraphs.

Theorem 7. *There are exactly eight 3-dicritical semi-complete digraphs. They are depicted in Figure 1.*

In particular, we can characterize all 3-dicritical tournaments.

Corollary 8. *There are exactly two 3-dicritical tournaments, namely $\mathcal{R}(\overrightarrow{C}_3, \overrightarrow{C}_3)$ and \mathcal{P}_7 .*

We finally investigate the maximum density of 3-dicritical digraphs. The *bidirected part* of a digraph D is the graph $B(D)$ with vertex set $V(D)$ in which two vertices are linked by an edge if and only if there is a digon between them in D . We prove in Proposition 25 that $B(D)$ is a forest for every 3-dicritical digraph D . From this result, one can easily deduce that $m(D) \leq \binom{n}{2} + n - 1$ holds for every 3-dicritical digraph D different from \overleftrightarrow{K}_3 . We slightly improve this upper bound on $m(D)$ as follows (the digraph \overrightarrow{W}_3 is depicted in Figure 1).

Theorem 9. *If D is a 3-dicritical digraph distinct from \overleftrightarrow{K}_3 and \overrightarrow{W}_3 , then $m(D) \leq \binom{n}{2} + \frac{2}{3}n$.*




Figure 1: The 3-dicritical semi-complete digraphs, namely the bidirected complete graph \overleftrightarrow{K}_3 , the directed wheel \overrightarrow{W}_3 , the digraph \mathcal{H}_5 , the rotative digraphs $\mathcal{R}(H_1, H_2)$ for every $H_1, H_2 \in \{\overleftrightarrow{K}_2, \overrightarrow{C}_3\}$ and the Paley tournament on seven vertices \mathcal{P}_7 . A big arrow linking two sets of vertices indicates that there is exactly one arc from every vertex in the first set to every vertex in the second set.

The rest of this article is structured as follows: In Section 2, we give a collection of preliminary results which will be used in the proofs of Theorems 6, 7, and 9. In Section 3, we prove Theorem 6. In Section 4, we prove Theorem 7, with all code we use being shifted to the appendix. Section 5 is devoted to the proof of Theorem 9. Finally, in Section 6, we conclude our work and give some directions for further research.

2 Useful lemmas

In this section, we give a collection of preliminary results we need in the proof of Theorem 6. Most of them will be reused in the proof of Theorem 7. We first describe 2-dicolourings with some important extra properties.

Let D be a digraph and uv be an arc of D . A uv -colouring of D is a 2-colouring $\phi : V(D) \rightarrow [2]$ such that:

- ϕ is a 2-dicolouring of $D \setminus uv$,

- $\phi(u) = \phi(v) = 1$, and
- $D \setminus uv$, coloured with ϕ , does not contain any monochromatic directed uv -path.

There is a close relationship between 3-dicritical digraphs and uv -colourings.

Lemma 10. *Let D be a 3-dicritical digraph and uv be an arc of D . Then D admits a uv -colouring.*

Proof. As D is 3-dicritical, there is a 2-dicolouring $\phi : V(D) \rightarrow [2]$ of $D \setminus uv$. By symmetry, we may suppose $\phi(u) = 1$. As ϕ is not a 2-dicolouring of D , we obtain that $\phi(v) = 1$ and there is a directed vu -path P in D such that $\phi(x) = 1$ for all $x \in V(P)$. If there is also a directed uv -path Q in $D \setminus uv$ such that $\phi(x) = 1$ for all $x \in V(Q)$, then the subdigraph of $D \setminus uv$ induced by $V(P) \cup V(Q)$ contains a monochromatic directed cycle. This contradicts ϕ being a 2-dicolouring of $D \setminus uv$. \square

The next result showing that every arc of a 3-dicritical semi-complete digraph is contained in a short directed cycle will play a crucial role in the upcoming proofs.

Lemma 11. *Let D be a 3-dicritical semi-complete digraph. Then every arc $a \in A(D)$ either belongs to a digon or is contained in an induced directed triangle.*

Proof. As D is 3-dicritical, there is a 2-dicolouring ϕ of $D \setminus a$. As ϕ is not a 2-dicolouring of D , there exists a directed cycle C in D such that C is monochromatic with respect to ϕ . We may suppose that C is chosen to be of minimum length with this property. As D is semi-complete, we obtain that C is either a digon or an induced directed triangle. As C is not a monochromatic directed cycle of $D \setminus a$ with respect to ϕ , we obtain that $a \in A(C)$. \square

We define O_5 as the oriented graph which consists of a directed triangle xyz and two additional vertices u, v , one arc from u to every vertex of the directed triangle, one arc from every vertex of the directed triangle to v , and the arc uv . An illustration can be found in Figure 2.




Figure 2: The oriented graph O_5 .

The following result is a consequence of Lemma 10.

Lemma 12. *Let D be a 3-dicritical digraph. Then D does not contain O_5 as a subdigraph.*

Proof. Assume for a contradiction that D contains O_5 as a subdigraph and let $V(O_5) = \{u, v, x, y, z\}$ be the labelling depicted in Figure 2. By Lemma 10, there exists a uv -colouring ϕ of D . Since there exists no monochromatic directed uv -path, we have $\phi(x) = \phi(y) = \phi(z) = 2$. Hence $D \setminus uv$ contains a monochromatic directed triangle with respect to ϕ , a contradiction. \square

Let S be a transitive subtournament of a digraph $D = (V, A)$. We denote by v_1, \dots, v_s the unique acyclic ordering of S . For some $i, j \in [s]$, we say that $\{v_i, \dots, v_j\}$ is an *interval* of S . Observe that \emptyset is an interval. For $i_0, j_0, i_1, j_1 \in [s]$ with $j_0 < i_1$, we say that the interval $\{v_{i_0}, \dots, v_{j_0}\}$ is *smaller* than the interval $\{v_{i_1}, \dots, v_{j_1}\}$. By convention \emptyset is both smaller and greater than any other interval. A sequence of intervals P_1, \dots, P_t is called *increasing* if P_i is smaller than P_j for all $i, j \in [t]$ with $i < j$.

Lemma 13. *Let T be a subtournament of a 3-dicritical digraph D and let S be a transitive subtournament of T with acyclic ordering v_1, \dots, v_s . For any $x \in V(T) \setminus S$, there is an increasing sequence of intervals (I_1, I_2, I_3, I_4) with $\bigcup_{i=1}^4 I_i = S$ such that, in T , x dominates $I_1 \cup I_3$ and is dominated by $I_2 \cup I_4$.*

Proof. Assume this is not the case. Then there exists an increasing sequence of indices (i_1, i_2, i_3, i_4) such that, in T , x is dominated by v_{i_1} and v_{i_3} and dominates v_{i_2} and v_{i_4} . Then the subdigraph of D induced by $\{v_{i_1}, v_{i_2}, x, v_{i_3}, v_{i_4}\}$ contains O_5 as a subdigraph, a contradiction to Lemma 12. \square

We finally need a well-known theorem which can be found in many basic textbooks on graph theory, see for example [7, Theorem 9.1.3].

Theorem 14 (MULTI-COLOUR RAMSEY THEOREM). *Let a and b be positive integers. There exists a smallest integer $R_a(b)$ such that for G being a copy of $K_{R_a(b)}$ and for every mapping $\psi : E(G) \rightarrow [a]$, there is a set $S \subseteq V(G)$ of cardinality b and $i \in [a]$ such that $\psi(e) = i$ for all $e \in E(G[S])$.*

3 A simple proof for finiteness

In this section, we prove that the number of 3-dicritical semi-complete digraphs is finite. Let us first restate this result.

Theorem 6. *There is a finite number of 3-dicritical semi-complete digraphs.*

Proof. Let $D = (V, A)$ be a 3-dicritical semi-complete digraph. We will show that $n(D) \leq 12R_6(3) + 1$, where $R_6(3)$ refers to the Ramsey number in Theorem 14. Assume for the sake of a contradiction that $n(D) \geq 12R_6(3) + 2$. Let $S \subseteq V$ be a maximum set of vertices such that $D[S]$ is acyclic. Let v_1, \dots, v_s be the unique acyclic ordering of S . Since D is 3-dicritical, for an arbitrary vertex $x \in V$, we have that $D - x$ is 2-dicolourable. This yields $s \geq \left\lceil \frac{n(D)-1}{2} \right\rceil \geq 6R_6(3) + 1$.

By Lemma 11, for every $i \in [s - 1]$, the arc $v_i v_{i+1}$ belongs to a digon or an induced directed triangle. Therefore, since $D[S]$ is acyclic, we know that there exists a vertex $x_i \in V \setminus S$ such that $v_i v_{i+1} x_i v_i$ is an induced directed triangle C_i .

Let T be an arbitrary spanning subtournament of D . Observe that $T[S] = D[S]$ as $D[S]$ is acyclic. Further, the directed triangle C_i is contained in T for $i \in [s - 1]$ as C_i is induced in D . For any vertex x in $V \setminus S$ and $i \in [s - 1]$, we say that x *switches at i* if x dominates v_i and is dominated by v_{i+1} in T or x is dominated by v_i and dominates v_{i+1} in T .

Let H be the digraph with vertex set $V(H) = [s - 1]$ and arc set $A(H) = A_1 \cup A_2$ with $A_1 = \{(i, i + 1) \mid i \in [s - 2]\}$ and $A_2 = \{(i, j) \mid i \neq j \text{ and } x_i \text{ switches at } j.\}$

By Lemma 13, for $i \in [s - 1]$, we have that x_i switches at at most three indices in $[s - 1]$. Further, as C_i is a directed triangle, x_i switches at i which yields that x_i switches at at most two indices in $[s - 1] \setminus \{i\}$. Thus every $i \in [s - 1]$ is the tail of at most two arcs in A_2 .

For every subset J of $[s - 1]$, observe that $H[J]$ contains at most $|J| - 1$ arcs in A_1 and at most $2|J|$ arcs in A_2 , hence at most $3|J| - 1$ arcs in total. Thus $\text{UG}(H)[J]$ has a vertex of degree at most 5. Hence $\text{UG}(H)$ is 5-degenerate, and so it is 6-colourable. Therefore H has an independent set I of size $\left\lceil \frac{1}{6}(s - 1) \right\rceil \geq R_6(3)$.

By definition of I , for any $i, j \in I$ with $i \neq j$, we have that $V(C_i)$ and $V(C_j)$ are disjoint. Moreover, either $\{v_j, v_{j+1}\}$ dominates x_i in T or $\{v_j, v_{j+1}\}$ is dominated by x_i in T . Hence, if $i < j$, the subdigraph of T induced by $V(C_i) \cup V(C_j)$ is one of the eight tournaments depicted in Figure 3. For $(\alpha) \in \{(a), \dots, (h)\}$, we say that (i, j) is an (α) -*configuration* if $T[V(C_i) \cup V(C_j)]$ is the tournament depicted in Figure 3 (α) .

Let us fix a pair $i, j \in I$ with $i < j$. We know that it is not a (g) -configuration, for otherwise $D[v_{i+1}, v_j, v_{j+1}, x_j, x_i]$ contains O_5 as a subdigraph, a contradiction to Lemma 12. We also know




Figure 3: A listing of all possible configurations for $i, j \in I$ with $i < j$. For the sake of better readability, the arcs in $A(C_i) \cup A(C_j)$ and the arcs from $V(C_j)$ to $V(C_i)$ are solid, and the arcs from $V(C_i)$ to $V(C_j)$ are dashed.

that it is not an (h) -configuration, for otherwise $D[x_j, v_i, v_{i+1}, x_i, v_j]$ contains O_5 as a subdigraph, a contradiction to Lemma 12.

Since $|I| \geq R_6(3)$, and by definition of $R_6(3)$, we know that there exist $\{i, j, h\} \subseteq I$, $i < j < h$, and $(\alpha) \in \{(a), \dots, (e)\}$ such that the three pairs $(i, j), (j, h), (i, h)$ are (α) -configurations. We show that each of the six cases yields a contradiction, implying the result.

- If $(\alpha) = (a)$, let ϕ be a $v_{i+1}v_h$ -colouring of D , the existence of which is guaranteed by Lemma 10. Recall that $\phi(v_{i+1}) = \phi(v_h) = 1$, ϕ is a 2-dicolouring of $D \setminus v_{i+1}v_h$ and D coloured with ϕ contains no monochromatic directed $v_{i+1}v_h$ -path. Then $\phi(v_j) = \phi(v_{j+1}) = 2$ because $\{v_j, v_{j+1}\} \subseteq N_D^+(v_{i+1}) \cap N_D^-(v_h)$. Thus, since C_j is not monochromatic in ϕ , we have $\phi(x_j) = 1$. We obtain that $\phi(x_i) = \phi(v_i) = 2$, for otherwise $v_{i+1}x_ix_jv_{i+1}$ or $v_iv_hx_jv_i$ is monochromatic. We deduce that $v_iv_jx_iv_i$ is monochromatic, a contradiction.
- If $(\alpha) = (b)$, then $D[x_h, v_j, v_{j+1}, x_j, x_i]$ contains O_5 as a subdigraph, a contradiction to Lemma 12.
- If $(\alpha) = (c)$, then $D[x_i, v_j, v_{j+1}, x_j, v_h]$ contains O_5 as a subdigraph, a contradiction to Lemma 12.

- If $(\alpha) = (d)$, then $D[v_i, v_j, v_{j+1}, x_j, x_h]$ contains O_5 as a subdigraph, a contradiction to Lemma 12.
- If $(\alpha) = (e)$, then $D[v_i, v_j, v_{j+1}, x_j, v_h]$ contains O_5 as a subdigraph, a contradiction to Lemma 12.
- If $(\alpha) = (f)$, then $D[v_i, v_j, v_{j+1}, x_j, v_h]$ contains O_5 as a subdigraph, a contradiction to Lemma 12.

□

4 The 3-dicritical semi-complete digraphs

This section is devoted to a computer-assisted proof of Theorem 7. It follows a similar line as the one of Theorem 6, but it needs some refined arguments. Further, due to the significant number of necessary computations, several parts of the proof are computer-assisted. We used codes implemented using SageMath. They are available on the third author's GitHub page.

We first restrain the structure of 3-dicritical semi-complete digraphs. To prove Theorem 6, we only needed the fact that O_5 does not occur as a subdigraph. To prove Theorem 7, we need to prove that several other digraphs cannot be subdigraphs or induced subdigraphs of a 3-dicritical digraph. One of these digraphs is the transitive tournament of size at least 8. While already parts of this proof are computer-assisted, the most intense computation part is carried out after. We generate all semi-complete digraphs satisfying these properties and check that none of them has dichromatic number 3, except the ones depicted in Figure 1.

Before dealing with the collection of digraphs which are not contained in 3-dicritical semi-complete digraphs as subdigraphs, we first give the following simple observation on matchings in graphs on seven vertices which will prove useful later on.

Lemma 15. *Let H be a graph that is obtained from a path $w_1 \dots w_7$ by adding the edges of a matching M on $\{w_1, \dots, w_7\}$. Then there is a stable set $S \subseteq V(H)$ with $|S| = 3$ and $\{w_1, w_7\} \setminus S \neq \emptyset$.*

Proof. Suppose otherwise. If w_1 is not incident to an edge of M , then, as none of $\{w_1, w_3, w_5\}$ and $\{w_1, w_3, w_6\}$ is an independent set, we obtain that $w_3w_5, w_3w_6 \in E(M)$, a contradiction to M being a matching. Hence M contains an edge e_1 incident to w_1 . Similarly, M contains an edge e_7 incident to w_7 . Further, M contains an edge e_0 both of whose endvertices are contained in $\{w_2, w_4, w_6\}$. As none of e_0 and e_7 are contained in one of $\{w_1, w_3, w_6\}$ and $\{w_1, w_3, w_5\}$, we obtain that e_1 needs to be contained in both of them. This yields $e_1 = w_1w_3$. Similarly, we obtain $e_7 = w_5w_7$. By symmetry, we may suppose that $e_0 \neq w_4w_6$. But then $\{w_1, w_4, w_6\}$ is an independent set, a contradiction. □

We now start excluding some subdigraphs of 3-dicritical semi-complete digraphs. We first define O_4 as the digraph which consists of a copy of \overleftrightarrow{K}_2 and two additional vertices u, v , one arc from u to every vertex of \overleftrightarrow{K}_2 , one arc from every vertex of \overleftrightarrow{K}_2 to v , and the arc uv . An illustration can be found in Figure 4.

The digraph O_4 plays a similar role as O_5 . Also, the proof of the following result is similar to the one of Lemma 12.

Lemma 16. *Let D be a 3-dicritical digraph. Then D does not contain O_4 as a subdigraph.*

Proof. Assume for the purpose of contradiction that D contains O_4 as a subdigraph and let $V(O_4) = \{u, v, x, y\}$ be the labelling depicted in Figure 4. By Lemma 10, there exists a 2-dicolouring ϕ of $D \setminus uv$ with $\phi(x) = \phi(y)$. Hence $D \setminus uv$ contains a monochromatic digon with respect to ϕ , a contradiction. □

In the following, let \overleftrightarrow{S}_4 be the bidirected star on 4 vertices, see Figure 5. The following result shows that \overleftrightarrow{S}_4 cannot be the subdigraph of any large 3-dicritical semi-complete digraph.




Figure 4: The digraph O_4 .




Figure 5: The bidirected star on 4 vertices \overleftrightarrow{S}_4 .

Lemma 17. *Let D be a semi-complete digraph containing \overleftrightarrow{S}_4 as a subdigraph. Then D is 3-dicritical if and only if D is \overrightarrow{W}_3 .*

Proof. It is easy to see that \overrightarrow{W}_3 is 3-dicritical and contains \overleftrightarrow{S}_4 . For the other direction, let D be a 3-dicritical semi-complete digraph such that D contains a vertex u linked by digons to three distinct vertices x, y, z .

Then, as D is semi-complete, we have that $D[\{x, y, z\}]$ needs to contain \overrightarrow{C}_3 or TT_3 as a subdigraph. If it is TT_3 , then D contains O_4 as a subdigraph, a contradiction to Lemma 16. Hence $D[\{u, x, y, z\}]$ contains \overrightarrow{W}_3 as a subdigraph. Since both D and \overrightarrow{W}_3 are 3-dicritical, we have $D = \overrightarrow{W}_3$. \square

We now prove a similar result for a collection of four digraphs. Given two digraphs H_1 and H_2 , let $H_1 \Rightarrow H_2$ denote the *directed join* of H_1 and H_2 , that is the digraph obtained from disjoint copies of H_1 and H_2 by adding all arcs from the copy of H_1 to the copy of H_2 . If we further add all the arcs from H_2 to H_1 , we obtain the *bidirected join* of H_1 and H_2 , denoted by $H_1 \boxplus H_2$. It is straightforward that $\vec{\chi}(H_1 \boxplus H_2) = \vec{\chi}(H_1) + \vec{\chi}(H_2)$, see [3].

Lemma 18. *Let H_1, H_2 be two digraphs in $\{\overrightarrow{K}_2, \overrightarrow{C}_3\}$ and let D be a semi-complete digraph containing $H_1 \Rightarrow H_2$ as a subdigraph. Then D is 3-dicritical if and only if D is exactly $\mathcal{R}(H_1, H_2)$.*

Proof. It is easy to see that $\mathcal{R}(H_1, H_2)$ is 3-dicritical. For the other direction, let us fix $H_1, H_2 \in \{\overrightarrow{K}_2, \overrightarrow{C}_3\}$ and let D be a 3-dicritical semi-complete digraph containing $H_1 \Rightarrow H_2$.

Let $X = V(H_1)$ and $Y = V(H_2)$. Let us first prove that $V(D) \setminus (X \cup Y) \neq \emptyset$, so assume for a contradiction that $V(D) = X \cup Y$. We claim that there exists a simple arc uv from X to Y . If this is not the case, then D is exactly $H_1 \boxplus H_2$, so it has dichromatic number 4, a contradiction. This simple arc uv belongs to an induced directed triangle by Lemma 11. This directed triangle uses an arc from Y to X , which is necessarily in a digon, a contradiction since it must be induced. Henceforth we assume that $V(D) \setminus (X \cup Y) \neq \emptyset$.

First suppose that there exists some $v \in V(D) \setminus (X \cup Y)$ having at least one in-neighbour and one out-neighbour in both X and Y . Since H_1 and H_2 are strongly connected, there exist four distinct vertices $x_1, x_2 \in X$ and $y_1, y_2 \in Y$ such that $\{x_1x_2, y_1y_2, x_1v, vx_2, y_1v, vy_2\}$ are all arcs of D . Then $D[\{x_1, v, x_2, y_1, y_2\}]$ contains O_5 as a subdigraph, a contradiction to Lemma 12. Henceforth we may assume that every vertex $v \in V(D) \setminus (X \cup Y)$ has no out-neighbour or no in-neighbour in one of $\{X, Y\}$.

Now suppose that there exists some $v \in V(T) \setminus (X \cup Y)$ that dominates X . If v has an out-neighbour y in Y , then $D[X \cup \{v, y\}]$ contains O_4 as a subdigraph if $H_1 = \overleftarrow{K}_2$ and O_5 otherwise, a contradiction to Lemma 16 or 12, respectively. Hence v has no out-neighbour in Y . Since D is semi-complete, this implies that Y dominates v . Hence D contains $\mathcal{R}(H_1, H_2)$, implying that D is exactly $\mathcal{R}(H_1, H_2)$ since both D and $\mathcal{R}(H_1, H_2)$ are 3-dicritical.

Henceforth we assume that for every vertex $v \in V(D) \setminus (X \cup Y)$, there exists in D a simple arc from X to v . By directional duality, there exists also a simple arc from v to Y . Recall that every vertex $v \in V(D) \setminus (X \cup Y)$ has no out-neighbour or no in-neighbour in one of $\{X, Y\}$. We conclude on the existence of a partition (V_1, V_2) of $V(D) \setminus (X \cup Y)$ such that there is no arc from V_1 to X and there is no arc from Y to V_2 .

By symmetry, we may assume that V_2 is non-empty. Let us fix $v_2 \in V_2$ and $y_1 \in Y$. Since $v_2 y_1$ is a simple arc, by Lemma 11, there exists a vertex v_1 such that $v_2 y_1 v_1 v_2$ is an induced directed triangle in D . Note that $v_1 \notin Y$ since there is no arc from Y to V_2 . Also note that $v_1 \notin X$ for otherwise $v_2 y_1 v_1 v_2$ is not induced since X dominates Y . Further note that $v_1 \notin V_2$ since it is an out-neighbour of y_1 . This implies $v_1 \in V_1$. As v_2 does not dominate X , there is some vertex in X , say x_1 that dominates v_2 . Note that x_1 dominates v_1 by definition of V_1 . Let y_2 be the unique out-neighbour of y_1 in Y .

If v_1 dominates y_2 , we obtain that $D[\{x_1, v_1, v_2, y_1, y_2\}]$ contains O_5 as a subdigraph, a contradiction to Lemma 12. We may hence suppose that y_2 dominates v_1 . As Y does not dominate v_1 , this implies that H_2 is \overrightarrow{C}_3 and the out-neighbour y_3 of y_2 is dominated by v_1 . Then $D[\{x_1, v_1, v_2, y_2, y_3\}]$ contains O_5 as a subdigraph, a contradiction to Lemma 12. \square

The rest of the preparatory results before the main proof of Theorem 7 aims to exclude a collection of tournaments \mathcal{T}_8 as induced subdigraphs and another digraph F as a (not necessarily induced) subdigraph. As the proofs of these results contain several common preliminaries, we give them together. While the exact definition of \mathcal{T}_8 is postponed, we now give the definition of F . Let F be the oriented graph with vertex set $\{u_1, \dots, u_6, x_1, x_2, x_3\}$ such that:

- $\{u_1, \dots, u_6\}$ induces a copy of TT_6 the unique acyclic ordering of which is exactly u_1, \dots, u_6 , and
- for every $i \in [3]$, F contains the arcs $u_{2i}x_i$ and $x_i u_{2i-1}$.

See Figure 6 for an illustration of F .




Figure 6: The oriented graph F .

We let \mathcal{F} be the set of tournaments T with vertex set $V(T) = V(F)$ and such that $A(F) \subseteq A(T)$. Note that \mathcal{F} contains 2^{15} tournaments since F has exactly 15 pairs of non-adjacent vertices. Four of them are of special interest and we denote them by T^1, \dots, T^4 . We give their adjacency matrices in Appendix A.

Lemma 19. *None of the tournaments in $\mathcal{F} \setminus \{T^1, T^2, T^3, T^4\}$ is a subdigraph of a 3-dicritical semi-complete digraph.*

Proof. For every tournament $T \in \mathcal{F}$, we check, using the code of Appendix B.1, if it contains $\overrightarrow{C_3} \Rightarrow \overrightarrow{C_3}$ as a subdigraph or if it admits no uv -colouring for an arc uv . This is always the case except when $T \in \{T^1, T^2, T^3, T^4\}$. The claim then follows by Lemmas 10 and 18. \square

Let F^+ be the oriented graph obtained from F by adding a vertex u_0 and the arcs of $\{u_0u_i \mid i \in [6]\}$. Analogously, let F^- be the oriented graph obtained from F by adding a vertex u_7 and the arcs of $\{u_iu_7 \mid i \in [6]\}$. See Figure 7 for an illustration.




Figure 7: The oriented graphs F^+ (left) and F^- (right).

Lemma 20. *Let D be a 3-dicritical semi-complete digraph. Then D does not contain a digraph in $\{F^+, F^-\}$ as a subdigraph.*

Proof. Observe that the digraph obtained from F^- by reversing all its arcs is isomorphic to F^+ . As the digraph obtained from a 3-dicritical, semi-complete digraph by reversing all arcs is 3-dicritical and semi-complete, it suffices to prove the statement for F^+ .

In order to do so, suppose for the sake of a contradiction, that there is a 3-dicritical semi-complete digraph D containing F^+ . By Lemma 19, $D - u_0$ contains some $T' \in \{T^1, T^2, T^3, T^4\}$. Now consider the collection \mathcal{T} of tournaments on $\{u_0, \dots, u_6, v_1, v_2, v_3\}$ that have one of T^1, T^2, T^3, T^4 as a labelled subdigraph and in which u_0 dominates $\{u_1, \dots, u_6\}$. Observe that by assumption, D contains a tournament in \mathcal{T} as a spanning subdigraph. Further, \mathcal{T} contains exactly $4 \times 2^3 = 32$ digraphs. Using the code in Appendix B.2, we check that each of them contains $\overrightarrow{C_3} \Rightarrow \overrightarrow{C_3}$ or contains an arc uv with no uv -colouring. We conclude that the same holds for D , a contradiction to Lemmas 10 or 18. \square

We are now ready to show that 3-dicritical semi-complete digraphs do not contain large transitive tournaments as induced subdigraphs.

Lemma 21. *Let D be a 3-dicritical semi-complete digraph. Then D does not contain TT_8 as an induced subdigraph.*

Proof. For a contradiction, assume that $D = (V, A)$ is a 3-dicritical semi-complete digraph containing TT_8 as an induced subdigraph. We will prove that D contains F^+ or F^- , which is a contradiction to Lemma 20.

Let $S \subseteq V$ be such that $D[S]$ is isomorphic to TT_8 . Let v_1, \dots, v_8 be the unique acyclic ordering of S . By Lemma 11, for every $i \in [7]$, there exists a vertex $x_i \in V \setminus S$ such that $v_i v_{i+1} x_i v_i$ forms an induced directed triangle C_i .

Let H be the graph with vertex set $V(H) = [7]$ and that contains an edge linking i and j if $V(C_i) \cap V(C_j) \neq \emptyset$. For any $i, j \in [7]$ with $ij \in E(H)$ and $|i - j| \geq 2$, we have $x_i = x_j$. By Lemma 13, there is no set $\{i, j, k\} \subseteq [7]$ such that $x_i = x_j = x_k$. This yields that H is obtained from a path on 7 vertices by adding a matching. We deduce from Lemma 15 that there is a set $I \subseteq [7]$ with $|I| = 3$ such that the following hold:

- (a) $\{1, 7\} \setminus I \neq \emptyset$,

(b) C_i and C_j are vertex-disjoint for all $\{i, j\} \subseteq I$.

This shows that D contains F^+ or F^- , yielding a contradiction to Lemma 20. \square

Given an integer k and a semi-complete digraph D , a k -extension of D is a semi-complete digraph on $n(D) + k$ vertices containing D as an induced subdigraph. Given a set S of semi-complete digraphs, a k -extension of S is a semi-complete digraph that is a k -extension of some $D \in S$. We are now ready to prove that no 3-dicritical semi-complete digraph contains F as a subdigraph.

Lemma 22. *Let D be a 3-dicritical semi-complete digraph. Then D does not contain F as a subdigraph.*

Proof. By Lemma 19, it remains to show that D does not contain a graph in $\{T^1, T^2, T^3, T^4\}$ as a subdigraph. Assume for a contradiction that D contains at least one of $\{T^1, T^2, T^3, T^4\}$ as a subtournament.

We use the code in Appendix B.3. In a first part, we compute the set \mathcal{L} of all semi-complete digraphs L on nine vertices such that each of the following holds:

- (i) L contains some $T \in \{T^1, T^2, T^3, T^4\}$ as a subdigraph,
- (ii) L does not contain any digraph in $\{\overrightarrow{S}_4, \overleftarrow{K}_2 \Rightarrow \overleftarrow{K}_2, \overleftarrow{K}_2 \Rightarrow \overleftarrow{C}_3, \overleftarrow{C}_3 \Rightarrow \overleftarrow{K}_2, \overleftarrow{C}_3 \Rightarrow \overleftarrow{C}_3, O_4, O_5\}$ as a subdigraph,
- (iii) L admits a uv -colouring for every arc $uv \in A(L)$, and
- (iv) L does not contain TT_8 as an induced subdigraph.

By Lemmas 10, 12, 16, 17, and 18, we know that D contains some $L \in \mathcal{L}$ as an induced subdigraph. In the second part of the code, we check that every 2-extension L' of \mathcal{L} does not satisfy at least one of the properties (ii), (iii) and (iv).

This shows, by Lemmas 10, 12, 16 and 18, that either $D \in \mathcal{L}$ or D is a 1-extension of \mathcal{L} . Finally, we check that every $L \in \mathcal{L}$ has dichromatic number at most two, and that every 1-extension L' satisfying (ii), (iii) and (iv) has dichromatic number at most two. This yields a contradiction. \square

We now give the definition of \mathcal{T}_8 and show that no digraph in \mathcal{T}_8 can be contained in a 3-dicritical semi-complete digraph as an induced subdigraph. Let \mathcal{T}_8 be the set of tournaments obtained from TT_8 by reversing exactly one arc. Observe that TT_8 belongs to \mathcal{T}_8 .

Lemma 23. *Let D be a 3-dicritical semi-complete digraph. Then D does not contain any digraph in \mathcal{T}_8 as an induced subdigraph.*

Proof. Assume for a contradiction that D contains some $T' \in \mathcal{T}_8$ as an induced subtournament. Let $X \subseteq V(T')$ be such that $D[X]$ is isomorphic to T' . By definition of \mathcal{T}_8 , let x_1, \dots, x_8 be an ordering of X such that D contains every arc $x_i x_j$ when $i < j$, except for exactly one pair $\{k, \ell\}$, $k < \ell$.

Assume first that $k = \ell - 1$. Then observe that T' is isomorphic to TT_8 , with the acyclic ordering obtained from x_1, \dots, x_8 by swapping x_ℓ and $x_{\ell-1}$. This contradicts Lemma 21.

Henceforth assume that $k \leq \ell - 2$. If $k \geq 2$ and $\ell \leq 7$ then $D[\{x_1, x_k, x_{\ell-1}, x_\ell, x_8\}]$ is isomorphic to O_5 , a contradiction to Lemma 12. Henceforth we assume that $k = 1$ or $\ell = 8$. By directional duality, we assume without loss of generality that $k = 1$. Let S be the transitive induced subtournament of D on vertices $X \setminus \{x_1, x_2, x_\ell\}$. We denote its acyclic ordering by y_1, \dots, y_5 , which exactly corresponds to $x_3, \dots, x_{\ell-1}, x_{\ell+1}, \dots, x_8$. By Lemma 11, for every $k \in [4]$, there exists a vertex z_k such that $y_k y_{k+1} z_k y_k$ forms a directed triangle C_k . As S is induced, z_k must be in $V \setminus V(S)$. Moreover, $z_k \notin \{x_1, x_2, x_\ell\}$ because both $X \setminus \{x_1\}$ and $X \setminus \{x_\ell\}$ are acyclic.

Let H be the graph with vertex set $V(H) = [4]$ and that contains an edge linking i and j if $V(C_i) \cap V(C_j) \neq \emptyset$. For any $i, j \in [4]$ with $ij \in E(H)$ and $|i - j| \geq 2$, we have $z_i = z_j$. By Lemma 13,

there is no set $\{h, i, j\} \subseteq [4]$ such that $z_i = z_j = z_h$. This yields that H is obtained from a path on 4 vertices by adding a matching containing at most 2 edges. Hence H contains two non-adjacent vertices, corresponding to two disjoint directed triangles C_i and C_j in D . Together with the directed cycle $C_h = x_1x_2x_\ell$, we deduce that D contains F as a subdigraph. This contradicts Lemma 22. \square

We have now proved all necessary structural properties of 3-dicritical semi-complete digraphs. The following result contains the decisive step of the proof and it requires heavy computation. For every $i \in [7]$, let \mathcal{D}_i be the set of semi-complete digraphs D such that each of the following holds:

- the maximum acyclic set $S \subseteq V(D)$ of D has size exactly i ,
- for every arc uv of D , D admits a uv -colouring,
- D does not contain any digraph of $\{\overleftarrow{S}_4, \overleftarrow{K}_2 \Rightarrow \overleftarrow{K}_2, \overleftarrow{K}_2 \Rightarrow \overrightarrow{C}_3, \overrightarrow{C}_3 \Rightarrow \overleftarrow{K}_2, \overrightarrow{C}_3 \Rightarrow \overrightarrow{C}_3, O_4, O_5, F\}$ as a subdigraph,
- D does not contain any digraph of \mathcal{T}_8 as an induced subdigraph,

Lemma 24. *The 3-dicritical digraphs in $\bigcup_{i=1}^7 \mathcal{D}_i$ are exactly \overleftarrow{K}_3 , \mathcal{H}_5 , and \mathcal{P}_7 .*

Proof. For every $i \in [7]$, we compute \mathcal{D}_i by starting from the singleton $\{TT_i\}$ which is clearly the only digraph in \mathcal{D}_i on at most i vertices. Using the code in Appendix B.4, we first successively compute the digraphs in \mathcal{D}_i on $j \geq i$ vertices by generating every possible 1-extension of the digraphs in \mathcal{D}_i on $j-1$ vertices, and saving only the ones satisfying the conditions on \mathcal{D}_i . When j is large enough, it turns out that the set of digraphs in \mathcal{D}_i on j vertices is empty, implying that \mathcal{D}_i is finite.

We then consider every digraph $D \in \mathcal{D}_i$ and check whether D is 2-dicolourable. When it is not, since it admits a uv -colouring for every arc uv , we conclude that D is 3-dicritical. \square

We are now ready to conclude the proof of Theorem 7.

Proof. By Lemmas 10, 12, 16, 17, 18, 22, and 23, we have that every 3-dicritical semi-complete digraph that is not contained in $\bigcup_{i=1}^7 \mathcal{D}_i$ is one of \overrightarrow{W}_3 , $\mathcal{R}(\overleftarrow{K}_2, \overleftarrow{K}_2)$, $\mathcal{R}(\overleftarrow{K}_2, \overrightarrow{C}_3)$, $\mathcal{R}(\overrightarrow{C}_3, \overleftarrow{K}_2)$, and $\mathcal{R}(\overrightarrow{C}_3, \overrightarrow{C}_3)$. The statement then follows directly from Lemma 24. \square

5 Maximum number of arcs in 3-dicritical digraphs

This section is devoted to the proof of Theorem 9. We need a collection of intermediate results. We first show that the bidirected part of a 3-dicritical digraph is a forest unless D is a bidirected odd cycle.

Proposition 25. *Let D be a 3-dicritical digraph that is not a bidirected odd cycle. Then $B(D)$ is a forest.*

Proof. Assume for a contradiction that $B(D)$ is not a forest. Then it contains a cycle $C = u_1u_2 \dots u_pu_1$. Let \overleftrightarrow{C} be the bidirected cycle in D corresponding to C . The cycle C cannot be odd, for otherwise \overleftrightarrow{C} would be a bidirected odd cycle, and $D = \overleftrightarrow{C}$ because a bidirected odd cycle is 3-dicritical, a contradiction. Hence C is an even cycle. By Lemma 10, there exists a 2-dicolouring ϕ of $D \setminus \{u_1u_p\}$. Necessarily, u_1 and u_p are coloured differently because there is a bidirected path of odd length between u_1 and u_p . Thus ϕ is a 2-dicolouring of D , a contradiction. \square

For the remainder of this section we need a few specific definitions. Let T be a tree and $V_3(T)$ be the set of vertices of degree at least 3 in T . Two vertices $u, v \in V(T)$ form an *odd pair* if they are non-adjacent and $\text{dist}_T(u, v)$ is odd, where $\text{dist}_T(u, v)$ denotes the length of the unique path between

u and v in T . The set of odd pairs of T is denoted by $\text{OP}(T)$ and its cardinality is denoted by $\text{op}(T)$. We finally define the *dearth* of T as follows:

$$\text{dearth}(T) = \sum_{v \in V_3(T)} \frac{1}{6} d(v)(d(v) - 1) + \text{op}(T).$$

We first prove that the dearth of a tree is always at least a fraction of its order.

Lemma 26. *Let T be a tree on n vertices for some positive integer n . Then $\text{dearth}(T) \geq \frac{1}{3}n - 1$.*

Proof. For the sake of a contradiction, suppose that T is a counterexample to the statement whose number of vertices is minimum. Clearly, we have $n \geq 4$. The following claim excludes a collection of simple structures of T .

Claim 27. T is neither a path nor a tree.

Proof of the claim. The statement follows from the following simple case distinction.

Case 1: T is a path of even length.

For every odd $i \in \{1, \dots, n-3\}$, as $n \geq 4$, there are exactly i distinct pairs of vertices at distance exactly $n-i$ in T . Hence $\text{dearth}(T) \geq \text{op}(T) = \sum_{i=1}^{\frac{n-2}{2}} (2i-1) = \left(\frac{n-2}{2}\right)^2 \geq \frac{1}{3}n - 1$.

Case 2: T is a path of odd length.

For every odd $i \in \{1, \dots, n-2\}$, as $n \geq 4$, there are exactly i distinct pairs of vertices at distance exactly $n-i$ in T . Hence $\text{dearth}(T) \geq \text{op}(T) = \sum_{i=1}^{\frac{n-3}{2}} 2i = \left(\frac{n-3}{2}\right) \left(\frac{n-1}{2}\right) \geq \frac{1}{3}n - 1$.

Case 3: T is a star on $n \geq 4$ vertices.

As $n \geq 4$, we obtain that $\text{dearth}(T)$ is exactly $\frac{1}{6}(n-1)(n-2)$, and so $\text{dearth}(T) \geq \frac{1}{3}n - 1$.

In either case, we obtain a contradiction to the choice of T . ◇

By Claim 27, we obtain that T is neither a path nor a star. In particular, it follows that T contains an edge uv such that $d_T(u) \geq 2$ and $d_T(v) \geq 3$. Let v_1, \dots, v_r be the neighbours of v in T , where $v_1 = u$ and $r = d_T(v) \geq 3$. For each $i \in [r]$, let T_i be the component of $T - v$ containing v_i . By the choice of T , we have $\text{dearth}(T_i) \geq \frac{1}{3}n(T_i) - 1$. Since the T_i s are pairwise disjoint and none of them contains v , and because u has a neighbour in T_1 at distance exactly 3 from v_2, \dots, v_r we obtain:

$$\begin{aligned} \text{dearth}(T) &\geq \sum_{i=1}^r \text{dearth}(T_i) + \frac{1}{6}r(r-1) + (r-1) \\ &\geq \frac{1}{3}(n(T) - 1) - r + \frac{1}{6}r(r-1) + (r-1) \\ &\geq \frac{1}{3}n(T) - 1, \end{aligned}$$

where in the last inequality we used $r \geq 3$. This contradicts the choice of T . □

Lemma 28. *Let D be a 3-dicritical digraph distinct from $\{\overleftrightarrow{K}_3, \overrightarrow{W}_3\}$ and \overleftrightarrow{T} be a bidirected tree contained in D . Then we have*

$$|\{\{u, v\} \subseteq V(T) \mid \{uv, vu\} \cap A(D) = \emptyset\}| \geq \text{dearth}(T).$$

Proof. Set $\mathcal{O} = \{\{u, v\} \subseteq V(T) \mid \{uv, vu\} \cap A(D) = \emptyset\}$. For every vertex $v \in V_3(T)$, let $\mathcal{O}_v = \mathcal{O} \cap (N_T(v) \times N_T(v))$. Finally let $\mathcal{O}_{\text{odd}} = \mathcal{O} \cap \text{OP}(T)$.

Let us first show that these sets are pairwise disjoint. Let $u, v \in V_3(T)$ be two vertices of degree at least 3 in T . Since T is a tree, we have that $N_T(u) \cap N_T(v)$ contains at most one vertex, implying that $\mathcal{O}_u \cap \mathcal{O}_v = \emptyset$. Also note that vertices in $N_T(v)$ are at distance exactly 2 from each other, so $\mathcal{O}_v \cap \mathcal{O}_{\text{odd}} = \emptyset$. This implies

$$|\mathcal{O}| \geq \sum_{v \in V_3(T)} |\mathcal{O}_v| + |\mathcal{O}_{\text{odd}}|.$$

Hence it is sufficient to prove $|\mathcal{O}_v| \geq \frac{1}{6}d_T(v)(d_T(v) - 1)$ for every $v \in V_3(T)$ and $\mathcal{O}_{\text{odd}} = \text{OP}(T)$ to prove Lemma 28.

Let $v \in V_3(T)$ and u, x, z be three distinct vertices in $N_T(v)$. We claim that $D[\{u, x, z\}]$ contains at most two arcs. If this is not the case, then $D[\{u, x, z\}]$ contains a digon, a directed triangle or a transitive tournament on three vertices. This implies that $D[\{u, x, z, v\}]$ contains \overrightarrow{K}_3 , \overrightarrow{W}_3 or O_4 . By Theorem 7 and Lemma 16, in each case, we obtain a contradiction to the choice of D . Since this holds for every choice of three distinct vertices in $N_T(v)$ and each pair of vertices in $N_T(v)$ is contained in $d_T(v) - 2$ triples, we deduce the following inequality

$$m(D[N_T(v)]) \cdot (d_T(v) - 2) = \sum_{\substack{X \subseteq N_T(v), \\ |X|=3}} m(D[X]) \leq 2 \cdot \binom{d_T(v)}{3},$$

implying that $m(D[N_T(v)]) \leq \frac{1}{3}d_T(v)(d_T(v) - 1)$. Therefore, we obtain $|\mathcal{O}_v| = \binom{d_T(v)}{2} - m(D[N_T(v)]) \geq \frac{1}{6}d_T(v)(d_T(v) - 1)$ as desired.

To show $\mathcal{O}_{\text{odd}} = \text{OP}(T)$, it is sufficient to show that if $\{u, v\}$ is an odd pair then $\{uv, vu\} \cap A(D) = \emptyset$. Assume this is not the case, then by Lemma 10 $D' = D \setminus \{uv, vu\}$ admits a 2-dicolouring ϕ in which $\phi(u) = \phi(v)$, a contradiction since u and v are connected by a bidirected odd path in D' . This shows the claim. \square

We are now ready to prove Theorem 9 that we first restate here for convenience.

Theorem 9. *If D is a 3-dicritical digraph distinct from \overrightarrow{K}_3 and \overrightarrow{W}_3 , then $m(D) \leq \binom{n}{2} + \frac{2}{3}n$.*

Proof. Let D be such a digraph. If D is a bidirected odd cycle, we have $n \geq 5$ and hence D has $2n \leq \binom{n}{2} + \frac{2}{3}n$ arcs, so the result trivially holds. Henceforth assume D is not a bidirected cycle. Then, by Proposition 25, $B(D)$ is a forest. Let T_1, \dots, T_s be the connected components of $B(D)$. For every $i \in [s]$, the number of digons in $D[V(T_i)]$ is exactly $n(T_i) - 1$, whereas the number of pairs of non-adjacent vertices is at least $\text{dearth}(T_i)$ by Claim 28. Hence, since there is no digon between the T_i s, we obtain

$$\begin{aligned} m(D) &\leq \binom{n}{2} + \sum_{i=1}^s ((n(T_i) - 1) - \text{dearth}(T_i)) \\ &\leq \binom{n}{2} + \sum_{i=1}^s \left((n(T_i) - 1) - \left(\frac{1}{3}n(T_i) - 1\right) \right) \quad \text{by Claim 26} \\ &= \binom{n}{2} + \frac{2}{3}n, \end{aligned}$$

which concludes the proof. \square

6 Conclusion

In this paper, we showed that the number of 3-dicritical semi-complete digraphs is finite and with a computer-assisted proof, we gave a full characterization of them. This result seems to be only the tip of an iceberg, and natural generalizations in several directions can be considered.

First, the conjecture of Hoshino and Kawarabayashi on the maximum density of 3-dicritical oriented graphs remains widely open.

We believe that almost all 3-dicritical digraphs are sparser than tournaments. We thus propose the following conjecture which would imply Theorem 6 and asymptotically improve on Theorem 9.

Conjecture 29. There is only a finite number of 3-dicritical digraphs D on n vertices that satisfy $m(D) \geq \binom{n}{2}$.

It is an interesting challenge to generalize the results obtained in this article to $k \geq 4$. In particular, we would be interested in a confirmation of the following statement.

Conjecture 30. For every $k \geq 4$, there is only a finite number of k -dicritical semi-complete digraphs.

Finally, it is also natural to consider a different notion of criticality. A digraph D is called *3-vertex-dicritical* if D is not 3-dicolourable, but $D - v$ is for all $v \in V(D)$. Observe that every 3-dicritical digraph is 3-vertex-dicritical, but the converse is not necessarily true. One can hence wonder whether an analogue of Theorem 6 is true for 3-vertex-dicritical digraphs. However, this turns out not to be the case. Chen et al. proved in [6] that it is NP-hard to decide whether a given tournament is 2-dicolourable. An infinite collection of 3-vertex-dicritical tournaments can easily be derived from their proof.

References

- [1] P. Aboulker. Personal communication, 2023.
- [2] P. Aboulker, T. Bellitto, F. Havet, and C. Rambaud. On the minimum number of arcs in k -dicritical oriented graphs. [arXiv preprint arXiv:2207.01051](#), 2022.
- [3] J. Bang-Jensen, T. Bellitto, T. Schweser, and M. Stiebitz. Hajós and Ore constructions for digraphs. [The Electronic Journal of Combinatorics](#), 27(1), Mar. 2020.
- [4] J. Bang-Jensen and G. Z. Gutin. [Digraphs: Theory, Algorithms and Applications](#). Springer-Verlag, London, 2nd edition, 2009.
- [5] T. Bellitto, N. Bousquet, A. Kabela, and T. Pierron. The smallest 5-chromatic tournament. [Mathematics of Computation](#), 93(345):443–458, 2024.
- [6] X. Chen, X. Hu, and W. Zang. A min-max theorem on tournaments. [SIAM Journal on Computing](#), 37(3):923–937, Jan. 2007.
- [7] R. Diestel. [Graph Theory](#). Graduate texts in mathematics. Springer, Berlin, Germany, 5th edition, June 2017.
- [8] G. A. Dirac. A property of 4-chromatic graphs and some remarks on critical graphs. [Journal of the London Mathematical Society](#), s1-27(1):85–92, 1952.
- [9] G. A. Dirac. Circuits in critical graphs. [Monatshefte für Mathematik](#), 59(3):178–187, 1955.

- [10] G. A. Dirac. A theorem of R. L. Brooks and a conjecture of H. Hadwiger. Proceedings of the London Mathematical Society, s3-7(1):161–195, 1957.
- [11] G. A. Dirac. On the structure of k -chromatic graphs. Mathematical Proceedings of the Cambridge Philosophical Society, 63(3):683–691, 1967.
- [12] P. Erdos. Problems and results in chromatic graph theory. Proof techniques in graph theory, pages 27–35, 1969.
- [13] T. Gallai. Kritische Graphen I. Publications of the Mathematical Institute of the Hungarian Academy of Sciences, 8:165–192, 1963.
- [14] T. Gallai. Kritische Graphen II. Magyar Tudományos Akadémia Matematikai Kutató Intézetének Közleményei, 8:373–395, 1963.
- [15] G. Hajós. Über eine Konstruktion nicht n -farbbarer Graphen. Wissenschaftliche Zeitschrift der Martin-Luther-Universität Halle-Wittenberg. Mathematisch-Naturwissenschaftliche Reihe, 10:116–117, 1961.
- [16] F. Havet, L. Picasarri-Arrieta, and C. Rambaud. On the minimum number of arcs in 4-dicritical oriented graphs. arXiv preprint arXiv:2306.10784, 2023.
- [17] R. Hoshino and K.-i. Kawarabayashi. The edge density of critical digraphs. Combinatorica, 35(5):619–631, 2015.
- [18] A. V. Kostochka and M. Stiebitz. On the number of edges in colour-critical graphs and hypergraphs. Combinatorica, 20(4):521–530, Apr. 2000.
- [19] A. V. Kostochka and M. Stiebitz. A new lower bound on the number of edges in colour-critical graphs and hypergraphs. Journal of Combinatorial Theory, Series B, 87(2):374–402, 2003.
- [20] A. V. Kostochka and M. Stiebitz. The minimum number of edges in 4-critical digraphs of given order. Graphs and Combinatorics, 36(3):703–718, May 2020.
- [21] A. V. Kostochka and M. Yancey. Ore’s conjecture on color-critical graphs is almost true. Journal of Combinatorial Theory, Series B, 109:73–101, Nov. 2014.
- [22] M. Krivelevich. On the minimal number of edges in color-critical graphs. Combinatorica, 17(3):401–426, 1997.
- [23] C.-H. Liu and L. Postle. On the minimum edge-density of 4-critical graphs of girth five. Journal of Graph Theory, 86(4):387–405, 2017.
- [24] C. Luo, J. Ma, and T. Yang. On the maximum number of edges in k -critical graphs. Combinatorics, Probability and Computing, 32(6):900–911, July 2023.
- [25] V. Neumann-Lara. The dichromatic number of a digraph. Journal of Combinatorial Theory, Series B, 33(3):265–270, 1982.
- [26] V. Neumann-Lara. The 3 and 4-dichromatic tournaments of minimum order. Discrete Mathematics, 135(1):233–243, 1994.
- [27] L. Postle. On the minimum number of edges in triangle-free 5-critical graphs. European Journal of Combinatorics, 66:264–280, 2017. Selected papers of EuroComb15.
- [28] M. Stiebitz. Subgraphs of colour-critical graphs. Combinatorica, 7:303–312, 1987.
- [29] B. Toft. On the maximal number of edges of critical k -chromatic graphs. Studia Scientiarum Mathematicarum Hungarica, 5:461–470, 1970.

B Code used in the proof of Theorem 7

This appendix contains the code used in the proof of Theorem 7. In Appendix B.0, we give a collection of useful subroutines we use in the main part of the code. In Appendices B.1, B.2, B.3, and B.4, we give the code use in the proofs of Lemmas 19 and 20, and Lemmas 22 and 24, respectively.

B.0 Preliminaries for the code

In the following code, we give a collection of subroutines we use in our code.

```
1 # The following function displays a progress bar
2 def printProgressBar (iteration, total):
3     percent = ("{0:.1f}").format(100 * (iteration / float(total)))
4     filledLength = int(50 * iteration // total)
5     bar = "#" * filledLength + "-" * (50 - filledLength)
6     print(f"\rProcess: |{bar}| {percent}% Complete", end = "\r")
7     # Print New Line on Complete
8     if iteration == total:
9         print()
10
11 # k,n : integers such that k < 3**n
12 # Returns: the decomposition of k in base 3 of length n
13 def ternary(k,n):
14     b = 3**(n-1)
15     res=""
16     for i in range(n):
17         if(k >= 2*b):
18             k -= 2*b
19             res = res + "2"
20         elif(k >= b):
21             k -= b
22             res = res + "1"
23         else:
24             res = res + "0"
25         b /= 3
26     return res
27
28 # d: DiGraph
29 # u: vertex
30 # v: vertex
31 # Returns: True if and only if d contains a directed path from u to v
32 def contains_directed_path(d,u,v):
33     to_be_treated = [u]
34     i=0
35     while(len(to_be_treated) != i):
36         x = to_be_treated[i]
37         if (x==v):
38             return True
39         for y in d.neighbors_out(x):
40             if (not y in to_be_treated):
41                 to_be_treated.append(y)
42         i+=1
43     return False
44
45 # d: DiGraph
```

```

46 # u: vertex of d
47 # v: vertex of d
48 # current_colouring: partial 2-dicolouring with colours {0,1} of d such that
   current_colouring[u] = current_colouring[v] = 0
49 # Returns: True if and only if current_colouring can be extended into a 2-
   dicolouring of d with no monochromatic directed path from u to v.
50 def can_be_subgraph_of_3_dicritical_aux(d,u,v, current_colouring):
51     #build the colour classes
52     colours = {}
53     colours[0] = []
54     colours[1] = []
55     for (x,i) in current_colouring.items():
56         colours[i].append(x)
57     #check whether both colour classes are acyclic
58     for i in range(2):
59         d_i = d.subgraph(colours[i])
60         if(not d_i.is_directed_acyclic()):
61             return False
62     #check whether there is a monochromatic directed path from u to v
63     d_0 = d.subgraph(colours[0])
64     if(contains_directed_path(d_0,u,v)):
65         return False
66     #check whether current_colouring is partial
67     if(len(current_colouring) == d.order()):
68         return True
69     else:
70         #find a vertex x that is not coloured yet
71         x = len(current_colouring)
72         while(x in current_colouring):
73             x-=1
74         #check recursively whether current_colouring can be extended to x
75         for i in range(2):
76             current_colouring[x] = i
77             if(can_be_subgraph_of_3_dicritical_aux(d,u,v,current_colouring)):
78                 return True
79             current_colouring.pop(x, None)
80         return False
81
82 # d: DiGraph
83 # forbidden_subtournaments: list of DiGraphs
84 # Returns: True if and only if d is {forbidden_subdigraphs}-free, {
   forbidden_induced_subdigraphs}-free and, for every arc (u,v) of d, d
   admits a uv-colouring.
85 def can_be_subgraph_of_3_dicritical(d, forbidden_subdigraphs,
   forbidden_induced_subdigraphs):
86     #check whether d contains a forbidden subgraph
87     for T in forbidden_subdigraphs:
88         if(d.subgraph_search(T, False) != None):
89             return False
90     #check whether d contains a forbidden induced subgraph
91     for T in forbidden_induced_subdigraphs:
92         if(d.subgraph_search(T, True) != None):
93             return False
94     #check for every arc uv if d admits a uv-colouring.
95     for e in d.edges():

```

```

96     d_aux = DiGraph(len(d.vertices()))
97     d_aux.add_edges(d.edges())
98     d_aux.delete_edge(e)
99     current_colouring = {}
100    current_colouring[e[0]] = 0
101    current_colouring[e[1]] = 0
102    if(not can_be_subgraph_of_3_dicritical_aux(d_aux,e[0],e[1],
current_colouring)):
103        return False
104    return True
105
106 # d: DiGraph
107 # Returns: True if and only if d is 2-dicolourable
108 def is_two_dicolourable(d):
109     n = d.order()
110     for bipartition in range(2**n):
111         #build the binary word corresponding to the bipartition
112         binary = bin(bipartition)[2:]
113         while(len(binary)<(n)):
114             binary = "0" + binary
115         #build the bipartition
116         V1 = []
117         V2 = []
118         for v in range(n):
119             if(binary[v] == '0'):
120                 V1.append(v)
121             else:
122                 V2.append(v)
123         #check whether (V1,V2) is actually a dicolouring
124         d1 = d.subgraph(V1)
125         d2 = d.subgraph(V2)
126         if(d1.is_directed_acyclic() and d2.is_directed_acyclic()):
127             return True
128     return False
129
130 #C3_C3 is the digraph made of two disjoint directed triangles, the vertices
of one dominating the vertices of the other
131 C3_C3 = DiGraph(6)
132 for i in range(3):
133     C3_C3.add_edge(i,(i+1)%3)
134     C3_C3.add_edge(i+3,((i+1)%3)+3)
135 for j in range(3,6):
136     C3_C3.add_edge(i,j)
137
138 #F is the digraph on nine vertices made of a TT6 u1,...,u6 and the arcs of
the directed triangles u1u2x1u1, u3u4x2u3, u5u6x3u5.
139 F = DiGraph(9)
140 for i in range(6):
141     for j in range(i):
142         F.add_edge(j,i)
143 F.add_edge(6,0)
144 F.add_edge(1,6)
145 F.add_edge(7,2)
146 F.add_edge(3,7)
147 F.add_edge(8,4)

```

```

148 F.add_edge(5,8)
149
150 #TT8 is the transitive tournament on 8 vertices
151 TT8 = DiGraph(8)
152 for i in range(8):
153     for j in range(i):
154         TT8.add_edge(j,i)
155
156 #reversed_TT8 is the set of tournaments, up to isomorphism, obtained from TT8
    by reversing exactly one arc
157 reversed_TT8 = []
158 for e in TT8.edges():
159     rev = DiGraph(8)
160     rev.add_edges(TT8.edges())
161     rev.delete_edge(e)
162     rev.add_edge(e[1],e[0])
163     check = True
164     for T in reversed_TT8:
165         check = check and (not T.is_isomorphic(rev))
166     if(check):
167         reversed_TT8.append(rev)
168
169 #K2 is the complete digraph on 2 vertices
170 K2 = DiGraph(2)
171 K2.add_edge(0,1)
172 K2.add_edge(1,0)
173
174 #S4 is the bidirected star on 4 vertices
175 S4 = DiGraph(4)
176 for i in range(1,4):
177     S4.add_edge(i,0)
178     S4.add_edge(0,i)
179
180 #C3_K2 is the digraph with a directed triangle dominating a digon.
181 C3_K2 = DiGraph(5)
182 for i in range(3):
183     C3_K2.add_edge(i,(i+1)%3)
184     for j in range(3,5):
185         C3_K2.add_edge(i,j)
186 C3_K2.add_edge(3,4)
187 C3_K2.add_edge(4,3)
188
189 #K2_C3 is the digraph with a digon dominating a directed triangle
190 K2_C3 = DiGraph(5)
191 for i in range(3):
192     K2_C3.add_edge(i,(i+1)%3)
193     for j in range(3,5):
194         K2_C3.add_edge(j,i)
195 K2_C3.add_edge(3,4)
196 K2_C3.add_edge(4,3)
197
198 #K2_K2 is the digraph with a digon dominating a digon
199 K2_K2 = DiGraph(4)
200 for i in range(2):
201     K2_K2.add_edge(i,(i+1)%2)

```



```

202     K2_K2.add_edge(i+2,((i+1)%2)+2)
203     for j in range(2,4):
204         K2_K2.add_edge(i,j)
205
206 #04 and 05 are the obstructions described in the paper.
207 04 = DiGraph(4)
208 for i in range(1,4):
209     04.add_edge(0,i)
210 for i in range(1,3):
211     04.add_edge(i,3)
212 04.add_edge(1,2)
213 04.add_edge(2,1)
214
215 05 = DiGraph(5)
216 for i in range(1,5):
217     05.add_edge(0,i)
218 for i in range(1,4):
219     05.add_edge(i,4)
220 05.add_edge(1,2)
221 05.add_edge(2,3)
222 05.add_edge(3,1)

```

B.1 The proof of Lemma 19

We here give the code used in the proof of Lemma 19.

Proof. For every tournament $T \in \mathcal{F}$, we check, using the following code, if it contains $\vec{C}_3 \Rightarrow \vec{C}_3$ as a subdigraph or if it admits no uv -colouring for an arc uv . This is always the case except when $T \in \{T^1, T^2, T^3, T^4\}$. The Claim then follows by Lemmas 10 and 18. \square

```

1 load("tools.sage")
2
3 # binary_code: a string of fifteen characters '0' and '1'
4 # Returns: a tournament of  $\mathcal{F}$ . The orientations of the fifteen non-
5   forced arcs correspond to the characters of binary_code.
6 def digraph_blowup_TT3(binary_code):
7     iterator_binary_code = iter(binary_code)
8     d = DiGraph(9)
9     #the vertices 0,...,8 correspond respectively to u_1,...,u_6,x_1,x_2,x_3
10
11     #add the arcs of the TT_6
12     for i in range(6):
13         for j in range(i):
14             d.add_edge(j,i)
15
16     #add the arcs of the directed triangles
17     for i in range(3):
18         d.add_edge(6+i, 2*i)
19         d.add_edge(2*i+1, 6+i)
20
21     missing_edges=[(6,2),(6,3),(6,4),(6,5),(6,7),(6,8),(7,0),(7,1),(7,4)
22                   ,(7,5),(7,8),(8,0),(8,1),(8,2),(8,3)]
23     #we orient the missing_edges according to binary_code
24     for e in missing_edges:

```

```

23     if(next(iterator_binary_code) == '0'):
24         d.add_edge(e[0],e[1])
25     else:
26         d.add_edge(e[1],e[0])
27     return d
28
29 print("Computing all possible candidates of \mathcal{F} for being a
    subtournament of a 3-dicritical semi-complete digraph...")
30 list_candidates = []
31 list_forbidden_induced_subdigraphs = [C3_C3]
32
33 #print progress bar
34 printProgressBar(0, 2**15)
35 for i in range(2**15):
36     binary_value = bin(i)[2:]
37     while(len(binary_value)<15):
38         binary_value = '0' + binary_value
39     d = digraph_blowup_TT3(binary_value)
40     if(can_be_subgraph_of_3_dicritical(d,[],
list_forbidden_induced_subdigraphs)):
41         list_candidates.append(d)
42     #update progress bar
43     printProgressBar(i + 1, 2**15)
44
45 print("Number of candidates: ",len(list_candidates),".")
46 for i in range(len(list_candidates)):
47     print("Candidate ",i+1,": ")
48     list_candidates[i].export_to_file("T"+str(i+1)+".pajek")
49     print(list_candidates[i].adjacency_matrix())

```

Running this code produces the following output after roughly 2 minutes of execution on a standard desktop computer:

```

1 Computing all possible candidates of \mathcal{F} for being a subtournament of
  a 3-dicritical semi-complete digraph...
2 Process: |#####| 100.0% Complete
3 Number of candidates: 4 .
4 Candidate 1 :
5 [0 1 1 1 1 1 0 1 1]
6 [0 0 1 1 1 1 1 1 1]
7 [0 0 0 1 1 1 0 0 0]
8 [0 0 0 0 1 1 0 1 0]
9 [0 0 0 0 0 1 0 1 0]
10 [0 0 0 0 0 0 1 1 1]
11 [1 0 1 1 1 0 0 0 1]
12 [0 0 1 0 0 0 1 0 1]
13 [0 0 1 1 1 0 0 0 0]
14 Candidate 2 :
15 [0 1 1 1 1 1 0 1 1]
16 [0 0 1 1 1 1 1 1 1]
17 [0 0 0 1 1 1 1 0 0]
18 [0 0 0 0 1 1 0 1 0]
19 [0 0 0 0 0 1 0 1 0]
20 [0 0 0 0 0 0 0 1 1]
21 [1 0 0 1 1 1 0 1 0]
22 [0 0 1 0 0 0 0 0 1]

```

```

23 [0 0 1 1 1 0 1 0 0]
24 Candidate 3 :
25 [0 1 1 1 1 1 0 0 0]
26 [0 0 1 1 1 1 1 0 1]
27 [0 0 0 1 1 1 1 0 1]
28 [0 0 0 0 1 1 1 1 1]
29 [0 0 0 0 0 1 0 0 0]
30 [0 0 0 0 0 0 0 0 1]
31 [1 0 0 0 1 1 0 1 1]
32 [1 1 1 0 1 1 0 0 0]
33 [1 0 0 0 1 0 0 1 0]
34 Candidate 4 :
35 [0 1 1 1 1 1 0 0 1]
36 [0 0 1 1 1 1 1 0 1]
37 [0 0 0 1 1 1 1 0 1]
38 [0 0 0 0 1 1 1 1 0]
39 [0 0 0 0 0 1 0 0 0]
40 [0 0 0 0 0 0 0 0 1]
41 [1 0 0 0 1 1 0 1 0]
42 [1 1 1 0 1 1 0 0 1]
43 [0 0 0 1 1 0 1 0 0]

```

The graphs in the output are exactly T_1, T_2, T_3 , and T_4 .

B.2 The proof of Lemma 20

We here give the code used in the proof of Lemma 20.

```

1 import networkx
2 load("tools.sage")
3
4 list_candidates = []
5 list_forbidden_induced_subdigraphs = [C3_C3]
6
7 #import T1, T2, T3 and T4
8 for i in range(1,5):
9     candidate = DiGraph(9)
10    nx = networkx.read_pajek("T"+str(i)+".pajek")
11    for e in nx.edges():
12        candidate.add_edge(int(e[0]),int(e[1]))
13    list_candidates.append(candidate)
14
15 print("We start from the ", len(list_candidates), " candidates on 9 vertices.
16    ")
17 #We want to prove that a 3-dicritical semi-complete digraph does not contain
18    a digraph in {F+,F-}. By directional duality, it is sufficient to prove
19    that it does not contain F+.
20 #For each candidate computed above, we try to add a new vertex that dominates
21    the transitive tournament, and then we build every possible orientation
22    between this vertex and the three other vertices.
23 print("Computing for F+...")
24 list_Fp = []
25 printProgressBar(0, 8)
26 for orientation in range(2**3):
27     binary = bin(orientation)[2:]

```

```

24 while(len(binary)<3):
25     binary = '0' + binary
26 for T9 in list_candidates:
27     iterator = iter(binary)
28     T10 = DiGraph(10)
29     T10.add_edges(T9.edges())
30     for v in range(6):
31         T10.add_edge(9,v)
32     for v in range(6,9):
33         if(next(iterator) == '0'):
34             T10.add_edge(v,9)
35         else:
36             T10.add_edge(9,v)
37     check = can_be_subgraph_of_3_dicritical(T10,[],
list_forbidden_induced_subdigraphs)
38     if(check):
39         list_Fm.append(T2)
40     printProgressBar(orientation+1, 8)
41
42 print("Number of 1-extensions of {T1,T2,T3,T4} containing F+: ",len(list_Fp))

```

Running this code produces the following output after roughly 1 second of execution on a standard desktop computer:

```

1 We start from the 4 candidates on 9 vertices.
2 Computing for F+...
3 Process: |#####| 100.0% Complete
4 Number of 1-extensions of {T1,T2,T3,T4} containing F+: 0

```

B.3 The proof of Lemma 22

We here give the code used in the proof of Lemma 22.

```

1 import networkx
2 load("tools.sage")
3
4 all_candidates = []
5 current_candidates = []
6 next_candidates = []
7
8 list_forbidden_subdigraphs = [S4, K2_K2, O4, O5, K2_C3, C3_K2, C3_C3]
9 list_forbidden_induced_subdigraphs = [TT8]
10
11 #import the candidates T1, ..., T4 on 9 vertices:
12 for i in range(1,5):
13     candidate = DiGraph(9)
14     nx = networkx.read_pajek("T"+str(i)+".pajek")
15     for e in nx.edges():
16         candidate.add_edge(int(e[0]),int(e[1]))
17     current_candidates.append(candidate)
18 print("We start from the tournaments {T^1,T^2,T^3,T^4} on 9 vertices, and
look for every possible completion of them that is potentially a
subdigraph of a larger 3-dicritical semi-complete digraphs.\n")
19
20
21 all_candidates.extend(current_candidates)

```

```

22 #completions of T1, ..., T4
23 while(len(current_candidates)>0):
24     for old_D in current_candidates:
25         for e in old_D.edges():
26             #we try to complete old_D by replacing e by a digon. It actually
                makes sense only if e is not already in a digon.
27             if(not (e[1],e[0],None) in old_D.edges()):
28                 new_D = DiGraph(9)
29                 new_D.add_edges(old_D.edges())
30                 new_D.add_edge(e[1],e[0])
31                 #we check whether this completion of old_D is potentially a
                subdigraph of a larger 3-dicritical semi-complete digraph.
32                 check = can_be_subgraph_of_3_dicritical(new_D,
                list_forbidden_subdigraphs, list_forbidden_induced_subdigraphs)
33                 for D in next_candidates:
34                     check = check and (not D.is_isomorphic(new_D))
35                 if(check):
36                     next_candidates.append(new_D)
37             all_candidates.extend(next_candidates)
38             current_candidates = next_candidates
39             next_candidates=[]
40
41 print("-----")
42 print("There are",len(all_candidates),"possible completions (up to
                isomorphism) of {T^1,T^2,T^3,T^4} that are potentially subdigraphs of a
                larger 3-dicritical semi-complete digraphs.\n")
43
44
45 count_dic_3 = 0
46 for D in all_candidates:
47     if(not is_two_dicolourable(D)):
48         count_dic_3 += 1
49 print(count_dic_3, " of them have dichromatic number at least 3. In
                particular,",count_dic_3,"of them are 3-dicritical.\n")
50
51 current_candidates = all_candidates
52 next_candidates = []
53 for n in range(10,12):
54     #computes the extensions on n vertices of {T1,T2,T3,T4}
55     print("-----")
56     print("Computing "+str(n-9)+"-extensions of the candidates on 9 vertices
                that are potentially subtournaments of 3-dicritical tournaments (up to
                isomorphism).")
57     printProgressBar(0, 3**(n-1))
58
59     for orientation in range(3**(n-1)):
60         ternary_code = ternary(orientation,n-1)
61         #build every 1-extension of current_candidates
62         for old_D in current_candidates:
63             new_D = DiGraph(n)
64             new_D.add_edges(old_D.edges())
65             for v in range(n-1):
66                 if(ternary_code[v] == '0'):
67                     new_D.add_edge(v,n-1)
68                 elif(ternary_code[v]=='1'):

```

```

69         new_D.add_edge(n-1,v)
70     else:
71         new_D.add_edge(v,n-1)
72         new_D.add_edge(n-1,v)
73     check = can_be_subgraph_of_3_dicritical(new_D,
list_forbidden_subdigraphs ,list_forbidden_induced_subdigraphs)
74     for D in next_candidates:
75         check = check and (not new_D.is_isomorphic(D))
76     if(check):
77         next_candidates.append(new_D)
78     printProgressBar(orientation+1, 3**(n-1))
79
80     print("Number of ", n-9,"-extensions up to isomorphism: ",len(
next_candidates))
81     #check if one of the candidates has dichromatic number at least 3.
82     count_dic_3 = 0
83     for D in next_candidates:
84         if(not is_two_dicolourable(D)):
85             count_dic_3 += 1
86     print(count_dic_3, " of them have dichromatic number at least 3. In
particular,", count_dic_3,"of them are 3-dicritical.\n")
87     current_candidates = next_candidates
88     next_candidates = []

```

Running this code produces the following output after roughly 12 minutes of execution on a standard desktop computer:

```

1 We start from the tournaments {T^1,T^2,T^3,T^4} on 9 vertices, and look for
every possible completion of them that is potentially a subdigraph of a
larger 3-dicritical semi-complete digraphs.
2
3 -----
4 There are 14 possible completions (up to isomorphism) of {T^1,T^2,T^3,T^4}
that are potentially subdigraphs of a larger 3-dicritical semi-complete
digraphs.
5
6 0 of them have dichromatic number at least 3. In particular, 0 of them are
3-dicritical.
7
8 -----
9 Computing 1-extensions of the candidates on 9 vertices that are potentially
subtournaments of 3-dicritical tournaments (up to isomorphism).
10 Process: |#####| 100.0% Complete
11 Number of 1 -extensions up to isomorphism: 34
12 0 of them have dichromatic number at least 3. In particular, 0 of them are
3-dicritical.
13
14 -----
15 Computing 2-extensions of the candidates on 9 vertices that are potentially
subtournaments of 3-dicritical tournaments (up to isomorphism).
16 Process: |#####| 100.0% Complete
17 Number of 2 -extensions up to isomorphism: 0
18 0 of them have dichromatic number at least 3. In particular, 0 of them are
3-dicritical.

```

B.4 The proof of Lemma 24

We here give the code used in the proof of Lemma 24.

```
1 load("tools.sage")
2
3 def possible_completions(graph_to_complete, nb_vertices,
4 list_forbidden_subdigraphs, list_forbidden_induced_subdigraphs, progress
5 =0):
6     if(progress == nb_vertices-1):
7         return [graph_to_complete]
8     else:
9         result = []
10        for i in range(3):
11            #we make a copy of the graph_to_complete
12            new_D = DiGraph(nb_vertices)
13            new_D.add_edges(graph_to_complete.edges())
14
15            #we consider every possible orientation between the vertices (
16            nb_vertices-1) and (progress)
17            if(i==0):
18                new_D.add_edge(nb_vertices-1, progress)
19            elif(i==1):
20                new_D.add_edge(progress, nb_vertices-1)
21            else:
22                new_D.add_edge(nb_vertices-1, progress)
23                new_D.add_edge(progress, nb_vertices-1)
24
25            #for each of the 3 possible orientations, we check whether the
26            obtained digraph is already an obstruction. If it is not, we compute all
27            possible completions recursively
28            if(can_be_subgraph_of_3_dicritical(new_D,
29 list_forbidden_subdigraphs, list_forbidden_induced_subdigraphs)):
30                result.extend(possible_completions(new_D, nb_vertices,
31 list_forbidden_subdigraphs, list_forbidden_induced_subdigraphs, progress
32 +1))
33        return result
34
35 for tt in range(1,8):
36     transitive_tournament = DiGraph(tt)
37     for i in range(tt):
38         for j in range(i):
39             transitive_tournament.add_edge(j,i)
40
41     next_transitive_tournament = DiGraph(tt+1)
42     for i in range(tt+1):
43         for j in range(i):
44             next_transitive_tournament.add_edge(j,i)
45
46     list_forbidden_subdigraphs = [S4, K2_K2, 04, 05, K2_C3, C3_K2, C3_C3, F]
47     list_forbidden_induced_subdigraphs = []
48     if(tt<7):
49         list_forbidden_induced_subdigraphs = [next_transitive_tournament]
50     else:
51         list_forbidden_induced_subdigraphs = reversed_TT8
```

```

45 print("\n-----\n")
46 print("Generating all 3-dicritical semi-complete digraphs with maximum
acyclic induced subdigraph of size exactly " + str(tt) + ".")
47
48 n=tt+1
49 candidates = [transitive_tournament]
50 next_candidates = []
51 while(len(candidates)>0):
52     print("\nComputing candidates on "+str(n)+" vertices.")
53     printProgressBar(0, len(candidates))
54     for i in range(len(candidates)):
55         old_D = candidates[i]
56         new_D = DiGraph(n)
57         new_D.add_edges(old_D.edges())
58         all_possible_completions_new_D = possible_completions(new_D, n,
list_forbidden_subdigraphs, list_forbidden_induced_subdigraphs)
59         for candidate in all_possible_completions_new_D:
60             check = True
61             for D in next_candidates:
62                 check = not D.is_isomorphic(candidate)
63                 if(not check):
64                     break
65             if(check):
66                 next_candidates.append(candidate)
67             printProgressBar(i + 1, len(candidates))
68
69             #check the candidates that are actually 3-dicritical.
70             print("We found",len(next_candidates),"candidates on "+str(n)+"
vertices.")
71             dicriticals = []
72             for D in next_candidates:
73                 if(not is_two_dicolourable(D)):
74                     dicriticals.append(D)
75             print(len(dicriticals), " of them are actually 3-dicritical.\n")
76             for D in dicriticals:
77                 print("adjacency matrix of a 3-dicritical digraph that we found:"
)
78                 print(D.adjacency_matrix())
79
80             candidates = next_candidates
81             next_candidates = []
82             n+=1

```

Running this code produces the following output after roughly 2 hours of execution on a standard desktop computer:

```

1 -----
2
3 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
induced subdigraph of size exactly 1.
4
5 Computing candidates on 2 vertices.
6 Process: |#####| 100.0% Complete
7 We found 1 candidates on 2 vertices.
8 0 of them are actually 3-dicritical.
9

```



```

10
11 Computing candidates on 3 vertices.
12 Process: |#####| 100.0% Complete
13 We found 1 candidates on 3 vertices.
14 1 of them are actually 3-dicritical.
15
16 adjacency matrix of a 3-dicritical digraph that we found:
17 [0 1 1]
18 [1 0 1]
19 [1 1 0]
20
21 Computing candidates on 4 vertices.
22 Process: |#####| 100.0% Complete
23 We found 0 candidates on 4 vertices.
24 0 of them are actually 3-dicritical.
25
26
27 -----
28
29 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
    induced subdigraph of size exactly 2.
30
31 Computing candidates on 3 vertices.
32 Process: |#####| 100.0% Complete
33 We found 5 candidates on 3 vertices.
34 0 of them are actually 3-dicritical.
35
36
37 Computing candidates on 4 vertices.
38 Process: |#####| 100.0% Complete
39 We found 5 candidates on 4 vertices.
40 0 of them are actually 3-dicritical.
41
42
43 Computing candidates on 5 vertices.
44 Process: |#####| 100.0% Complete
45 We found 0 candidates on 5 vertices.
46 0 of them are actually 3-dicritical.
47
48
49 -----
50
51 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
    induced subdigraph of size exactly 3.
52
53 Computing candidates on 4 vertices.
54 Process: |#####| 100.0% Complete
55 We found 13 candidates on 4 vertices.
56 0 of them are actually 3-dicritical.
57
58
59 Computing candidates on 5 vertices.
60 Process: |#####| 100.0% Complete
61 We found 37 candidates on 5 vertices.
62 1 of them are actually 3-dicritical.

```

```

63 adjacency matrix of a 3-dicritical digraph that we found:
64 [0 1 1 0 0]
65 [0 0 1 0 1]
66 [0 0 0 1 1]
67 [1 1 0 0 1]
68 [1 0 1 1 0]
69
70
71 Computing candidates on 6 vertices.
72 Process: |#####| 100.0% Complete
73 We found 8 candidates on 6 vertices.
74 0 of them are actually 3-dicritical.
75
76
77 Computing candidates on 7 vertices.
78 Process: |#####| 100.0% Complete
79 We found 1 candidates on 7 vertices.
80 1 of them are actually 3-dicritical.
81
82 adjacency matrix of a 3-dicritical digraph that we found:
83 [0 1 1 0 0 0 1]
84 [0 0 1 0 1 1 0]
85 [0 0 0 1 0 1 1]
86 [1 1 0 0 0 1 0]
87 [1 0 1 1 0 0 0]
88 [1 0 0 0 1 0 1]
89 [0 1 0 1 1 0 0]
90
91 Computing candidates on 8 vertices.
92 Process: |#####| 100.0% Complete
93 We found 0 candidates on 8 vertices.
94 0 of them are actually 3-dicritical.
95
96
97 -----
98
99 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
    induced subdigraph of size exactly 4.
100
101 Computing candidates on 5 vertices.
102 Process: |#####| 100.0% Complete
103 We found 27 candidates on 5 vertices.
104 0 of them are actually 3-dicritical.
105
106
107 Computing candidates on 6 vertices.
108 Process: |#####| 100.0% Complete
109 We found 116 candidates on 6 vertices.
110 0 of them are actually 3-dicritical.
111
112
113 Computing candidates on 7 vertices.
114 Process: |#####| 100.0% Complete
115 We found 10 candidates on 7 vertices.
116 0 of them are actually 3-dicritical.

```

```

117
118
119 Computing candidates on 8 vertices.
120 Process: |#####| 100.0% Complete
121 We found 0 candidates on 8 vertices.
122 0 of them are actually 3-dicritical.
123
124
125 -----
126
127 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
    induced subdigraph of size exactly 5.
128
129 Computing candidates on 6 vertices.
130 Process: |#####| 100.0% Complete
131 We found 49 candidates on 6 vertices.
132 0 of them are actually 3-dicritical.
133
134
135 Computing candidates on 7 vertices.
136 Process: |#####| 100.0% Complete
137 We found 266 candidates on 7 vertices.
138 0 of them are actually 3-dicritical.
139
140
141 Computing candidates on 8 vertices.
142 Process: |#####| 100.0% Complete
143 We found 20 candidates on 8 vertices.
144 0 of them are actually 3-dicritical.
145
146
147 Computing candidates on 9 vertices.
148 Process: |#####| 100.0% Complete
149 We found 0 candidates on 9 vertices.
150 0 of them are actually 3-dicritical.
151
152
153 -----
154
155 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
    induced subdigraph of size exactly 6.
156
157 Computing candidates on 7 vertices.
158 Process: |#####| 100.0% Complete
159 We found 80 candidates on 7 vertices.
160 0 of them are actually 3-dicritical.
161
162
163 Computing candidates on 8 vertices.
164 Process: |#####| 100.0% Complete
165 We found 500 candidates on 8 vertices.
166 0 of them are actually 3-dicritical.
167
168
169 Computing candidates on 9 vertices.

```

```

170 Process: |#####| 100.0% Complete
171 We found 39 candidates on 9 vertices.
172 0 of them are actually 3-dicritical.
173
174
175 Computing candidates on 10 vertices.
176 Process: |#####| 100.0% Complete
177 We found 0 candidates on 10 vertices.
178 0 of them are actually 3-dicritical.
179
180
181 -----
182
183 Generating all 3-dicritical semi-complete digraphs with maximum acyclic
    induced subdigraph of size exactly 7.
184
185 Computing candidates on 8 vertices.
186 Process: |#####| 100.0% Complete
187 We found 110 candidates on 8 vertices.
188 0 of them are actually 3-dicritical.
189
190
191 Computing candidates on 9 vertices.
192 Process: |#####| 100.0% Complete
193 We found 459 candidates on 9 vertices.
194 0 of them are actually 3-dicritical.
195
196
197 Computing candidates on 10 vertices.
198 Process: |#####| 100.0% Complete
199 We found 16 candidates on 10 vertices.
200 0 of them are actually 3-dicritical.
201
202
203 Computing candidates on 11 vertices.
204 Process: |#####| 100.0% Complete
205 We found 0 candidates on 11 vertices.
206 0 of them are actually 3-dicritical.

```

The adjacency matrices in the output are exactly those of the digraphs \overleftrightarrow{K}_3 , \mathcal{H}_5 and \mathcal{P}_7 .