



HAL
open science

Functional Extreme-PLS

Stéphane Girard, Cambyse Pakzad

► **To cite this version:**

| Stéphane Girard, Cambyse Pakzad. Functional Extreme-PLS. 2024. hal-04488561

HAL Id: hal-04488561

<https://inria.hal.science/hal-04488561v1>

Preprint submitted on 4 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

FUNCTIONAL EXTREME-PLS

STÉPHANE GIRARD AND CAMBYSE PAKZAD

Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, 38000 Grenoble, France

ABSTRACT. We propose an extreme dimension reduction method extending the Extreme-PLS approach to the case where the covariate lies in a possibly infinite-dimensional Hilbert space. The ideas are partly borrowed from both Partial Least-Squares and Sliced Inverse Regression techniques. As such, the method relies on the projection of the covariate onto a subspace and maximizes the covariance between its projection and the response conditionally to an extreme event to capture most of the tail information. Moreover, we link the covariate and the heavy-tailed response through a non-linear inverse single-index model and our goal is to infer the index in this regression framework. We propose a new family of estimators and show its asymptotic consistency with convergence rates under the model. Assuming mild conditions on the noise, most of the assumptions are stated in terms of regular variation unlike the standard literature on SIR and single-index regression. Finally, we illustrate our results on a finite-sample study with synthetic functional data.

1. INTRODUCTION

Over the last decades, the amount of available data has massively increased, urging the statistical community to shift focus toward the high-dimensional regime where the dimension of the observations is large compared to the sample size, or even infinite for the functional case. This situation turns out to be challenging for many reasons, see the monographs [Bosq, 2000](#); [Ramsay and Silverman, 2005](#); [Ferraty and Vieu, 2006](#). First, from the computational point of view, the time and energy costs may exceed the feasible limits. Second, a deeper phenomena called the curse of dimensionality occurs, whose one facet is sparsity among the observations. A prejudice may be overfitting in the regression context where standard estimation methods exhibit high variance and hence may be untrustworthy.

Regression analysis aims at extracting a relationship between a real response variable Y and a high-dimensional predictor/covariate X , based on their observations. Doing so, it has inherent connections with *sufficient dimension reduction* (SDR), a generalization of the single-index model (see *e.g.*, [Cook, 1998](#)), whose generic purpose is to alleviate the curse of dimensionality in a regression setting. In essence, the idea of dimension reduction is to substitute X by its orthogonal projection $P_S X$ onto a low-dimensional subspace S that carries the same amount of information on $Y|X$. Specifically, one seeks a subspace S such that Y conditionally to X has same distribution as Y conditionally to $P_S X$, or equivalently Y and X are independent conditionally to $P_S X$ ([Cook, 2007](#)). As an example, a powerful, flexible and abundantly studied tool is assuming a single-index model which itself extends standard linear models. Such a setting prescribes $\dim(S) = 1$ and a relationship between X and Y through an unknown link function in such a way that one may explain the values of Y

E-mail address: `stephane.girard@inria.fr` - `cambyse.pakzad@inria.fr`.

Date: March 4, 2024.

2020 Mathematics Subject Classification. 60G70 Extreme value theory; extremal stochastic processes; 62G08 Nonparametric regression and quantile regression; 62G32 Statistics of extreme values and tail inference; 62R10 Functional data analysis.

Key words and phrases. Extremes; Dimension reduction; Functional data analysis; Partial least squares; Inverse regression; Single-index model.

using a linear combination of the X coordinates. The literature is rich and we only mention [Horowitz \(2009\)](#); [Chen et al. \(2011\)](#) on this topic.

Among the popular methods for dimension reduction, the Partial Least Squares (PLS) method, initiated by [Wold \(1975\)](#), has proven to be successful in both theoretical and practical sides; especially in chemometrics ([Martens and Naes, 1992](#)) where it takes root. It combines characteristics of Principal Component Analysis (PCA) and regression. Its core idea is to find a projection of X having high variance and correlation with Y (namely high covariance) meaning that the information in the covariate X that impacts the response Y is encompassed in $P_S X$ while being the most exhaustive.

The literature on high-dimensional PLS regression is scarce. A breakthrough is [Chun and Keleş \(2010\)](#) which show that, under some model, the PLS estimator in linear regression is not consistent unless the dimension of X grows to infinity much slower than the sample size. Interestingly and in the opposite direction, [Cook and Forzani \(2018, 2019\)](#) retrieve consistency with standard rates in the multi-index model and without any regime on the dimension. When the predictor/covariate X is functional, *i.e.*, X lies in an infinite-dimensional space, [Preda and Saporta \(2005\)](#) introduce the first PLS regression method adapted to the particular $L^2([0, 1])$ -case. Most of developments focus on estimating the slope in functional linear models and we refer to [Reiss and Ogden \(2007\)](#); [Delaigle and Hall \(2012\)](#); [Aguilera et al. \(2016\)](#).

Another standard line of research is Sliced Inverse Regression (SIR), first introduced by [Li \(1991\)](#) in the context of SDR, which targets the central subspace $S_{Y|X}$ defined as the minimal subspace S such that Y and X are independent conditionally to $P_S X$. It hinges on the inverse regression 'X against Y' that happens to be simpler than the forward regression, usually at the price of a certain condition of linearity and constant variance. This somewhat contradicts the philosophy of Fisher, as noted and encouraged by [Cook \(2007\)](#), where the inference is rather conditioned on the predictors/covariates X . The literature on SIR is vast and has seen many developments; we refer to [Li et al. \(2007\)](#) for an alternative method, [Lin et al. \(2018, 2021\)](#) for the fine properties (optimality and minimax rates) of SIR in high-dimension, [Tan et al. \(2020\)](#) in the sparse case and more recently [Huang et al. \(2023\)](#) when $\dim(S_{Y|X})$ may not be bounded.

Concerning the functional framework (FSIR), the seminal papers introducing the topic are [Dauxois et al. \(2001\)](#); [Ferré and Yao \(2003\)](#) which prove the consistency (without rate) provided that the true dimension of the central subspace is known. In a complementary way, [Li and Hsing \(2010\)](#) propose a test on the value of the reduced dimension. The question of consistency is complemented by [Forzani and Cook \(2007\)](#); [Cook et al. \(2010\)](#). Herein, it is proven that the central subspace may only be consistently estimated with rate \sqrt{n} by FSIR under strict restrictions on the covariance operator of X . Next, [Lian \(2015\)](#) proves the consistency of FSIR and show that the convergence rate coincide with the minimax rate for the functional linear regression. This connection is explored by [Wang and Lian \(2020\)](#) in the framework of reproducing kernel Hilbert spaces (RKHS) which has already found success in SDR (see *e.g.*, [Fukumizu et al. \(2009\)](#); [Lee and Li \(2022\)](#)). Let us mention two generalizations of FSIR using RKHS to nonlinear spaces: non-Hilbertian sample paths for X in [Hsing and Ren \(2009\)](#), and nonlinear spaces in [Li and Song \(2017\)](#) (see also [Li and Song \(2022\)](#)). For the more realistic situation where the functional data are longitudinal, meaning they are sparsely observed, we may mention [Jiang et al. \(2014\)](#) or very recently [Chatla and Liu \(2024\)](#). About the latest development of FSIR, we refer to [Chen et al. \(2023\)](#) on optimality and minimax rates, and [Tian et al. \(2023\)](#) for a slice-free version.

Now, one may wonder what happens at the extreme regime, *i.e.*, when the focus is on the conditional tail of Y given a high-dimensional covariate X . This question belongs to the vivid research branch of the association of dimension reduction and extreme-value statistics. In this case, the sparsity is doubled in the sense that both the curse of dimensionality and the rarity of tail observations operate. Such hindrance may apply in the estimation of

certain statistics or risk measures in the functional extreme regime (see *e.g.*, Gardes and Girard (2012); Girard et al. (2022)). Thereby, one main aspect in this branch builds up on the existence of a projection $P_S X$ that captures all the information on the extreme values of Y . Instead of using functional estimation methods, one would then expect that replacing X by $P_S X$ as the covariate, and hence opting for a lower-dimensional method, yields comparable or even better performances. Generally speaking, one may distinguish two classes of method, the *unsupervised* case where the variables at hand are somehow exchangeable in their roles, and the *supervised* case where different statuses exist (such as response or covariate). Concerning the former setting, many developments are available in the literature and we refer to Aghbalou et al. (2024) for the latest bibliography material.

Adversely, the supervised situation, especially from a regression perspective, is mostly untouched. This branch builds up on the existence of a projection of X that captures all the information on the extreme values of Y . The pioneer work may be identified as Gardes (2018), later followed up by Gardes (2020), which proposes a new notion of tail conditional independence, the latter property underlying SDR. Applying this to the extremal quantile regression, the author then studies the inference under a single and multi-index model when the dimension is large but finite and when the dimension reduction subspace is known. Still, some theoretical properties are lacking and the computational cost is too high. In this direction, Xu et al. (2020) develop a three-step estimation procedure for such large conditional quantiles and show the \sqrt{n} -consistency under a new tail single-index model when the indexes are unknown; yielding better computational performances. Alternatively and having in sight the prediction of tail events, Aghbalou et al. (2024) introduce a new mathematical framework, including a new definition for the tail conditional independence, for estimating with theoretical guarantees, in the extreme regime and by means of inverse regression, the sufficient dimension reduction subspace, under the standard assumptions for SIR. Intersecting dimension reduction and conditional extreme value theory, a novel method based on the PLS in the extreme regime was recently introduced by Bousebata et al. (2023). Herein, the authors introduce an inverse single-index model. This means that one expresses the covariate in terms of the response and some unknown link function (instead of the converse) which results in a reduction of dimension. In particular, under some conditions and within the regular variation framework, this inverse point of view may also be interpreted as a forward single-index model. Next, they propose an estimator of the index which is shown to be consistent with explicit rate. The nature of their assumptions is quite different from the standard literature of dimension reduction as no conditions on the inverse conditional mean $\mathbb{E}(X|Y)$, such as linearity, nor noise independence or centering are required.

In this work, we take up the methodology introduced in Bousebata et al. (2023) for the supervised branch of extreme dimension reduction and convert it to the functional setting. Specifically, we project the predictor/covariate onto a one-dimensional subspace while capturing most of the information explaining the extreme values of the response variable. As such, we consider covariates that lie in an infinite-dimensional Hilbert space H and the projection is expressed in terms of the associated inner product. Our results equally apply when $H = \mathbb{R}^d$ with d large but finite, thereby positioning Bousebata et al. (2023) as a particular sub-case. Following the philosophy of PLS, we maximize the covariance between the projected predictor/covariate and the response conditionally to the response being larger than an increasing threshold. This corresponds to the extreme framework where the response is assumed to be heavy-tailed. As for the covariate, we echo to SIR by modelling its relationship to the response through a non-linear inverse regression model which relies on a single index, *i.e.*, a deterministic vector in H . We may again recover a forward single-index model interpretation. Thereby, the method at hand induces an effective reduction of dimension as we provide an estimation of the aforementioned index based on the observations. Using a simple geometrical interpretation, we in fact propose a new family of estimators that

generalizes the one in [Bousebata et al. \(2023\)](#) and hence allows more freedom to the statistician. Our theoretical guarantees take form of results on the asymptotic consistency with rates under similar conditions as in [Bousebata et al. \(2023\)](#). In particular, the assumptions are expressed in the language of regular variation and we require minimal conditions about the noise, unlike the standard literature on SIR and forward single-index model. Finally, the final-sample properties of the method are studied by numerical simulations on synthetic functional data.

The remaining of the paper is organized as follows. First, we introduce in [Section 2](#) the mathematical framework for the Functional Extreme Partial Least-Squares (FEPLS in short) and, more specifically, the inverse single-index model. The estimators are presented in [Section 3](#) with the associated consistency results. Next, the FEPLS finite-sample properties are illustrated in [Section 4](#). The respective code in `Python` is publicly available on the website of the authors¹. The mathematical details needed for the derivation of our main results are gathered in [Section 5](#).

2. THEORETICAL FRAMEWORK

Let $(H, \langle \cdot, \cdot \rangle)$ be a separable Hilbert space with its Borel σ -field $\mathcal{B}(H)$ and let $(\Omega, \mathcal{A}, \mathbb{P})$ be a complete probability space. Consider a random pair (Y, X) , where $Y : \Omega \rightarrow \mathbb{R}$ and $X : \Omega \rightarrow H$ are random variables, *i.e.*, Y is Borel measurable and X is strongly \mathbb{P} -measurable in the sense of Bochner, see ([Hytönen et al., 2016](#), Definition 1.1.3).

We define the FEPLS direction as

$$w(y) := \arg \max_{\|w\|=1} \text{Cov}(\langle w, X \rangle, Y \mid Y \geq y), \quad y \in \mathbb{R}, \quad (2.1)$$

where $\|\cdot\|$ is the norm induced by the inner product $\langle \cdot, \cdot \rangle$ and $y \in \mathbb{R}^+$ is a large deterministic threshold. The expectation underlying the covariance in [\(2.1\)](#) is understood in the sense of Lebesgue. The interpretation of [\(2.1\)](#) follows the core idea of PLS but in the extreme regime. Using an optimization framework, we seek the element in H which maximizes the tail information, by considering the covariance (and not the correlation) conditionally to the tail event of Y , shared between Y and the projection of X on a single direction.

In the sequel, we denote the tail-moment of a random object W by $m_W(y) := \mathbb{E}(W 1_{\{Y \geq y\}})$ whenever it exists. In the case where $W \in H$, we interpret the expectation at hand as the Bochner expectation denoted by $\mathbb{E}_{\mathcal{B}}$. Note that, since we consider a separable space, we could equivalently use the Pettis integral instead. We refer to [Hytönen et al. \(2016\)](#) for more details about the notion.

A key fact is that the solution of the optimization problem [\(2.1\)](#) is explicit as we have:

Proposition 2.1. *Suppose that Y is integrable and X is Bochner-integrable. Then, the unique solution of the optimization problem [\(2.1\)](#) is given by, for any $y \in \mathbb{R}$,*

$$w(y) = v(y)/\|v(y)\| \text{ where } v(y) = \bar{F}(y)m_{XY}(y) - m_X(y)m_Y(y). \quad (2.2)$$

This result allows us to undertake an inference procedure based on the empirical counterparts of tail-moments in [\(2.2\)](#). Before that, let us introduce the characteristics of our model. To make sense of the conditional tail event in [\(2.1\)](#), we consider an extreme framework where the response variable Y is heavy-tailed. Equivalently, its associated survival function is such that

$$y \mapsto \bar{F}(y) := \mathbb{P}(Y \geq y) \in \mathcal{RV}_{-1/\gamma}(+\infty), \quad \gamma \in (0, 1), \quad (2.3)$$

where $\mathcal{RV}_a(+\infty)$, $a \in \mathbb{R}$, is the set of real-valued measurable functions f defined on a neighbourhood $[x_0, +\infty)$ of infinity and such that, for any $x > x_0 \geq 0$,

$$\lim_{t \rightarrow +\infty} \frac{f(tx)}{f(t)} = x^a.$$

¹<https://cpakzad.github.io/download/FEPLS.zip>

In such a case, f is said to be regularly-varying at infinity with index a and we refer to [Bingham et al. \(1989\)](#) for further details.

Next, in order to provide theoretical guarantees on the inference method described in the next section and based on (2.1)–(2.2), we consider the following (non-linear) inverse single-index functional model,

$$X = g(Y)\beta + \varepsilon, \quad \beta \in H, \quad \|\beta\| = 1, \quad (2.4)$$

with $\varepsilon : \Omega \rightarrow H$ a random variable being the noise and an (unknown) deterministic link function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $g \in \mathcal{RV}_\kappa(+\infty)$ with $\kappa > 0$.

The appeal of the model (2.4) stems from the fact that, for instance when ε is independent of X and centered, the FEPLS direction (2.1) coincides with the true index: $w(y) = \beta$ for any $y \geq 0$. Heuristically, one expects, assuming more generally that Y and ε are dependent but ε has a small contribution in the tail regime of Y , that

$$\arg \max_{\|w\|=1} \text{Cov}(\langle w, X \rangle, Y \mid Y \geq y) \rightarrow \beta, \quad \text{as } y \rightarrow +\infty.$$

To deal with the conditional expectation involving the tail event $\{Y > y\}$ for large y , we need some regularity on the distribution of Y . Hence, let us assume that the density of Y exists and denote it by $f := (1 - \bar{F})'$. Concerning the noise, we require that there exists $q > 2$ such that

$$\mathbb{E}(\|\varepsilon\|^q) < +\infty. \quad (2.5)$$

In view of the norms injection, the noise ε also satisfies $\mathbb{E}(\|\varepsilon\|) < +\infty$. By ([Hytönen et al., 2016](#), Proposition 1.2.2), since ε is strongly \mathbb{P} -measurable as a random variable, it is also Bochner-integrable.

3. INFERENCE AND CONSISTENCY

Let $(X_i, Y_i)_{1 \leq i \leq n}$ be independent copies of (X, Y) with $X_i \in H$ and $Y_i \in \mathbb{R}$. Let also $y_n \rightarrow +\infty$ be a deterministic sequence such that

$$n\bar{F}(y_n) \xrightarrow{n \rightarrow +\infty} +\infty. \quad (3.1)$$

The empirical counterpart of the tail-moment $m_W(y)$ is defined by

$$\hat{m}_W(y_n) := \frac{1}{n} \sum_{i=1}^n W_i 1_{\{Y_i \geq y_n\}}$$

basing on the n -sample. It is then possible to estimate $w(y_n)$ by $\hat{w}(y_n)$, obtained by replacing all four tail moments in (2.2) by their empirical counterparts. Note that $\hat{m}_Y(y_n)$ is sometimes referred to as the empirical Conditional Tail Expectation (see [El-Methni et al., 2014](#)). It readily appears from (2.2) that $w(y_n)$ is a linear combination of $m_{XY}(y_n) \in H$ and $m_X(y_n) \in H$. Since both of them point in the same direction in H which is given by X , it seems reasonable to investigate the asymptotic behaviour of more general empirical tail moments defined as

$$\hat{v}_\varphi(y_n) = \hat{m}_{\varphi(Y)X}(y_n) = \frac{1}{n} \sum_{i=1}^n \varphi(Y_i) X_i 1_{\{Y_i \geq y_n\}}, \quad \hat{\beta}_\varphi(y_n) := \frac{\hat{v}_\varphi(y_n)}{\|\hat{v}_\varphi(y_n)\|}.$$

where $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is some test function such that $\varphi \in \mathcal{RV}_\tau(+\infty)$ with $\tau \in \mathbb{R}$.

We may now state our consistency result under the inverse single-index model. From now on, we adopt the notation $a_n \ll b_n$ for two real sequences if and only if $a_n/b_n \rightarrow 0$ as $n \rightarrow +\infty$ and $b_n \neq 0$ eventually.

Proposition 3.1. *Suppose (2.3), (2.4) and (2.5) hold. Let (y_n) be a deterministic sequence such that $y_n \rightarrow +\infty$ and (3.1) holds. Assume moreover that*

$$2(\kappa + \tau)\gamma < 1, \quad (3.2)$$

$$q\kappa\gamma > 1. \quad (3.3)$$

If $\tau > 0$, assume in addition that

$$q(1 - 2\tau\gamma) > 2. \quad (3.4)$$

Let $\delta_n := (g(y_n)\bar{F}^{1/q}(y_n))^{-1} \ll 1$. Then,

$$\|\hat{\beta}_\varphi(y_n) - \beta\| = O_{\mathbb{P}}(\delta_n).$$

In the next result, we show that the same amount of the tail information is asymptotically preserved when projecting on $\text{Span}(\hat{\beta}_\varphi(y_n))$ instead of $\text{Span}(\beta)$.

Proposition 3.2. *Let (X_0, Y_0) with same distribution as (X, Y) and independent of $(X_i, Y_i)_i$. Under the conditions of Proposition 3.1, it holds that, as $n \rightarrow +\infty$,*

$$\left| \frac{\text{Cov}(Y_0, \langle \hat{\beta}_\varphi(y_n), X_0 \rangle \mid Y_0 \geq y_n)}{\text{Cov}(Y_0, \langle \beta, X_0 \rangle \mid Y_0 \geq y_n)} - 1 \right| = O(\delta_n),$$

where it is recalled that $\delta_n = (g(y_n)\bar{F}^{1/q}(y_n))^{-1} \ll 1$.

Remark 3.1. We state some comments about the set of hypotheses and their implications.

- The assumptions may be divided into two classes. First, there are the model hypotheses which are constituted by (2.3), (2.4), (2.5) and (3.3). Second, the hypotheses related to the estimation are (3.1), (3.2), and (3.4).
- The regime (3.1) entails that the average number of tail observations $Y_i > y_n$ increases as the sample size grows. This roughly ensures that the threshold y_n does not grow too fast so we have enough data points in our inference scheme. In particular, rephrasing (3.1) and using (2.3) in terms of regular variations yield $y_n \ll n^{\gamma-\eta}$ for any small $\eta > 0$.
- The set of $(\gamma, \tau, \kappa, q)$ with $\gamma \in (0, 1)$, $\kappa > 0$, $0 < \tau < \frac{1}{2\gamma}$ and $q > 2$ such that (3.2), (3.3) and (3.4) hold is non-empty.
- The convergence rate δ_n only depends on the tail distribution of Y , the integrability order of the noise ε and the asymptotic behaviour of the deterministic link function g .

4. ILLUSTRATION ON SIMULATED DATA

In this section, we assess the validity of our method by a Monte Carlo simulation experiment with $N = 500$ independent repetitions. The ambient space is $H = L^2([0, 1])$ endowed with its usual inner product.

Throughout this numerical study, all functions are discretized on a regular grid $0 = x_1 < \dots < x_d = 1$ with $d = 101$ and the $L^2([0, 1])$ -inner product of two functions W_1, W_2 is approximated by

$$\langle W_1, W_2 \rangle_d := \frac{1}{d} \sum_{k=1}^d W_1(x_k) W_2(x_k).$$

We draw a data sample of $n = 500$ independent replications (X_i, Y_i) of (X, Y) using the following scheme.

- Y has a Burr distribution, *i.e.*, $\bar{F}(y) = (1 + y^{-\rho})^{\frac{1}{\gamma\rho}}$ with $y \geq 0$, and $\gamma \in (0, 1)$ being the tail-index and $\rho < 0$ being the second-order parameter.
- $\varepsilon \mid Y = y$ has the same distribution as $\sigma(y)B_y^H + \mu(y)$ where B_y^H is the fractional Brownian motion (fBm) on $[0, 1]$ with Hurst parameter $H(y) \in (0, 1)$, $\sigma(y)$ is the conditional noise deviation and $\mu(y)$ the conditional mean.
- $\beta : t \in [0, 1] \mapsto \sqrt{2} \sin(2\pi t)$ and $g : y \in \mathbb{R}^+ \mapsto y^\kappa$ with $\kappa \in \{1, 3/2, 2\}$.

Hence, conditionally on $Y = y$, the noise process based on B_y^H is centered Gaussian with conditional covariance function

$$(s, t) \mapsto \frac{\sigma^2(y)}{2} \left(t^{2H(y)} + s^{2H(y)} - |t - s|^{2H(y)} \right).$$

Note that $H(y) = 1/2$ gives a Brownian motion B_y with mean $\mu(y)$ and variance $\sigma^2(y)$.

Once both the response variable Y and the noise ε are simulated, and the deterministic functions β and g are chosen, the covariate sampling $(X_i)_{1 \leq i \leq n}$ is readily derived from (2.4). The choice of σ is guided by the following reasonable requirement: One should ensure that the contribution in (2.4) of ε does not overwhelm $g(Y)$, but also conversely, in order to avoid a trivial case where the noise is not impacting. To this end, we propose to pick $\sigma(y) = g(y)/10$. Moreover, we fix in our study case $(H, \mu) \equiv (1/3, 200)$ regardless of $Y = y$.

As usual in extreme value theory, we parametrize the diverging threshold by the quantiles of $(Y_i)_{1 \leq i \leq n}$. Hence, we set up y_n of the form $Y_{n-k+1:n}$ where $\min(Y_i) = Y_{1:n} \leq \dots \leq Y_{n:n} = \max(Y_i)$ are the order statistics of (Y_i) . We consider the following data-driven selection procedure on the order k , based on the maximization of the correlation between Y and $\langle \hat{\beta}_\varphi(y_n), X \rangle$ when Y is large.

For any $1 \leq k \leq n$, we introduce $\text{PCC}(k, n)$ the Pearson correlation coefficient between $(Y_{n-i+1:n})_{1 \leq i \leq k}$ and $(\langle \hat{\beta}_\varphi(Y_{n-k+1:n}), X_{(n-i+1,n)} \rangle)_{1 \leq i \leq k}$ where, for any $1 \leq i \leq k$, $X_{(n-i+1,n)}$ denotes the concomitant of $Y_{n-i+1:n}$, *i.e.*, the value of X_s with $s \in \{1, \dots, n\}$ being the unique index such that $Y_s = Y_{n-i+1:n}$ (since Y is continuous, there is no tie). The measure of association $\text{PCC}(j, n)$ is explicitly written as

$$\frac{1}{k} \sum_{i=1}^k Y_{n-i+1,n} \langle \hat{\beta}_\varphi(Y_{n-k+1:n}), X_{(n-i+1,n)} \rangle - \frac{1}{k} \sum_{i=1}^k Y_{n-i+1,n} \frac{1}{k} \sum_{i=1}^k \langle \hat{\beta}_\varphi(Y_{n-k+1:n}), X_{(n-i+1,n)} \rangle.$$

We select the order k defined as

$$k := \arg \max_{5 \leq j \leq n} \text{PCC}(j, n).$$

Under the described simulation model, we compute the values of $\text{PCC}(k, n)$ for $1 \leq k \leq 100$ and $n = 500$ for different admissible τ . The results are represented in Figure 1. As one may see, the estimator should be the most performing at the highest admissible value of τ where it always capture the most of the correlation in the extremes between Y and $\langle \hat{\beta}_\varphi, X \rangle$.

Using the selection method for k , we may now set the threshold y_n according to the data and infer the single-index parameter β thanks to $\hat{\beta}_\varphi$. The results are shown in Figure 2 for multiple tail indexes. One may observe that the estimation is more accurate when γ or (mostly) τ increases.




FIGURE 1. Finite sample behaviour comparison of $\hat{\beta}_\varphi(Y_{n-k-1:n})$ for different admissible τ under the inverse model with Burr response and conditional fBm noise. The x -axis represents the number k of exceedances and the y -axis is the average value over $N = 500$ Monte Carlo repetitions of the correlation between $(Y_{n-i+1:n})_{1 \leq i \leq k}$ et $(\langle \hat{\beta}_\varphi(Y_{n-k+1:n}), X_{(n-i+1,n)} \rangle)_{1 \leq i \leq k}$. Here, the tail indexes κ and γ are varying and $\rho = -1/2$.




FIGURE 2. Simulation results on the inverse model with Burr response and conditional fBm noise. The orange curve is the graph of β on $[0, 1]$ while the blue one is the averaged value of $\hat{\beta}_\varphi(Y_{n-k+1:n})$ over $N = 500$ Monte Carlo repetitions, where $k = \arg \max_{5 \leq j \leq n} \text{PCC}(j, n)$. The light blue area corresponds to the confidence region comprising the top 5 – 95% values of the Monte Carlo repetitions. Here, the tail indexes κ and γ are varying while we fix $\rho = -1/2$ and $\tau = -2$.

5. TECHNICAL RESULTS

We gather in this section all the mathematical details behind Section 2 and Section 3. We start with Section 5.1 and an useful technical tool which, among other things, enables the proof of Proposition 2.1. We also give a result on the asymptotic equivalent of theoretical tail-moments, which will be extensively used throughout the paper. Next, we derive a central limit theorem on the univariate empirical tail-moments in Section 5.2. The relevant asymptotics of empirical tail-moments involving the noise and its inner product with β are respectively investigated in Section 5.3 and Section 5.4. Finally, we establish our main consistency results in Section 5.5.

5.1. Preliminaries. A standard result in the literature of Bochner spaces, *e.g.*, (Hytönen et al., 2016, Eq (1.2)), states that Bochner integrals commute with bounded linear operators such as orthogonal projections. Note that this is in fact the definition of Pettis integrals. In the Hilbert framework, projections are generally expressed in terms of the inner product. The following technical tool considers this commutation operation with the inner product when the two random variables are stochastically independent. The case of one variable being deterministic will also often occur in the proofs.

Lemma 5.1. *For any W_1, W_2 independent Bochner-integrable random variables having values in $(H, \langle \cdot, \cdot \rangle)$ a separable Hilbert space,*

$$\mathbb{E}(\langle W_1, W_2 \rangle) = \langle \mathbb{E}_{\mathcal{B}}(W_1), \mathbb{E}_{\mathcal{B}}(W_2) \rangle.$$

Proof. Let $i \in \{1, 2\}$. We start by observing that $\mathbb{E}(\|W_i\|) < +\infty$ in view of (Hytönen et al., 2016, Proposition 1.2.2). Since H is separable, it admits an orthonormal countable basis, *i.e.*, there exists $(e_n)_{n \geq 1} \in H$ such that $\|e_n\| = 1$ for any $n \geq 1$ and $W_i = \sum_{n \geq 1} \langle W_i, e_n \rangle e_n$ as well as $\|W_i\| = \|\langle W_i, e_n \rangle\|_{\ell^2(\mathbb{N})}$. Accordingly, $\mathbb{E}(\|W_i\|) < +\infty$ entails $\mathbb{E}(\sum_{n \geq 1} \langle W_i, e_n \rangle^2) < +\infty$ which is the same as $\|\mathbb{E}(\langle W_i, e_n \rangle^2)\|_{\ell^1(\mathbb{N})} = \sum_{n \geq 1} \mathbb{E}(\langle W_i, e_n \rangle^2) < +\infty$ by a straightforward application of the standard Fubini's theorem. Using norm injections, we conclude that $\|\mathbb{E} \langle W_i, e_n \rangle\|_{\ell^1(\mathbb{N})} \leq \sum_{n \geq 1} \mathbb{E} |\langle W_i, e_n \rangle| < +\infty$. We may now invoke Fubini's theorem (Hytönen et al., 2016, Proposition 1.2.7) for Bochner integrals to decompose

$$\mathbb{E}_{\mathcal{B}}(W_i) = \sum_{n \geq 1} \mathbb{E}(\langle W_i, e_n \rangle) e_n.$$

Now, $w \mapsto \langle w, z \rangle$ is a bounded linear operator for any fixed $z \in H$ and hence Hille's theorem (Hytönen et al., 2016, Theorem 1.2.4) (which is more than what we actually need) yields $\mathbb{E} \langle W_i, e_n \rangle = \langle \mathbb{E}_{\mathcal{B}}(W_i), e_n \rangle$ which in turn gives

$$\mathbb{E}_{\mathcal{B}}(W_i) = \sum_{n \geq 1} \langle \mathbb{E}_{\mathcal{B}}(W_i), e_n \rangle e_n.$$

Overall,

$$\langle \mathbb{E}_{\mathcal{B}}(W_1), \mathbb{E}_{\mathcal{B}}(W_2) \rangle = \sum_{n \geq 1} \langle \mathbb{E}_{\mathcal{B}}(W_1), e_n \rangle \langle \mathbb{E}_{\mathcal{B}}(W_2), e_n \rangle$$

holds by continuity of the inner product on H^2 . Commuting again the Bochner expectation with the inner product and independence, it follows that

$$\langle \mathbb{E}_{\mathcal{B}}(W_1), \mathbb{E}_{\mathcal{B}}(W_2) \rangle = \sum_{n \geq 1} \mathbb{E}(\langle W_1, e_n \rangle \langle W_2, e_n \rangle).$$

The last step is to exchange the standard Lebesgue expectation and the infinite sum in the RHS of the last equality. This would conclude the proof as it implies that

$$\langle \mathbb{E}_{\mathcal{B}}(W_1), \mathbb{E}_{\mathcal{B}}(W_2) \rangle = \mathbb{E}(\sum_{n \geq 1} \langle W_1, e_n \rangle \langle W_2, e_n \rangle) = \mathbb{E}(\langle W_1, W_2 \rangle).$$

Whence, we seek to apply the standard Fubini's theorem and we thus need to show that $\sum_{n \geq 1} \mathbb{E} |\langle W_1, e_n \rangle \langle W_2, e_n \rangle| < +\infty$. Using independence, Cauchy-Schwarz's inequality in $\ell^2(\mathbb{N})$, Jensen's inequality and the integrability of W_1, W_2 , one may write

$$\sum_{n \geq 1} \mathbb{E} |\langle W_1, e_n \rangle \langle W_2, e_n \rangle| = \sum_{n \geq 1} \mathbb{E} |\langle W_1, e_n \rangle| \mathbb{E} |\langle W_2, e_n \rangle|$$

and bound this quantity by

$$\|\mathbb{E}(|\langle W_1, e_n \rangle|)\|_{\ell^2(\mathbb{N})} \|\mathbb{E}(|\langle W_2, e_n \rangle|)\|_{\ell^2(\mathbb{N})} \leq \|\mathbb{E}(\langle W_1, e_n \rangle^2)\|_{\ell^1(\mathbb{N})}^{1/2} \|\mathbb{E}(\langle W_2, e_n \rangle^2)\|_{\ell^1(\mathbb{N})}^{1/2} < +\infty$$

which ends the proof. \square

We may now establish the explicit solution of the optimization problem (2.1) under the inverse single-index model (2.4). The following proof is a simpler version of (Bousebata et al., 2023, Proposition 1) but it extends to the functional case.

Proof of Proposition 2.1. We start by rewriting the conditional covariance. For any $y \in \mathbb{R}$,

$$\text{Cov}(\langle w, X \rangle, Y | Y \geq y) = \frac{\mathbb{E}(\langle w, X \rangle Y \mathbf{1}_{\{Y \geq y\}})}{\bar{F}(y)} - \frac{\mathbb{E}(\langle w, X \rangle \mathbf{1}_{\{Y \geq y\}}) \mathbb{E}(Y \mathbf{1}_{\{Y \geq y\}})}{\bar{F}(y)^2}.$$

By (Hytönen et al., 2016, Proposition 1.2.2), we have $\mathbb{E}(\|X\|) < +\infty$. Since $\|w\| = 1$, a straightforward application of Cauchy-Schwarz's inequality and monotony of expectations imply that the expectations in the last display are finite. Next, by Lemma 5.1, one has

$$\text{Cov}(\langle w, X \rangle, Y | Y \geq y) = \bar{F}(y) \langle w, m_{XY}(y) \rangle - \langle w, m_Y(y) m_X(y) \rangle = \langle w, v(y) \rangle.$$

The optimization problem now becomes $\arg \max_{\|w\|=1} \langle w, v(y) \rangle$ for which the solution is known to be $v(y)/\|v(y)\|$. Indeed, for any $w \in H$ such that $\|w\| = 1$, Cauchy-Schwarz's inequality implies that $\langle w, v(y) \rangle \leq \|v(y)\|$, while taking $w = v(y)/\|v(y)\|$ realizes the latter upper-bound. \square

The next result is a slight extension to (Bousebata et al., 2023, Lemma 2) that we quote here for the sake for self-containedness.

Lemma 5.2. *Let any $h \in \mathcal{RV}_\rho(+\infty)$ with $\rho \in \mathbb{R}$ and any real random variable Y with density $f \in \mathcal{RV}_{-1/\gamma-1}(+\infty)$ with $\gamma > 0$. Assume that $\rho\gamma < 1$. Then, as $y \rightarrow +\infty$,*

$$m_{h(Y)}(y) := \mathbb{E}(h(Y) \mathbf{1}_{\{Y \geq y\}}) \sim \frac{1}{1 - \rho\gamma} h(y) \bar{F}(y),$$

where $\bar{F}(y) := 1 - \int_{-\infty}^y f(t) dt \in \mathcal{RV}_{-1/\gamma}(+\infty)$ is the survival distribution function of Y .

Proof. By transfer formula, one may write

$$\mathbb{E}(h(Y) \mathbf{1}_{\{Y \geq y\}}) = \int_y^{+\infty} h(t) f(t) dt.$$

Since $h \cdot f \in \mathcal{RV}_{\rho-1-1/\gamma}(+\infty)$, we may write $h(x) f(x) = x^{\rho-1-1/\gamma} \ell(x)$ for any x where ℓ is slowly-varying. Now, since $\rho\gamma < 1$ ensures that $\rho - 1 - 1/\gamma < -1$, one may apply (Bingham et al., 1989, Proposition 1.5.10) to get

$$\int_y^{+\infty} h(t) f(t) dt \sim \frac{1}{-\rho + \frac{1}{\gamma}} y^{\rho-1/\gamma} \ell(y) = \frac{1}{-\rho + \frac{1}{\gamma}} y h(y) f(y), \quad y \rightarrow +\infty.$$

Since $\gamma > 0$ so that $-1 - 1/\gamma < -1$ and since $f \in \mathcal{RV}_{-1-1/\gamma}(+\infty)$, we may apply once more (Bingham et al., 1989, Proposition 1.5.10) to get $y f(y) \sim \bar{F}(y)/\gamma$ as $y \rightarrow +\infty$. The conclusion follows by combining the previous asymptotics and transitivity of equivalents. \square

5.2. Empirical tail-moment of $g(Y)$.

Lemma 5.3. *Let $\varphi \in \mathcal{RV}_\tau(+\infty)$ with $\tau \in \mathbb{R}$ and $g \in \mathcal{RV}_\kappa(+\infty)$ with $\kappa > 0$. Let Y a real random variable with survival function such that (2.3) holds. Let (y_n) be a deterministic sequence such that (3.1) holds. Assume that (3.2) holds for $\tau \in \mathbb{R}$. Then,*

$$\sqrt{n\bar{F}(y_n)} \left(\frac{\hat{m}_{\varphi \cdot g(Y)}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} - 1 \right) \xrightarrow[n \rightarrow +\infty]{d} \mathcal{N} \left(0, \frac{(1 - (\kappa + \tau)\gamma)^2}{1 - 2(\kappa + \tau)\gamma} \right). \quad (5.1)$$

Proof. Let $D_{ni} := \frac{\varphi(Y_i)g(Y_i)}{m_{\varphi \cdot g(Y)}(y_n)} 1_{\{Y_i \geq y_n\}} - 1$ and $s_n^2 := \sum_{i=1}^n \text{Var}(D_{ni}) = n \left(\frac{m_{\varphi^2 \cdot g^2(Y)}(y_n)}{m_{\varphi \cdot g(Y)}^2(y_n)} - 1 \right)$.

First, observe that Lemma 5.2 yields, under condition (3.2), the following two first-order approximations:

$$\begin{aligned} m_{\varphi \cdot g(Y)}(y_n) &= \bar{F}(y_n) \mathbb{E}(\varphi(Y)g(Y) \mid Y \geq y_n) \sim \frac{1}{1 - (\kappa + \tau)\gamma} \varphi(y_n)g(y_n)\bar{F}(y_n), \\ m_{\varphi^2 \cdot g^2(Y)}(y_n) &= \bar{F}(y_n) \mathbb{E}(\varphi^2(Y)g^2(Y) \mid Y \geq y_n) \sim \frac{1}{1 - 2(\kappa + \tau)\gamma} \varphi^2(y_n)g^2(y_n)\bar{F}(y_n), \end{aligned}$$

as $n \rightarrow +\infty$. In particular, one has

$$\frac{m_{\varphi^2 \cdot g^2(Y)}(y_n)}{m_{\varphi \cdot g(Y)}^2(y_n)} \sim \frac{(1 - (\kappa + \tau)\gamma)^2}{1 - 2(\kappa + \tau)\gamma} \frac{1}{\bar{F}(y_n)} \xrightarrow[n \rightarrow +\infty]{} +\infty,$$

and thus,

$$s_n^2 \sim \frac{(1 - (\kappa + \tau)\gamma)^2}{1 - 2(\kappa + \tau)\gamma} \frac{n}{\bar{F}(y_n)}, \quad n \rightarrow +\infty.$$

Since $(D_{ni})_{1 \leq i \leq n}$ is a centered independent triangular array, and according to (Billingsley, 1995, Theorem 27.3), it is enough to show that

$$\exists \delta > 0 \text{ such that } \sum_{i=1}^n \frac{1}{s_n^{2+\delta}} \mathbb{E}(|D_{ni}|^{2+\delta}) \xrightarrow[n \rightarrow +\infty]{} 0. \quad (5.2)$$

This would provide that $\sum_{i=1}^n D_{ni}/s_n \xrightarrow[n \rightarrow +\infty]{d} \mathcal{N}(0, 1)$ which is equivalent to (5.1). Now, let $\delta \in (0, \frac{1}{\gamma(\kappa + \tau)} - 2)$ which is possible by hypothesis (3.2). We write, for $i \in \{1, \dots, n\}$:

$$D_{ni} = \frac{1}{m_{\varphi \cdot g(Y)}(y_n)} \left(\varphi(Y_i)g(Y_i)1_{\{Y_i \geq y_n\}} - m_{\varphi \cdot g(Y)}(y_n) \right).$$

Using the triangle inequality, Jensen inequality, the fact that the L^p -norms are ordered and that regularly-varying functions are eventually positive,

$$\begin{aligned} \mathbb{E}(|D_{ni}|^{2+\delta})^{\frac{1}{2+\delta}} &\leq m_{\varphi \cdot g(Y)}(y_n)^{-1} \left[\mathbb{E} \left((\varphi(Y_i)g(Y_i))^{2+\delta} 1_{\{Y_i \geq y_n\}} \right)^{\frac{1}{2+\delta}} + m_{\varphi \cdot g(Y)}(y_n) \right] \\ &\leq m_{\varphi \cdot g(Y)}(y_n)^{-1} \left[\mathbb{E} \left((\varphi(Y_i)g(Y_i))^{2+\delta} 1_{\{Y_i \geq y_n\}} \right)^{\frac{1}{2+\delta}} + \mathbb{E}(\varphi(Y_i)g(Y_i)1_{\{Y_i \geq y_n\}}) \right] \\ &\leq 2m_{\varphi \cdot g(Y)}(y_n)^{-1} \mathbb{E} \left((\varphi(Y_i)g(Y_i))^{2+\delta} 1_{\{Y_i \geq y_n\}} \right)^{\frac{1}{2+\delta}}. \end{aligned}$$

It follows that

$$0 \leq \sum_{i=1}^n \frac{1}{s_n^{2+\delta}} \mathbb{E}(|D_{ni}|^{2+\delta}) \leq \sum_{i=1}^n \frac{2^{2+\delta}}{s_n^{2+\delta} m_{\varphi \cdot g(Y)}(y_n)} \mathbb{E} \left((\varphi(Y_i)g(Y_i))^{2+\delta} 1_{\{Y_i \geq y_n\}} \right).$$

Using equidistribution, Lemma 5.2 again with the fact that $(\varphi \cdot g)^{2+\delta} \in \mathcal{RV}_{(2+\delta)(\kappa + \tau)}(+\infty)$, there exists a constant $c \in (0, +\infty)$ depending only on $\kappa, \tau, \gamma, \delta$ such that, as $n \rightarrow +\infty$,

$$\sum_{i=1}^n \frac{1}{s_n^{2+\delta} m_{\varphi \cdot g(Y)}(y_n)} \mathbb{E} \left(|\varphi(Y_i)g(Y_i)1_{\{Y_i \geq y_n\}}|^{2+\delta} \right) = \frac{n}{s_n^{2+\delta}} \frac{m_{(\varphi \cdot g)^{2+\delta}(Y)}(y_n)}{m_{\varphi \cdot g(Y)}^{2+\delta}(y_n)} \sim c (n\bar{F}(y_n))^{-\delta/2},$$

and the conclusion follows from (3.1). \square

5.3. Empirical tail-moments for ε .

Lemma 5.4. *Assume that (2.3), (2.5), (3.2) and (3.4) hold. Let (y_n) be a deterministic sequence such that (3.1) holds and let $\delta_n := (g(y_n)\bar{F}^{1/q}(y_n))^{-1}$. Then,*

$$\frac{\|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|}{m_{\varphi \cdot g(Y)}(y_n)} = O_{\mathbb{P}}(\delta_n), \quad n \rightarrow +\infty.$$

Remark 5.5. Observe that $\delta_n \ll 1$ under assumption (3.3).

Proof. Denoting $Z := \varphi(Y)\varepsilon$ for clarity, we start by computing the expectation of $\|\hat{m}_Z(y_n)\|^2$ which is by construction,

$$\|\hat{m}_Z(y_n)\|^2 = \frac{1}{n^2} \sum_{i_1, i_2=1}^n \langle Z_{i_1}, Z_{i_2} \rangle 1_{\{Y_{i_1} \geq y_n\}} 1_{\{Y_{i_2} \geq y_n\}}.$$

Taking the expectation and splitting the sum, we may write

$$\mathbb{E}(\|\hat{m}_Z(y_n)\|^2) = \frac{1}{n^2} \sum_{i_1 \neq i_2}^n \mathbb{E} \left(\langle Z_{i_1} 1_{\{Y_{i_1} \geq y_n\}}, Z_{i_2} 1_{\{Y_{i_2} \geq y_n\}} \rangle \right) + \frac{1}{n^2} \sum_{i=1}^n \mathbb{E} \left(\|Z_i\|^2 1_{\{Y_i \geq y_n\}} \right).$$

Lemma 5.1 and equidistribution entail

$$\mathbb{E}(\|\hat{m}_Z(y_n)\|^2) = \|m_Z(y_n)\|^2 + \frac{1}{n} \left(m_{\|Z\|^2}(y_n) - \|m_Z(y_n)\|^2 \right).$$

The main tool of the proof is Markov's inequality which gives, for any $\eta' > 0$,

$$\mathbb{P} \left(\frac{\|\hat{m}_Z(y_n)\|}{m_{\varphi \cdot g(Y)}(y_n)} > \eta' \right) \leq \frac{1}{\eta'^2} \mathbb{E} \left(\|\hat{m}_Z(y_n)\|^2 \right) m_{\varphi \cdot g(Y)}^{-2}(y_n). \quad (5.3)$$

According to Lemma 5.2 and under (3.2), there exists some constant $c \in (0, +\infty)$ independent of n such that, for n large enough, Markov's inequality (5.3) becomes

$$\mathbb{P} \left(\frac{\|\hat{m}_Z(y_n)\|}{m_{\varphi \cdot g(Y)}(y_n)} > \eta' \right) \leq \frac{c}{\eta'^2} (\varphi \cdot g \cdot \bar{F})^{-2}(y_n) \left(\|m_Z(y_n)\|^2 \vee \frac{1}{n} m_{\|Z\|^2}(y_n) \right).$$

The next step is to bound the term involving the tail-moments. To this end, observe that we may write by Lemma 5.1,

$$\|m_Z(y_n)\|^2 = \langle \mathbb{E}_{\mathcal{B}}(Z_1 1_{\{Y_1 \geq y_n\}}), \mathbb{E}_{\mathcal{B}}(Z_2 1_{\{Y_2 \geq y_n\}}) \rangle = \mathbb{E}(\langle Z_1, Z_2 \rangle 1_{\{Y_1, Y_2 \geq y_n\}}).$$

By independence and Cauchy-Scharwz's inequality in H and then by Cauchy-Schwarz's inequality in $L^2(\Omega, \mathcal{A}, \mathbb{P})$, we may bound

$$\|m_Z(y_n)\|^2 \leq m_{\varphi(Y)\|\varepsilon\|}^2(y_n) \leq m_{\|Z\|^2}(y_n) \bar{F}(y_n). \quad (5.4)$$

The regime (3.1) implies that there exists a constant $c \in (0, +\infty)$ independent of n such that for n large enough,

$$\frac{1}{n} m_{\|Z\|^2}(y_n) \leq c \bar{F}(y_n) m_{\|Z\|^2}(y_n).$$

It follows from (5.4) that

$$\|m_Z(y_n)\|^2 \vee \frac{1}{n} m_{\|Z\|^2}(y_n) \leq m_{\|Z\|^2}(y_n) \bar{F}(y_n). \quad (5.5)$$

The focus now turns to the tail-moment of $\|Z\|^2$ in the previous inequality. The q -moment of $\|\varepsilon\|$ is finite under (2.5) so that c 's inequality allows us to write,

$$m_{\|Z\|^2}(y_n) = \mathbb{E} \left(\varphi^2(Y) \|\varepsilon\|^2 1_{\{Y \geq y_n\}} \right) \leq \mathbb{E} (\|\varepsilon\|^q)^{2/q} m_{\varphi(Y) \frac{2q}{q-2}}^{1-2/q}(y_n).$$

Note that the tail moment $m_{\varphi(Y)^{\frac{2q}{q-2}}}(y_n)$ is finite if $\frac{2\tau q}{q-2} < \frac{1}{\gamma}$ which is satisfied for any $q > 2$ and $\gamma \in (0, 1)$ whenever $\tau \leq 0$; This existence condition is also satisfied when $\tau > 0$ under (3.4). Moreover, $\varphi^{\frac{2q}{q-2}} \in \mathcal{RV}_{\frac{2\tau q}{q-2}}(+\infty)$ so that Lemma 5.2 and (3.4) yield

$$m_{\varphi(Y)^{\frac{2q}{q-2}}}^{1-2/q}(y_n) \sim \frac{1}{(1 - \frac{2\tau q \gamma}{q-2})^{1-2/q}} \varphi^2(y_n) \bar{F}^{1-2/q}(y_n).$$

It follows that there exists a constant $c \in (0, +\infty)$ independent of n such that for n large enough,

$$m_{\|Z\|^2}(y_n) \leq \mathbb{E}(\|\varepsilon\|^q)^{2/q} m_{\varphi(Y)^{\frac{2q}{q-2}}}^{1-2/q}(y_n) \leq c \varphi^2(y_n) \bar{F}^{1-2/q}(y_n). \quad (5.6)$$

Combining (5.5) and (5.6), we have for some constant $c \in (0, +\infty)$ independent of n ,

$$\|m_Z(y_n)\|^2 \vee \frac{1}{n} m_{\|Z\|^2}(y_n) \leq c \varphi^2(y_n) \bar{F}^{2(1-1/q)}(y_n). \quad (5.7)$$

Going back to Markov's bound (5.3), for some constant $c \in (0, +\infty)$ independent of n ,

$$\mathbb{P}\left(\frac{\|\hat{m}_Z(y_n)\|}{m_{\varphi \cdot g(Y)}(y_n)} > \eta'\right) \leq \frac{c}{\eta'^2} g^{-2}(y_n) \bar{F}^{-2/q}(y_n) = \frac{c \delta_n^2}{\eta'^2}.$$

Now, let any $\eta > 0$. Since η' is arbitrary, we may pick η' of the form $\delta_n(c/\eta)^{-1/2}$ so that it concludes the proof as we have shown that

$$\mathbb{P}\left(\delta_n^{-1} \frac{\|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|}{m_{\varphi \cdot g(Y)}(y_n)} > \eta^{-1/2}\right) < \eta,$$

for all $\eta > 0$. □

5.4. Empirical tail-moment of the inner product.

Lemma 5.6. *Let any $\beta \in H$ with $\|\beta\| = 1$. Assume that (2.3), (2.5), (3.2) and (3.4) hold. Let (y_n) be a deterministic sequence such that (3.1) holds and let $\delta_n := (g(y_n) \bar{F}^{1/q}(y_n))^{-1}$. Then,*

$$\frac{\hat{m}_{\langle \beta, \varphi(Y)\varepsilon \rangle}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} = O_{\mathbb{P}}(\delta_n), \quad n \rightarrow +\infty.$$

Proof. We start with a preliminary computation of the expectation. One has

$$\hat{m}_{\langle \beta, \varphi(Y)\varepsilon \rangle}(y_n) = \frac{1}{n} \sum_{i=1}^n \langle \beta, \varphi(Y_i)\varepsilon_i \rangle 1_{\{Y_i \geq y_n\}}.$$

Since $\langle \beta, \cdot \rangle : H \rightarrow \mathbb{R}$ is a continuous linear mapping,

$$\mathbb{E}(\hat{m}_{\langle \beta, \varphi(Y)\varepsilon \rangle}(y_n)) = \frac{1}{n} \sum_{i=1}^n \langle \beta, m_{\varphi(Y_i)\varepsilon_i}(y_n) \rangle = \langle \beta, m_{\varphi(Y)\varepsilon}(y_n) \rangle.$$

We may now invoke the main tool for the proof, namely the Chebyshev's inequality which states that

$$\mathbb{P}\left(\left|\frac{\hat{m}_{\langle \beta, \varphi(Y)\varepsilon \rangle}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} - \frac{\langle \beta, m_{\varphi(Y)\varepsilon}(y_n) \rangle}{m_{\varphi \cdot g(Y)}(y_n)}\right| > \eta'\right) \leq \frac{\text{Var}(\hat{m}_{\langle \beta, \varphi(Y)\varepsilon \rangle}(y_n))}{m_{\varphi \cdot g(Y)}^2(y_n) \eta'^2}, \quad \forall \eta' > 0. \quad (5.8)$$

The first step is to show that the RHS in (5.8) converges to zero for large n . Focusing on the variance, it is straightforward to see that, by Lemma 5.1, independence and equidistribution,

$$\text{Var}(\hat{m}_{\langle \beta, \varphi(Y)\varepsilon \rangle}(y_n)) = \frac{1}{n} \text{Var}(\langle \beta, \varphi(Y)\varepsilon 1_{\{Y \geq y_n\}} \rangle).$$

By positivity, one has

$$\text{Var}(\langle \beta, \varphi(Y)\varepsilon 1_{\{Y \geq y_n\}} \rangle) \leq \mathbb{E}(\langle \beta, \varphi(Y)\varepsilon 1_{\{Y \geq y_n\}} \rangle^2) = \mathbb{E}(\varphi^2(Y) 1_{\{Y \geq y_n\}} \langle \beta, \varepsilon \rangle^2).$$

Now, Cauchy-Schwarz's inequality in H entails

$$\text{Var}(\hat{m}_{\langle\beta,\varepsilon\rangle}(y_n)) \leq \frac{1}{n} \mathbb{E} \left(\varphi^2(Y) 1_{\{Y \geq y_n\}} \|\varepsilon\|^2 \right) = \frac{1}{n} m_{\varphi^2(Y)\|\varepsilon\|^2}(y_n).$$

Next, using the bound (5.6) in the proof of Lemma 5.4, which is allowed since (2.5) and (3.4) hold, there exists a constant $c \in (0, +\infty)$ independent of n such that for n large enough,

$$\text{Var}(\hat{m}_{\langle\beta,\varepsilon\rangle}(y_n)) \leq cn^{-1} \varphi^2(y_n) \bar{F}^{1-2/q}(y_n).$$

The denominator of the RHS in (5.8) is controlled with Lemma 5.2:

$$m_{\varphi \cdot g(Y)}^2(y_n) \sim \frac{1}{(1 - (\kappa + \tau)\gamma)^2} \varphi^2(y_n) g^2(y_n) \bar{F}^2(y_n) \text{ as } n \rightarrow +\infty.$$

Overall, the Chebychev's inequality (5.8) becomes, for some constant $c \in (0, +\infty)$ independent of n , for any n large enough and any $\eta' > 0$,

$$\mathbb{P} \left(\left| \frac{\hat{m}_{\langle\beta,\varphi(Y)\varepsilon\rangle}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} - \frac{\langle\beta, m_{\varphi(Y)\varepsilon}(y_n)\rangle}{m_{\varphi \cdot g(Y)}(y_n)} \right| > \eta' \right) \leq \frac{c}{\eta'^2} (n \bar{F}(y_n) \delta_n)^{-1}.$$

Now, since η' is arbitrary, we may pick η' of the form $\delta_n (n \bar{F}(y_n) \eta/c)^{-1/2}$ for any $\eta > 0$ leading to

$$\frac{\hat{m}_{\langle\beta,\varphi(Y)\varepsilon\rangle}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} = \frac{\langle\beta, m_{\varphi(Y)\varepsilon}(y_n)\rangle}{m_{\varphi \cdot g(Y)}(y_n)} + O_{\mathbb{P}} \left(\delta_n (n \bar{F}(y_n))^{-1/2} \right).$$

The next step is to evaluate the first term of the RHS in the previous asymptotic expansion. While the denominator is controlled by Lemma 5.2, the numerator is bounded using Cauchy-Schwarz's inequality in H and then (5.7) from the proof of Lemma 5.4. This chain of arguments gives:

$$\left| \frac{\langle\beta, m_{\varphi(Y)\varepsilon}(y_n)\rangle}{m_{\varphi \cdot g(Y)}(y_n)} \right| \leq \frac{m^{1-1/q} \varphi^{q/q-1}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} \leq c \left(g(y_n) \bar{F}^{1/q}(y_n) \right)^{-1} = c \delta_n.$$

Finally, combining the previous bound with the regime (3.1) and the fact that we derived

$$\frac{\hat{m}_{\langle\beta,\varphi(Y)\varepsilon\rangle}(y_n)}{m_{\varphi \cdot g(Y)}(y_n)} = O_{\mathbb{P}} \left(\delta_n \vee \delta_n (n \bar{F}(y_n))^{-1/2} \right) = O_{\mathbb{P}}(\delta_n),$$

the conclusion follows. \square

5.5. Consistency of $\hat{\beta}_{\varphi}$.

Proof of Proposition 3.1. Under (2.4), the estimator \hat{v}_{φ} may be expressed as,

$$\hat{v}_{\varphi}(y_n) = \frac{1}{n} \sum_{i=1}^n \varphi(Y_i) g(Y_i) \beta 1_{\{Y_i \geq y_n\}} + \frac{1}{n} \sum_{i=1}^n \varphi(Y_i) \varepsilon_i 1_{\{Y_i \geq y_n\}} = \hat{m}_{\varphi g(Y)}(y_n) \beta + \hat{m}_{\varphi(Y)\varepsilon}(y_n).$$

So that, since $\|\beta\| = 1$, one may write the inner product between β and $\hat{v}_{\varphi}(y_n)$ as

$$\langle \hat{v}_{\varphi}(y_n), \beta \rangle = \hat{m}_{\varphi g(Y)}(y_n) + \hat{m}_{\langle\varphi(Y)\varepsilon,\beta\rangle}(y_n),$$

and $\|\hat{v}_{\varphi}(y_n)\|^2 = \hat{m}_{\varphi g(Y)}^2(y_n) + \|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|^2 + 2\hat{m}_{\varphi g(Y)}(y_n) \hat{m}_{\langle\varphi(Y)\varepsilon,\beta\rangle}(y_n)$. It follows that

$$\begin{aligned} 1 - \frac{\langle \hat{v}_{\varphi}(y_n), \beta \rangle^2}{\|\hat{v}_{\varphi}(y_n)\|^2} &= \frac{\|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|^2 - \hat{m}_{\langle\varphi(Y)\varepsilon,\beta\rangle}^2(y_n)}{\hat{m}_{\varphi g(Y)}^2(y_n) + 2\hat{m}_{\varphi g(Y)}(y_n) \hat{m}_{\langle\varphi(Y)\varepsilon,\beta\rangle}(y_n) + \|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|^2} \\ &= \frac{(\|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|/\hat{m}_{\varphi g(Y)}(y_n))^2 - (\hat{m}_{\langle\varphi(Y)\varepsilon,\beta\rangle}(y_n)/\hat{m}_{\varphi g(Y)}(y_n))^2}{1 + 2(\hat{m}_{\langle\varphi(Y)\varepsilon,\beta\rangle}(y_n)/\hat{m}_{\varphi g(Y)}(y_n)) + (\|\hat{m}_{\varphi(Y)\varepsilon}(y_n)\|/\hat{m}_{\varphi g(Y)}(y_n))^2}. \end{aligned}$$

In view of Lemma 5.3, we may write after a first order Taylor expansion, with respect to the convergence in probability,

$$1 - \frac{\langle \hat{v}_\varphi(y_n), \beta \rangle^2}{\|\hat{v}_\varphi(y_n)\|^2} \stackrel{\mathbb{P}}{\sim} \frac{(\|\hat{m}_{\langle \varphi(Y)\varepsilon} \rangle(y_n)\|/m_{\varphi \cdot g(Y)}(y_n))^2 - (\hat{m}_{\langle \varphi(Y)\varepsilon, \beta \rangle}(y_n)/m_{\varphi \cdot g(Y)}(y_n))^2}{1 + 2(\hat{m}_{\langle \varphi(Y)\varepsilon, \beta \rangle}(y_n)/m_{\varphi \cdot g(Y)}(y_n)) + (\|\hat{m}_{\langle \varphi(Y)\varepsilon} \rangle(y_n)\|/m_{\varphi \cdot g(Y)}(y_n))^2}.$$

The conditions to apply Lemma 5.4 and Lemma 5.6 are fulfilled so that a Taylor expansion yields

$$1 - \frac{\langle \hat{v}_\varphi(y_n), \beta \rangle^2}{\|\hat{v}_\varphi(y_n)\|^2} = O_{\mathbb{P}}(\delta_n^2) \xrightarrow{n \rightarrow +\infty} 0.$$

Finally, note that $\|\hat{\beta}_\varphi(y_n) - \beta\|^2 = 2(1 - \langle \hat{v}_\varphi(y_n), \beta \rangle^2 / \|\hat{v}_\varphi(y_n)\|^2)$ to conclude the proof. \square

Proof of Proposition 3.2. To ease the notations, let us introduce

$$\begin{aligned} \hat{C}_n &:= \text{Cov}(Y_0, \langle \hat{\beta}_\varphi(y_n), X_0 \rangle \mid Y_0 \geq y_n), \\ C_n &:= \text{Cov}(Y_0, \langle \beta, X_0 \rangle \mid Y_0 \geq y_n). \end{aligned}$$

Under the model (2.4), one may decompose

$$\begin{aligned} \hat{C}_n &= \text{Cov}(Y_0, g(Y_0) \langle \hat{\beta}_\varphi(y_n), \beta \rangle \mid Y_0 \geq y_n) + \text{Cov}(Y_0, \langle \hat{\beta}_\varphi(y_n), \varepsilon \rangle \mid Y_0 \geq y_n) \\ &= \mathbb{E}(Y_0 g(Y_0) \langle \hat{\beta}_\varphi(y_n), \beta \rangle \mid Y_0 \geq y_n) - \mathbb{E}(Y_0 \mid Y_0 \geq y_n) \mathbb{E}(g(Y_0) \langle \hat{\beta}_\varphi(y_n), \beta \rangle \mid Y_0 \geq y_n) \\ &\quad + \mathbb{E}(Y_0 \langle \hat{\beta}_\varphi(y_n), \varepsilon_0 \rangle \mid Y_0 \geq y_n) - \mathbb{E}(Y_0 \mid Y_0 \geq y_n) \mathbb{E}(\langle \varepsilon_0, \hat{\beta}_\varphi(y_n) \rangle \mid Y_0 \geq y_n), \end{aligned}$$

and similarly

$$\begin{aligned} C_n &= \text{Cov}(Y_0, g(Y_0) \mid Y_0 \geq y_n) + \text{Cov}(Y_0, \langle \beta, \varepsilon_0 \rangle \mid Y_0 \geq y_n) \\ &= \mathbb{E}(Y_0 g(Y_0) \mid Y_0 \geq y_n) - \mathbb{E}(Y_0 \mid Y_0 \geq y_n) \mathbb{E}(g(Y_0) \mid Y_0 \geq y_n) \\ &\quad + \mathbb{E}(Y_0 \langle \beta, \varepsilon_0 \rangle \mid Y_0 \geq y_n) - \mathbb{E}(Y_0 \mid Y_0 \geq y_n) \mathbb{E}(\langle \beta, \varepsilon_0 \rangle \mid Y_0 \geq y_n). \end{aligned}$$

Taking the difference yields $\hat{C}_n - C_n = \sum_{\ell=1}^4 A_{\ell,n}$ where,

$$\begin{aligned} A_{1,n} &:= \mathbb{E}(Y_0 g(Y_0) (\langle \hat{\beta}_\varphi(y_n), \beta \rangle - 1) \mid Y_0 \geq y_n), \\ A_{2,n} &:= \mathbb{E}(Y_0 \mid Y_0 \geq y_n) \mathbb{E}(g(Y_0) (1 - \langle \hat{\beta}_\varphi(y_n), \beta \rangle) \mid Y_0 \geq y_n), \\ A_{3,n} &:= \mathbb{E}(Y_0 \langle \hat{\beta}_\varphi(y_n) - \beta, \varepsilon_0 \rangle \mid Y_0 \geq y_n), \\ A_{4,n} &:= \mathbb{E}(Y_0 \mid Y_0 \geq y_n) \mathbb{E}(\langle \varepsilon_0, \beta - \hat{\beta}_\varphi(y_n) \rangle \mid Y_0 \geq y_n). \end{aligned}$$

The first step for deriving the statement is to control $\hat{C}_n - C_n$. To this end, we apply the triangular inequality which gives $|\hat{C}_n - C_n| \leq \sum_{\ell=1}^4 |A_{\ell,n}|$ and we subsequently bound each term $|A_{\ell,n}|$, $\ell = 1, \dots, 4$. Starting with $A_{1,n}$, one observes that $\hat{\beta}_\varphi(y_n)$ only depends on X_1, \dots, X_n so that, by independence, equidistribution and Jensen's inequality,

$$|A_{1,n}| \leq \frac{1}{\bar{F}(y_n)} \mathbb{E}(Y_0 g(Y_0) 1_{\{Y_0 \geq y_n\}}) \mathbb{E}(|\langle \hat{\beta}_\varphi(y_n), \beta \rangle - 1|).$$

Now, from Proposition 3.1 and a Taylor expansion of the square root, we have that $\langle \hat{\beta}_\varphi(y_n), \beta \rangle = 1 + O(\delta_n)$. Since, moreover, $|\langle \hat{\beta}_\varphi(y_n), \beta \rangle - 1| \leq 2$ holds almost surely by Cauchy-Schwarz's inequality in H , we may apply the dominated convergence theorem and Lemma 5.2, leading to $|A_{1,n}| = O(y_n g(y_n) \delta_n)$. Indeed, a general fact is that, if a generic sequence (W_n) of real random variables bounded almost surely by $M > 0$ converges to zero in probability with some rate, then it also converges in expectation to zero with the same rate. This may be derived by noticing that, for any $\eta > 0$,

$$\begin{aligned} \mathbb{E}(|W_n|) &= \int_0^{+\infty} \mathbb{P}(|W_n| \geq x) dx = \int_0^\eta \mathbb{P}(|W_n| \geq x) dx + \int_\eta^M \mathbb{P}(|W_n| \geq x) dx \\ &\leq \eta + M \mathbb{P}(|W_n| > \eta). \end{aligned}$$

Similarly, $|A_{2,n}| = O(y_n g(y_n) \delta_n)$. Focussing on the third term, we give a first bound using independence between (Y_0, ε_0) and $(X_i)_{1 \leq i \leq n}$, Lemma 5.1, Cauchy-Schwarz's inequality in H and equidistribution:

$$\begin{aligned} |A_{3,n}| &= \frac{1}{\bar{F}(y_n)} \left| \mathbb{E} \left(\langle \hat{\beta}_\varphi(y_n) - \beta, Y_0 \varepsilon_0 1_{\{Y_0 \geq y_n\}} \rangle \right) \right| = \frac{1}{\bar{F}(y_n)} \left| \langle \mathbb{E}_{\mathcal{B}}(\hat{\beta}_\varphi(y_n) - \beta), \mathbb{E}_{\mathcal{B}}(Y_0 \varepsilon_0 1_{\{Y_0 \geq y_n\}}) \rangle \right| \\ &\leq \frac{1}{\bar{F}(y_n)} \left\| \mathbb{E}_{\mathcal{B}}(Y_0 \varepsilon_0 1_{\{Y_0 \geq y_n\}}) \right\| \left\| \mathbb{E}_{\mathcal{B}}(\hat{\beta}_\varphi(y_n) - \beta) \right\|. \end{aligned}$$

Next, an application of Hölder's inequality and Lemma 5.2 yields, for some $c \in (0, +\infty)$,

$$\left\| \mathbb{E}_{\mathcal{B}}(Y_0 \varepsilon_0 1_{\{Y_0 \geq y_n\}}) \right\| \leq \mathbb{E} \left(Y_0 \|\varepsilon_0\| 1_{\{Y_0 \geq y_n\}} \right) \leq \mathbb{E} \left(\|\varepsilon_0\|^q \right)^{1/q} m_{Y_0}^{1-1/q}(y_n) \leq c y_n \bar{F}^{1-1/q}(y_n). \quad (5.9)$$

Besides, since $\hat{\beta}_\varphi(y_n)$ and β are unit vectors of H , using Jensen's inequality and the same argument based on the dominated convergence theorem altogether with Proposition 3.1, we may write

$$\begin{aligned} \left\| \mathbb{E}_{\mathcal{B}} \hat{\beta}_\varphi(y_n) - \beta \right\|^2 &\leq \mathbb{E} \left\| \hat{\beta}_\varphi(y_n) - \beta \right\|^2 = \mathbb{E} \left\| \hat{\beta}_\varphi(y_n) \right\|^2 + \mathbb{E} \|\beta\|^2 - 2 \mathbb{E} \langle \hat{\beta}_\varphi(y_n), \beta \rangle \\ &\leq 2 \left(1 - \mathbb{E} \langle \hat{\beta}_\varphi(y_n), \beta \rangle \right) = O_{\mathbb{P}}(\delta_n). \end{aligned}$$

Combining the last three inequalities, we have shown that $|A_{3,n}| = O(y_n \bar{F}^{-1/q}(y_n) \delta_n^{1/2})$. Lastly, for some constant $c \in (0, +\infty)$ independent of n and for n large enough,

$$\begin{aligned} |A_{4,n}| &= \frac{m_Y(y_n)}{\bar{F}^2(y_n)} \left| \mathbb{E} \left(\langle \hat{\beta}_\varphi(y_n) - \beta, \varepsilon_0 1_{\{Y_0 \geq y_n\}} \rangle \right) \right| = \frac{m_Y(y_n)}{\bar{F}^2(y_n)} \left| \langle \mathbb{E}_{\mathcal{B}}(\hat{\beta}_\varphi(y_n) - \beta), \mathbb{E}_{\mathcal{B}}(\varepsilon_0 1_{\{Y_0 \geq y_n\}}) \rangle \right| \\ &\leq \frac{m_Y(y_n)}{\bar{F}^2(y_n)} \left\| \mathbb{E}_{\mathcal{B}}(\varepsilon_0 1_{\{Y_0 \geq y_n\}}) \right\| \left\| \mathbb{E}_{\mathcal{B}}(\hat{\beta}_\varphi(y_n) - \beta) \right\| \leq c \frac{m_Y(y_n)}{\bar{F}^2(y_n)} \left\| \mathbb{E}_{\mathcal{B}}(\varepsilon_0 1_{\{Y_0 \geq y_n\}}) \right\| \delta_n. \end{aligned}$$

Applying Hölder's inequality once more so that, for some constant $c \in (0, +\infty)$ only depending on γ and q , we may write

$$\left\| \mathbb{E}_{\mathcal{B}}(\varepsilon_0 1_{\{Y_0 \geq y_n\}}) \right\| \leq \mathbb{E} \left(\|\varepsilon_0\| 1_{\{Y_0 \geq y_n\}} \right) \leq \mathbb{E} \left(\|\varepsilon_0\|^q \right)^{1/q} \bar{F}^{1-1/q}(y_n) \leq c \bar{F}^{1-1/q}(y_n), \quad (5.10)$$

It follows that $|A_{4,n}| = O(y_n \bar{F}^{-1/q}(y_n) \delta_n^{1/2}) = o(y_n g(y_n) \delta_n)$ under assumption (3.3). Overall, this yields

$$\left| \hat{C}_n - C_n \right| = O \left(\max_{\ell \in \{1,2,3,4\}} |A_{\ell,n}| \right) = O(y_n g(y_n) \delta_n) = O(y_n \bar{F}^{-1/q}(y_n)). \quad (5.11)$$

In a second step, we give an asymptotic equivalent of C_n and start by noticing that, by equidistribution, it may be written as,

$$C_n = \frac{m_{Yg(Y)}(y_n)}{\bar{F}(y_n)} - \frac{m_Y(y_n) m_{g(Y)}(y_n)}{\bar{F}^2(y_n)} + \frac{\mathbb{E} \left(Y \langle \beta, \varepsilon \rangle 1_{\{Y \geq y_n\}} \right)}{\bar{F}(y_n)} + \frac{\mathbb{E} \left(\langle \beta, \varepsilon \rangle 1_{\{Y \geq y_n\}} \right) m_Y(y_n)}{\bar{F}^2(y_n)}.$$

Lemma 5.2 yields the asymptotic expansion

$$\begin{aligned} C_n &= c_{\gamma,\kappa} y_n g(y_n) (1 + o(1)) + \frac{\mathbb{E} \left(Y \langle \beta, \varepsilon \rangle 1_{\{Y \geq y_n\}} \right)}{\bar{F}(y_n)} + \frac{1}{1-\gamma} \frac{\mathbb{E} \left(\langle \beta, \varepsilon \rangle 1_{\{Y \geq y_n\}} \right)}{\bar{F}(y_n)} (1 + o(1)) \\ &= c_{\gamma,\kappa} y_n g(y_n) \left[1 + \frac{\mathbb{E} \left(Y \langle \beta, \varepsilon \rangle 1_{\{Y \geq y_n\}} \right)}{c_{\gamma,\kappa} y_n g(y_n) \bar{F}(y_n)} + \frac{1}{(1-\gamma) c_{\gamma,\kappa}} \frac{\mathbb{E} \left(\langle \beta, \varepsilon \rangle 1_{\{Y \geq y_n\}} \right)}{y_n g(y_n) \bar{F}(y_n)} (1 + o(1)) + o(1) \right] \end{aligned}$$

where $c_{\gamma,\kappa} := \frac{1}{1-(c+1)\gamma} - \frac{1}{(1-\gamma)(1-c\gamma)}$. Next, combining (5.9)–(5.10) with the RHS of the previous asymptotic expansion readily implies that, as $n \rightarrow +\infty$,

$$C_n = c_{\gamma,\kappa} y_n g(y_n) [1 + O(\delta_n) + O(\delta_n/y_n) + o(1)] \sim c_{\gamma,\kappa} y_n g(y_n). \quad (5.12)$$

The final step consists in assembling the previous ingredients (5.11) and (5.12):

$$\left| \frac{\hat{C}_n}{C_n} - 1 \right| = \frac{|\hat{C}_n - C_n|}{|C_n|} = \frac{O(y_n \bar{F}^{-1/q}(y_n))}{c_{\gamma, \kappa} y_n g(y_n) (1 + o(1))} = O(\delta_n) \ll 1,$$

in view of (3.3). □

REFERENCES

- Aghbalou, A., F. Portier, A. Sabourin, and C. Zhou (2024). Tail inverse regression: Dimension reduction for prediction of extremes. *Bernoulli* 30(1), 503 – 533.
- Aguilera, A., M. Aguilera-Morillo, and C. Preda (2016). Penalized versions of functional PLS regression. *Chemometrics and Intelligent Laboratory Systems* 154, 80–92.
- Billingsley, P. (1995). *Measure and Probability (third edition)*. John Wiley & Sons, New-York.
- Bingham, N., C. Goldie, and J. Teugels (1989). *Regular Variation*. Number n° 1 in Encyclopedia of Mathematics and its Applications. Cambridge University Press.
- Bosq, D. (2000). *Linear Processes in Function Spaces: Theory and Applications*. Lecture Notes in Statistics. Springer New York.
- Bousebata, M., G. Enjolras, and S. Girard (2023). Extreme Partial Least-Squares. *Journal of Multivariate Analysis* 194, 105101.
- Chatla, S. B. and R. Liu (2024). Inverse regression for spatially distributed functional data. *arXiv:2402.03206*.
- Chen, D., P. Hall, and H.-G. Müller (2011). Single and multiple index functional regression models with nonparametric link. *The Annals of Statistics* 39(3), 1720 – 1747.
- Chen, R., S. Tian, D. Huang, Q. Lin, and J. S. Liu (2023). On the optimality of functional sliced inverse regression. *arXiv:2307.02777*.
- Chun, H. and S. Keleş (2010). Sparse partial least squares regression for simultaneous dimension reduction and variable selection. *Journal of the Royal Statistical Society Series B (Statistical Methodology)* 72(1), 3–25.
- Cook, R. (1998). *Regression Graphics: Ideas for Studying Regressions Through Graphics*. Wiley.
- Cook, R. D. (2007). Fisher lecture: Dimension reduction in regression. *Statistical Science* 22(1), 1–26.
- Cook, R. D. and L. Forzani (2018). Big data and partial least-squares prediction. *The Canadian Journal of Statistics* 46(1), 62–78.
- Cook, R. D. and L. Forzani (2019). Partial least squares prediction in high-dimensional regression. *The Annals of Statistics* 47(2), 884 – 908.
- Cook, R. D., L. Forzani, and A. F. Yao (2010). Necessary and sufficient conditions for consistency of a method for smoothed functional inverse regression. *Statistica Sinica* 20(1), 235–238.
- Dauxois, J., L. Ferré, and A.-F. Yao (2001). Un modèle semi-paramétrique pour variables aléatoires hilbertiennes. *Comptes Rendus de l'Académie des Sciences - Series I - Mathématiques* 333(10), 947–952.
- Delaigle, A. and P. Hall (2012). Methodology and theory for partial least squares applied to functional data. *The Annals of Statistics* 40(1), 322 – 352.
- El-Methni, J., L. Gardes, and S. Girard (2014). Non-parametric estimation of extreme risk measures from conditional heavy-tailed distributions. *Scandinavian Journal of Statistics* 41(4), 988–1012.
- Ferraty, F. and P. Vieu (2006). *Nonparametric Functional Data Analysis: Theory and Practice (Springer Series in Statistics)*. Berlin, Heidelberg: Springer-Verlag.
- Ferré, L. and A. F. Yao (2003). Functional sliced inverse regression analysis. *Statistics* 37(6), 475–488.
- Forzani, L. and R. D. Cook (2007). A note on smoothed functional inverse regression. *Statistica Sinica* 17(4), 1677–1681.

- Fukumizu, K., F. R. Bach, and M. I. Jordan (2009). Kernel dimension reduction in regression. *The Annals of Statistics* 37(4), 1871 – 1905.
- Gardes, L. (2018). Tail dimension reduction for extreme quantile estimation. *Extremes* 21(1), 57–95.
- Gardes, L. (2020). Nonparametric confidence intervals for conditional quantiles with large-dimensional covariates. *Electronic Journal of Statistics* 14(1), 661 – 701.
- Gardes, L. and S. Girard (2012). Functional kernel estimators of large conditional quantiles. *Electronic Journal of Statistics* 6, 1715 – 1744.
- Girard, S., G. Stupfler, and A. Usseglio-Carleve (2022). Functional estimation of extreme conditional expectiles. *Econometrics and Statistics* 21, 131–158.
- Horowitz, J. (2009, 01). *Semiparametric and Nonparametric Methods in Econometrics*, Volume 692.
- Hsing, T. and H. Ren (2009). An RKHS formulation of the inverse regression dimension-reduction problem. *The Annals of Statistics* 37(2), 726 – 755.
- Huang, D., S. Tian, and Q. Lin (2023). Sliced inverse regression with large structural dimensions. *arXiv:2305.04340*.
- Hytönen, T., J. van Neerven, M. Veraar, and L. Weis (2016). *Bochner spaces*, pp. 1–66. Cham: Springer International Publishing.
- Jiang, C.-R., W. Yu, and J.-L. Wang (2014). Inverse regression for longitudinal data. *The Annals of Statistics* 42(2), 563 – 591.
- Lee, K.-Y. and L. Li (2022). Functional sufficient dimension reduction through average Fréchet derivatives. *The Annals of Statistics* 50(2), 904 – 929.
- Li, B. and J. Song (2017). Nonlinear sufficient dimension reduction for functional data. *The Annals of Statistics* 45(3), 1059 – 1095.
- Li, B. and J. Song (2022). Dimension reduction for functional data based on weak conditional moments. *The Annals of Statistics* 50(1), 107 – 128.
- Li, K.-C. (1991). Sliced inverse regression for dimension reduction. *Journal of the American Statistical Association* 86(414), 316–327.
- Li, L., R. D. Cook, and C.-L. Tsai (2007). Partial inverse regression. *Biometrika* 94(3), 615–625.
- Li, Y. and T. Hsing (2010). Deciding the dimension of effective dimension reduction space for functional and high-dimensional data. *The Annals of Statistics* 38(5), 3028 – 3062.
- Lian, H. (2015). Functional sufficient dimension reduction: Convergence rates and multiple functional case. *Journal of Statistical Planning and Inference* 167, 58–68.
- Lin, Q., X. Li, D. Huang, and J. S. Liu (2021). On the optimality of sliced inverse regression in high dimensions. *The Annals of Statistics* 49(1), 1 – 20.
- Lin, Q., Z. Zhao, and J. S. Liu (2018). On consistency and sparsity for sliced inverse regression in high dimensions. *The Annals of Statistics* 46(2), 580 – 610.
- Martens, H. and T. Naes (1992). *Multivariate calibration*. John Wiley & Sons, New-York.
- Preda, C. and G. Saporta (2005). PLS regression on a stochastic process. *Computational Statistics & Data Analysis* 48(1), 149–158.
- Ramsay, J. O. and B. W. Silverman (2005). *Functional Data Analysis*. Springer.
- Reiss, P. T. and R. T. Ogden (2007). Functional principal component regression and functional partial least squares. *Journal of the American Statistical Association* 102(479), 984–996.
- Tan, K., L. Shi, and Z. Yu (2020). Sparse SIR: Optimal rates and adaptive estimation. *The Annals of Statistics* 48(1), 64 – 85.
- Tian, S., Z. Yu, and R. Chen (2023). Functional slicing-free inverse regression via martingale difference divergence operator. *arXiv:2307.12537*.
- Wang, G. and H. Lian (2020). Functional sliced inverse regression in a reproducing kernel hilbert space: A theoretical connection to functional linear regression. *Statistica Sinica* 30(1), 17–33.

- Wold, H. (1975). Soft modelling by latent variables: The non-linear iterative partial least squares (nipals) approach. *Journal of Applied Probability* 12(S1), 117–142.
- Xu, W., D. Li, and H. Wang (2020). Extreme quantile estimation based on the tail single-index model. *Statistica Sinica* 32, 1–22.