



HAL
open science

An analogue of Reed's conjecture for digraphs

Ken-Ichi Kawarabayashi, Lucas Picasarri-Arrieta

► **To cite this version:**

Ken-Ichi Kawarabayashi, Lucas Picasarri-Arrieta. An analogue of Reed's conjecture for digraphs. 2024. hal-04639956

HAL Id: hal-04639956

<https://inria.hal.science/hal-04639956v1>

Preprint submitted on 9 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

An analogue of Reed's conjecture for digraphs

Ken-ichi Kawarabayashi^{*,1} and Lucas Picasarri-Arrieta^{†,2}

¹National Institute of Informatics, The University of Tokyo, Japan

²Université Côte d'Azur, CNRS, Inria, I3S, Sophia Antipolis, France

Abstract

Reed in 1998 conjectured that every graph G satisfies $\chi(G) \leq \lceil \frac{\Delta(G)+1+\omega(G)}{2} \rceil$. As a partial result, he proved the existence of $\varepsilon > 0$ for which every graph G satisfies $\chi(G) \leq \lceil (1-\varepsilon)(\Delta(G)+1) + \varepsilon\omega(G) \rceil$. We propose an analogue conjecture for digraphs. Given a digraph D , we denote by $\bar{\chi}(D)$ the dichromatic number of D , which is the minimum number of colours needed to partition D into acyclic induced subdigraphs. We let $\vec{\omega}(D)$ denote the size of the largest biclique (a set of vertices inducing a complete digraph) of D and $\tilde{\Delta}(D) = \max_{v \in V(D)} \sqrt{d^+(v) \cdot d^-(v)}$. We conjecture that every digraph D satisfies $\bar{\chi}(D) \leq \lceil \frac{\tilde{\Delta}(D)+1+\vec{\omega}(D)}{2} \rceil$, which if true implies Reed's conjecture. As a partial result, we prove the existence of $\varepsilon > 0$ for which every digraph D satisfies $\bar{\chi}(D) \leq \lceil (1-\varepsilon)(\tilde{\Delta}(D)+1) + \varepsilon\vec{\omega}(D) \rceil$. This implies both Reed's result and an independent result of Harutyunyan and Mohar for oriented graphs.

To obtain this upper bound on $\bar{\chi}$, we prove that every digraph D with $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D)+1)$, where $\Delta_{\max}(D) = \max_{v \in V(D)} \max(d^+(v), d^-(v))$, admits an acyclic set of vertices intersecting each biclique of D , which generalises a result of King.

Keywords: Dichromatic number, colouring, biclique number.

1 Introduction

In 1998, Reed [30] posed the following celebrated conjecture, which draws a connection between the maximum degree $\Delta(G)$, the clique number $\omega(G)$, and the chromatic number $\chi(G)$ of a graph G .

Conjecture 1.1 (Reed [30]). *Every graph G satisfies $\chi(G) \leq \lceil \frac{\Delta(G)+1+\omega(G)}{2} \rceil$.*

As a partial result towards this conjecture, Reed proved the following.

Theorem 1.2 (Reed [30]). *There exists $\varepsilon > 0$ such that every graph G satisfies*

$$\chi(G) \leq \lceil (1-\varepsilon)(\Delta(G)+1) + \varepsilon\omega(G) \rceil.$$

Observe that Conjecture 1.1 is exactly Theorem 1.2 for $\varepsilon = \frac{1}{2}$. Restricted to graphs with sufficiently large maximum degree, Hurley *et al.* [20] proved that Theorem 1.2 holds for $\varepsilon \leq 0.199$, improving on an earlier result obtained by Bonamy *et al.* [9].

*Research supported by JSPS KAKENHI JP20A402 and 22H05001 and by JST ASPIRE JPMJAP2302.

†Research supported by the research grants DIGRAPHS ANR-19-CE48-0013 and ANR-17-EURE-0004.

The purpose of this work is to propose analogues of Conjecture 1.1 and Theorem 1.2 for digraphs. We thus need three digraph parameters that extend Δ , ω , and χ respectively. By extension, we mean that, if D is a symmetric digraph, these digraph parameters on D must coincide with Δ , ω , and χ on the underlying graph of D . One can easily check that this is actually the case for $\tilde{\Delta}$, $\tilde{\omega}$, and $\tilde{\chi}$, defined as follows.

Let D be a digraph. We let $\tilde{\Delta}(D)$ denote the maximum geometric mean of the in- and out-degrees of the vertices of D , that is $\tilde{\Delta}(D) = \max\{\sqrt{d^+(v) \cdot d^-(v)} \mid v \in V(D)\}$. A *biclique* of D is a set of vertices inducing a complete digraph. The *biclique number* $\tilde{\omega}(D)$ of D is the size of the largest biclique of D . The notions of *dicolouring* and *dichromatic number* were introduced in the late 1970s by Erdős and Neumann-Lara [27, 12]. A *k-dicolouring* of D is a function $c : V(D) \rightarrow [k]$ such that $c^{-1}(i)$ induces an acyclic subdigraph on D for each $i \in [k]$. The *dichromatic number* of D , denoted by $\tilde{\chi}(D)$, is the smallest $k \in \mathbb{N}$ such that D admits a k -dicolouring. These two concepts were rediscovered by Mohar in the 2000s [25, 8] and received a lot of attention since then. It turned out that many classical results on graph colouring can be extended to digraphs with these notions, which shows that dicolouring is somehow the right extension of proper colouring to digraphs. For instance, people generalised the Strong Perfect Graph Theorem [6], Gyárfás-Sumner's Conjecture [4, 3, 11], or Brooks-type results [17, 1, 2, 15].

Harutyunyan and Mohar [18] posed the following analogue of Conjecture 1.1 for oriented graphs, that are digraphs with biclique number at most 1.

Conjecture 1.3 (Harutyunyan and Mohar [18]). *Let D be an oriented graph, then*

$$\tilde{\chi}(D) \leq \left\lceil \frac{\tilde{\Delta}(D)}{2} \right\rceil + 1.$$

As a partial result towards their conjecture, they proved the following for $\varepsilon = e^{-13}$.

Theorem 1.4 (Harutyunyan and Mohar [18]). *There exists $\varepsilon > 0$ such that every oriented graph D with $\tilde{\Delta}(D)$ large enough satisfies $\tilde{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$.*

We actually believe that the analogue of Conjecture 1.1 holds not only for oriented graphs but for all digraphs. We thus propose the following conjecture, which if true implies both Conjectures 1.1 and 1.3.

Conjecture 1.5. *Every digraph D satisfies $\tilde{\chi}(D) \leq \left\lceil \frac{\tilde{\Delta}(D) + 1 + \tilde{\omega}(D)}{2} \right\rceil$.*

The main result of this work is the following, which implies both Theorem 1.2 (when restricted to symmetric digraphs) and Theorem 1.4 (when restricted to oriented graphs), and provide some support to Conjecture 1.5.

Theorem 1.6. *There exists $\varepsilon > 0$ such that every digraph D satisfies*

$$\tilde{\chi}(D) \leq \lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \tilde{\omega}(D) \rceil.$$

We made no attempt to maximize the value of ε in our proof. In particular, applying Theorem 1.6 to obtain Theorems 1.2 and 1.4 yields values of ε that are much smaller than the existing ones.

Our proof of Theorem 1.6 is inspired by a short proof of Theorem 1.2 due to King and Reed [23]. Along the way, we generalise to digraphs a collection of intermediate results that we find of independent interest. We refer the reader to the recent book of Stiebitz, Schweser, and Toft [31, Chapter 6] for a complete proof of Theorem 1.2 which includes proofs of all intermediate results (on undirected graphs) mentioned later on.

The paper is organised as follows. In Section 2, we give notation on digraphs and preliminary results used in the paper. In Section 3, using the probabilistic method, we prove the following upper bound on the dichromatic number of sparse digraphs, which generalises to digraphs a classical result of Molloy and Reed [26]. Given a digraph D with maximum degree $\Delta_{\max}(D) = \Delta$ and a vertex v of D , we denote by $m^+(v)$ and $m^-(v)$ the number of arcs of $D[N^+(v)]$ and $D[N^-(v)]$ respectively. We say that D is B -sparse if, for every vertex $v \in V(D)$, we have $\min(m^+(v), m^-(v)) \leq \Delta(\Delta - 1) - B$.

Theorem 1.7. *There exists an integer $\Delta_0 \geq 1$ such that the following holds. For every digraph D with maximum degree $\Delta_{\max}(D) = \Delta > \Delta_0$, if D is B -sparse with $B > \Delta \log^3 \Delta$, then*

$$\vec{\chi}(D) \leq (\Delta + 1) - \frac{B}{4e^7 \Delta}.$$

In Section 4, we give a sufficient condition for a digraph to admit an acyclic set of vertices intersecting each of its bicliques, by proving the following generalisation of a result of King [22].

Theorem 1.8. *Let D be a digraph with $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D) + 1)$. Then D has an acyclic set of vertices I such that $\vec{\omega}(D - I) = \vec{\omega}(D) - 1$.*

This implies that, in a minimum counterexample of Theorem 1.6, there must be a significant gap between the biclique number and the maximum degree. We indeed prove a more general result, as we characterise exactly the digraphs D with $\vec{\omega}(D) \geq \frac{2}{3}(\Delta_{\max}(D) + 1)$ that do not admit such an acyclic set of vertices. We thus extend a result due to Christofides *et al.* [10]. We also discuss the consequences of our result on list-dicolouring of digraphs, and show that we generalise a result due to Haxell [19] that we find of independent interest.

In Section 5, we deal with vertices having a dense neighbourhood in a counterexample of Theorem 1.6. We extend a result of King and Reed [23] by proving the following.

Theorem 1.9. *Let a and ε be constants satisfying $0 < \varepsilon \leq \frac{1}{6} - 4\sqrt{a}$. There exists $\Delta(a) \geq 1$ such that the following holds. Let D be a digraph with maximum degree $\Delta_{\max}(D) = \Delta \geq \Delta(a)$ satisfying $\vec{\omega}(D) \leq \frac{2}{3}(\Delta + 1)$ and let $v \in V(D)$ be such that $\max(m^-(v), m^+(v)) > (1 - a)\Delta(\Delta - 1)$. Then*

$$\vec{\chi}(D) \leq \max(\vec{\chi}(D - v), (1 - \varepsilon)(\Delta + 1)).$$

We put the pieces together and prove Theorem 1.6 in Section 6. In particular, we use a trick introduced by Harutyunyan and Mohar [18] to obtain the result with $\tilde{\Delta}$ instead of Δ_{\max} . We finally propose some further research directions in Section 7. In particular, we discuss possible analogues of Conjecture 1.5 with $\tilde{\Delta}(D)$ being replaced by $\Delta_{\min}(D) = \max_{v \in V(D)}(d_{\min}(v))$.

2 Preliminaries

2.1 General notation on digraphs

Our notation on digraphs follows [7]. A *digon* is a pair of arcs in opposite directions between the same vertices. A *simple arc* is an arc that is not in a digon. A digraph is *symmetric* if it contains

no simple arc. An *oriented graph* is a digraph with no digon. Given an undirected graph G , the digraph \overleftrightarrow{G} is the symmetric digraph obtained from G by replacing every edge with a digon. For an integer n , \overleftrightarrow{K}_n is called the *complete digraph* on n vertices.

Let D be a digraph. The *underlying graph* of D , denoted by $\text{UG}(D)$, is the undirected graph with vertex-set $V(D)$ in which uv is an edge if and only if uv or vu is an arc of D . The *symmetric part* of D , denoted by $S(D)$, is the undirected graph with vertex-set $V(D)$ in which uv is an edge if and only if $\{uv, vu\}$ is a digon of D . A *matching* of D is a set of pairwise disjoint arcs of D .

Let v be a vertex of D . The *out-degree* (resp. *in-degree*) of v , denoted by $d^+(v)$ (resp. $d^-(v)$), is the number of arcs leaving (resp. entering) v . We define the *maximum degree* of v as $d_{\max}(v) = \max\{d^+(v), d^-(v)\}$ and its *minimum degree* as $d_{\min}(v) = \min\{d^+(v), d^-(v)\}$. We can then define the corresponding maximum degrees of D :

$$\Delta_{\max}(D) = \max_{v \in V(D)} (d_{\max}(v)) \quad \text{and} \quad \Delta_{\min}(D) = \max_{v \in V(D)} (d_{\min}(v)).$$

The *maximum geometric mean* of D is defined as $\tilde{\Delta}(D) = \max_{v \in V(D)} \sqrt{d^-(v) \cdot d^+(v)}$. By definition, observe that we have $\Delta_{\min}(D) \leq \tilde{\Delta}(D) \leq \Delta_{\max}(D)$. We say that D is Δ -*diregular* if, for every vertex $v \in V(D)$, $d^-(v) = d^+(v) = \Delta$. Note that D being diregular implies $\Delta_{\min}(D) = \Delta_{\max}(D)$ but the converse is not true.

Let X, Y be two disjoint set of vertices of D . We denote by $A(X, Y)$ the set of arcs $uv \in A(D)$ such that $u \in X$ and $v \in Y$.

A *list assignment* L of D is a function that associates a list of colours to every vertex v of D . It is a *k -list assignment* if $|L(v)| \geq k$ holds for every vertex v of D . Let L be a list assignment. An *L -dicolouring* of D is a dicolouring α of D such that, for every vertex v of D , $\alpha(v) \in L(v)$. We say that D is *L -dicolourable* if D admits an L -dicolouring. Given an integer k , we say that D is *k -dichoosable* if, for every k -list assignment L of D , D is L -dicolourable.

2.2 Probabilistic tools

Our proof of Theorem 1.6 contains some probabilistic arguments. We briefly recall here two classical results used later. The reader unfamiliar with the probabilistic method is referred to [5] and [26]. We first need the symmetric version of the Lovász Local Lemma, due to Erdős and Lovász [13].

Lemma 2.1 (Lovász Local Lemma). *Let A_1, A_2, \dots, A_n be events in an arbitrary probability space. Suppose that each event A_i is mutually independent of a set of all the other events but at most d , and that $\Pr(A_i) \leq p$ for all $1 \leq i \leq n$. If*

$$ep(d+1) \leq 1,$$

then $\Pr(\bigwedge_{i=1}^n \overline{A_i}) > 0$.

The second result we need is the celebrated concentration bound due to Talagrand [32]. We use it in the following form (see [21, Theorem 6.3]).

Lemma 2.2 (Talagrand's Inequality). *Let X be a non-negative random variable, not identically 0, which is determined by n independent trials and satisfying the following for some $c, r > 0$:*

1. *Changing the outcome of any one trial can affect X by at most c , and*

2. for any s , if $X \geq s$ then there is a set of at most rs trials whose outcomes certify that $X \geq s$.

Then

$$\Pr(|X - \mathbf{E}(X)| > t) \leq 4 \exp\left(-\frac{t^2}{8rc^2(4\mathbf{E}(X) + t)}\right)$$

for any real number t with $t > 96c\sqrt{r\mathbf{E}(X)} + 128rc^2$.

3 Colouring sparse digraphs

Our first goal is to generalise the following result due to Molloy and Reed [26, Theorem 10.5] on the chromatic number of sparse graphs. For a vertex v of a graph G , we denote by $m(v)$ the number of edges of $G[N(v)]$. We say that G is B -sparse if $m(v) \leq \binom{\Delta(G)}{2} - B$ for every $v \in V(G)$.

Theorem 3.1 (Molloy and Reed [26]). *Let Δ be an integer large enough and G be a graph with maximum degree Δ . If G is B -sparse with $B > \Delta \log^3 \Delta$, then*

$$\chi(G) \leq (\Delta + 1) - \frac{B}{e^6 \Delta}.$$

The goal of this section is to prove Theorem 1.7, which is the directed analogue of Theorem 3.1. We first recall it here for convenience.

Theorem 1.7. *There exists an integer $\Delta_0 \geq 1$ such that the following holds. For every digraph D with maximum degree $\Delta_{\max}(D) = \Delta > \Delta_0$, if D is B -sparse with $B > \Delta \log^3 \Delta$, then*

$$\vec{\chi}(D) \leq (\Delta + 1) - \frac{B}{4e^7 \Delta}.$$

We will actually use Theorem 1.7 in the following form in Section 6.

Corollary 3.2. *There exists $\Delta_1 \geq 2$ such that the following holds. Let D be a digraph with maximum degree $\Delta_{\max}(D) = \Delta \geq \Delta_1$ and $a \in \mathbb{R}$ be such that $a > \frac{\log^3 \Delta}{\Delta - 1}$, and for every vertex $v \in V(D)$, $\min(m^+(v), m^-(v)) \leq (1 - a)\Delta(\Delta - 1)$. Then*

$$\vec{\chi}(D) \leq \left(1 - \frac{a}{5e^7}\right) (\Delta + 1).$$

Proof. By assumption D is B_a -sparse for $B_a = a\Delta(\Delta - 1) > \Delta \log^3 \Delta$. Hence, by taking $\Delta_1 \geq \Delta_0$, we obtain by Theorem 1.7 that

$$\vec{\chi}(D) \leq \left(1 - \frac{a}{4e^7}\right) (\Delta + 1) + \frac{a}{2e^7} \leq \left(1 - \frac{a}{5e^7}\right) (\Delta + 1),$$

where in the second inequality we used $\Delta_1 \geq 9$. □

The remainder of this section is devoted to the proof of Theorem 1.7. We first briefly justify that we only have to prove it for Δ -diregular digraphs.

Lemma 3.3. *Let D be a B -sparse digraph with maximum degree $\Delta_{\max}(D) = \Delta$. There exists a B -sparse Δ -diregular digraph H such that $\vec{\chi}(D) \leq \vec{\chi}(H)$.*

Proof. We repeat the following process until D is Δ -diregular. Add to D a disjoint copy of \bar{D} , the digraph obtained from D by reversing every arc. Then for every vertex $v \in V(D)$ such that $d_D^+(v) < \Delta$, we add the arc vv' , where v' is the copy of v in \bar{D} . Similarly, for every vertex v such that $d_D^-(v) < \Delta$, we add the arc $v'v$. Let D' be the obtained digraph. Note that $\bar{\chi}(D) \leq \bar{\chi}(D')$ trivially holds as D is a subdigraph of D' . Observe also that, for every vertex $v \in V(D)$, we have $m_D^+(v) = m_{D'}^+(v) = m_{D'}^-(v')$ and $m_D^-(v) = m_{D'}^-(v) = m_{D'}^+(v')$. Hence, as D is B -sparse, so is D' .

Observe finally that, for every vertex $v \in V(D)$, $d_{D'}^+(v) = d_{D'}^-(v') = \min(d_D^+(v) + 1, \Delta)$ and $d_{D'}^-(v) = d_{D'}^+(v') = \min(d_D^-(v) + 1, \Delta)$. Hence, repeating this process at most Δ times, we eventually find the desired digraph H . \square

A *partial* (k, ℓ) -dicolouring α of a digraph D is a k -dicolouring of an induced subdigraph of D such that, for every vertex v , either ℓ colours appear twice in $N^-(v)$ or ℓ colours appear twice in $N^+(v)$. We formally have

$$\begin{aligned} & |\{c \in [k] \mid \exists u, w \in N^-(v), u \neq w, \alpha(u) = \alpha(w) = c\}| \geq \ell \\ \text{or} \quad & |\{c \in [k] \mid \exists u, w \in N^+(v), u \neq w, \alpha(u) = \alpha(w) = c\}| \geq \ell. \end{aligned}$$

Lemma 3.4. *Let D be a Δ -diregular digraph and k, ℓ be two integers such that $k \leq \Delta + 1 - \ell$. If D admits a partial (k, ℓ) -dicolouring, then $\bar{\chi}(D) \leq \Delta + 1 - \ell$.*

Proof. Assume that D admits a partial (k, ℓ) -dicolouring. We let α be a partial $(\Delta + 1 - \ell, \ell)$ -dicolouring of D for which the number of uncoloured vertices is minimum, the existence of which is guaranteed as $k \leq \Delta + 1 - \ell$. We claim that α is a dicolouring of D , hence implying $\bar{\chi}(D) \leq \Delta + 1 - \ell$. Assume this is not the case, and there is a vertex v which is not coloured by α . By definition of a partial $(\Delta + 1 - \ell, \ell)$ -dicolouring, we have $|\alpha(N^-(v))| \leq \Delta - \ell$ or $|\alpha(N^+(v))| \leq \Delta - \ell$. Hence we can choose for v a colour of $[\Delta + 1 - \ell]$ that is not appearing either in its in- or in its out-neighbourhood. Colouring v with this specific colour extends α into a partial $(\Delta + 1 - \ell, \ell)$ -dicolouring of D with fewer uncoloured vertices, a contradiction to the choice of α . \square

We are now ready to prove Theorem 1.7.

Proof of Theorem 1.7. We do not give the explicit value of Δ_0 , we simply assume in what follows that it is arbitrarily large. We assume that D is Δ -diregular, for otherwise we may prove the result for the B -sparse Δ -diregular digraph H obtained by Lemma 3.3, and conclude that $\bar{\chi}(D) \leq \bar{\chi}(H) \leq (\Delta + 1) - \frac{B}{4e^7 \Delta}$.

We use the probabilistic method. Let us fix $k = \lfloor \frac{\Delta}{2} \rfloor$. We randomly colour every vertex of D with a colour of $[k]$. Then we simultaneously uncolour every vertex v having at least one in-neighbour and at least one out-neighbour sharing its colour. Our goal is to show that with non-zero probability, the obtained colouring α is a partial (k, ℓ) -dicolouring of D , where $\ell = \frac{B}{4e^7 \Delta}$. Observe that we have $k \leq \Delta + 1 - \ell$ as $B \leq \Delta(\Delta - 1)$. Hence, if α is indeed a partial (k, ℓ) -dicolouring of D , the result follows from Lemma 3.4. Note that α is necessarily a dicolouring of an induced subdigraph of D , for otherwise D , partially coloured with α , contains a monochromatic directed cycle, the vertices of which should have been uncoloured in the process. It remains to show that, for every vertex of D , ℓ colours are repeated either in its in-neighbourhood or in its out-neighbourhood.

For each vertex v , we let N_v be $N^+(v)$ if $m^+(v) \leq m^-(v)$ and $N^-(v)$ otherwise. We also let B_v be the number of missing arcs in $D[N_v]$, that is $B_v = \Delta(\Delta - 1) - \min(m^-(v), m^+(v))$. Note that $B_v \geq B > \Delta \log^3 \Delta$. Let X_v be the random variable corresponding to the number of colours that

are assigned to at least two vertices in N_v and are retained by all these vertices. We also define Y_v as the number of colours assigned to at least two vertices in N_v that are not linked by a digon, but not necessarily retained. Finally, we define Z_v as the number of colours assigned to at least two vertices in N_v that are not linked by a digon and one of these vertices does not retain its colour. We thus have $X_v = Y_v - Z_v$.

We let A_v be the event that $X_v < \ell$. Our goal is to show that none of the events A_v occurs with non-zero probability. To do so, we will prove that, for each vertex v , the probability that A_v occurs is small enough and then conclude by the Lovász Local Lemma (Lemma 2.1).

Claim 3.5. *For every vertex v , we have $\mathbf{E}(X_v) \geq \frac{B_v}{2e^{\tau}\Delta}$*

Proof of claim. For every pair of vertices $u, w \in N_v$ and every colour $c \in [k]$, we let $A_{u,w}^c$ be the event that u, w are exactly the vertices coloured c in N_v and they both retain their colour. We let $X_{u,w}^c$ be the binary random variable equals to 1 if $A_{u,w}^c$ occurs and 0 otherwise. We clearly have $X_v \geq \sum_{u,w,c} X_{u,w}^c$.

Assume u and w are not linked by a digon. If uw is a simple arc of D , then $A_{u,w}^c$ occurs if and only if both u and w are coloured c and no vertex in $N = (N_v \cup N^-(u) \cup N^+(w)) \setminus \{u, w\}$ receives colour c . If the arc uw is missing, then $A_{u,w}^c$ occurs in particular if both u and w are coloured c and no vertex in $N = (N_v \cup N^+(u) \cup N^-(w)) \setminus \{u, w\}$ receives colour c . In both cases, as $k = \lfloor \frac{\Delta}{2} \rfloor$, we have $|N| \leq 3\Delta \leq 7k$, which implies

$$\Pr(A_{u,w}^c) \geq \left(\frac{1}{k}\right)^2 \left(1 - \frac{1}{k}\right)^{7k}.$$

By definition of B_v we have at least $\frac{B_v}{2}$ distinct pair of vertices that are not linked by a digon in $N^+(v)$. We also have k choices for the colour c . By linearity of the expectation we thus obtain

$$\mathbf{E}(X_v) \geq k \frac{B_v}{2} \left(\frac{1}{k}\right)^2 \left(1 - \frac{1}{k}\right)^{7k} \geq \frac{B_v}{\Delta} \left(1 - \frac{1}{k}\right)^{7k} \geq \frac{B_v}{2e^{\tau}\Delta},$$

where in the last inequality we used that $k = \lfloor \frac{\Delta}{2} \rfloor$ is large enough and $\lim_{k \rightarrow +\infty} \left(1 - \frac{1}{k}\right)^{7k} = \frac{1}{e^{\tau}} > \frac{1}{2e^{\tau}}$. \diamond

Claim 3.6. *For every vertex v , we have $\mathbf{E}(X_v) \leq \mathbf{E}(Y_v) \leq \frac{3B_v}{\Delta}$*

Proof of claim. Since $X_v = Y_v - Z_v$ it is clear that $\mathbf{E}(X_v) \leq \mathbf{E}(Y_v)$. Given two vertices $u, w \in N_v$ that are not linked by a digon and a colour $c \in [k]$, we let $Y_{u,w}^c$ be the binary random variable equals to 1 if both u and w are coloured c and 0 otherwise. We have $Y_v \leq \sum_{u,w,c} Y_{u,w}^c$ and $\Pr(Y_{u,w}^c = 1) = \frac{1}{k^2}$. Since we have at most B_v choices for the pair u, w and exactly k for the colour c , by linearity of the expectation we obtain

$$\mathbf{E}(X_v) \leq \mathbf{E}(Y_v) \leq kB_v \left(\frac{1}{k}\right)^2 \leq \frac{3B_v}{\Delta},$$

where in the last inequality we used $k = \lfloor \frac{\Delta}{2} \rfloor \geq \frac{\Delta}{3}$. \diamond

Claim 3.7. *For every vertex v , $\Pr\left(|X_v - \mathbf{E}(X_v)| > \log \Delta \sqrt{\mathbf{E}(X_v)}\right) < \frac{1}{\Delta^5}$.*

Proof of claim. Let us set $t = \frac{1}{2} \log \Delta \sqrt{\mathbf{E}(X_v)}$. Recall that $X_v = Y_v - Z_v$, so by linearity of the expectation we have $\mathbf{E}(X_v) = \mathbf{E}(Y_v) - \mathbf{E}(Z_v)$, and $|X_v - \mathbf{E}(X_v)| \leq |Y_v - \mathbf{E}(Y_v)| + |Z_v - \mathbf{E}(Z_v)|$. In particular, this implies

$$\Pr(|X_v - \mathbf{E}(X_v)| > 2t) \leq \Pr(|Y_v - \mathbf{E}(Y_v)| > t) + \Pr(|Z_v - \mathbf{E}(Z_v)| > t). \quad (1)$$

We will use Talagrand's inequality (Lemma 2.2) to bound both $\Pr(|Y_v - \mathbf{E}(Y_v)| > t)$ and $\Pr(|Z_v - \mathbf{E}(Z_v)| > t)$. Observe that changing the colour of exactly one vertex affects Y_v by at most 1. Also if $Y_v \geq s$ then there exists a set of $2s$ vertices whose colours certify $Y_v \geq s$: for each of the s colours we may choose two vertices not linked by a digon sharing this colour. Combining Claims 3.5 and 3.6 we obtain $\mathbf{E}(Y_v) \leq 6e^7 \mathbf{E}(X_v)$. Hence, Δ being sufficiently large, we have $t > 96\sqrt{2\mathbf{E}(Y_v)} + 256$, which allows us to apply Talagrand's inequality to Y_v and obtain

$$\Pr(|Y_v - \mathbf{E}(Y_v)| > t) \leq 4 \exp\left(-\frac{t^2}{64\mathbf{E}(Y_v) + 16t}\right). \quad (2)$$

We now consider Z_v . Changing the colour of exactly one vertex affects Z_v by at most 2. Also if $Z_v \geq s$ then there exists a set of $4s$ vertices whose colours certify $Z_v \geq s$: for each of the s colours we may choose four vertices a, b, b^-, b^+ sharing this colour, $a \neq b$, such that a and b are not linked by a digon in D , and $b^-b, bb^+ \in A(D)$ (a, b^-, b^+ are not necessarily distinct). Observe that $Z_v \leq Y_v$, so in particular $\mathbf{E}(Z_v) \leq \mathbf{E}(Y_v) \leq 6e^7 \mathbf{E}(X_v)$ by Claims 3.5 and 3.6. Hence, Δ being sufficiently large, we have $t > 192\sqrt{4\mathbf{E}(Z_v)} + 2048$, which allows us to apply Talagrand's inequality to Z_v and obtain

$$\Pr(|Z_v - \mathbf{E}(Z_v)| > t) \leq 4 \exp\left(-\frac{t^2}{512\mathbf{E}(Z_v) + 128t}\right). \quad (3)$$

Combining (1), (2), and (3) together with $\mathbf{E}(Z_v) \leq \mathbf{E}(Y_v) \leq 6e^7 \mathbf{E}(X_v)$ we obtain

$$\begin{aligned} \Pr(|X_v - \mathbf{E}(X_v)| > 2t) &\leq 8 \exp\left(-\frac{t^2}{3072e^7 \mathbf{E}(X_v) + 128t}\right) = 8 \exp\left(-\frac{\log^2 \Delta}{12288e^7 + 256 \frac{\log \Delta}{\sqrt{\mathbf{E}(X_v)}}}\right) \\ &\leq 8 \exp\left(-\frac{\log^2 \Delta}{12288e^7 + 256\sqrt{2e^7} \frac{1}{\sqrt{\log \Delta}}}\right), \end{aligned}$$

where in the last inequality we used $\mathbf{E}(X) \geq \frac{B_v}{2e^7 \Delta}$ and $B_v \geq \Delta \log^3 \Delta$. As Δ is sufficiently large, we obtain

$$\Pr\left(|X_v - \mathbf{E}(X_v)| > \log \Delta \sqrt{\mathbf{E}(X_v)}\right) < \frac{1}{\Delta^5},$$

which concludes the proof of the claim. \diamond

By Claim 3.5 together with the fact that $B_v \geq B \geq \Delta \log^3 \Delta$, we have

$$\mathbf{E}(X_v) - \log \Delta \sqrt{\mathbf{E}(X_v)} \geq \frac{B_v}{2e^7 \Delta} \left(1 - \log \Delta \sqrt{\frac{2e^7 \Delta}{B_v}}\right) \geq \frac{B}{2e^7 \Delta} \left(1 - \sqrt{\frac{2e^7}{\log \Delta}}\right).$$

By choice of ℓ , we thus have $\mathbf{E}(X_v) - \log \Delta \sqrt{\mathbf{E}(X_v)} \geq 2\ell \left(1 - \sqrt{\frac{2e^7}{\log \Delta}}\right)$, which is larger than ℓ for Δ large enough. Hence we obtain

$$\Pr(A_v) = \Pr(X_v < \ell) \leq \Pr\left(X_v < \mathbf{E}(X) - \log \Delta \sqrt{\mathbf{E}(X)}\right) \leq \Pr\left(|X_v - \mathbf{E}(X)| > \log \Delta \sqrt{\mathbf{E}(X)}\right).$$

Therefore, by Claim 3.7, we have $\Pr(A_v) < \frac{1}{\Delta^5}$. Observe finally that the value of X_v depends only on the colour of vertices at distance at most 2 of v in $\text{UG}(D)$. Hence, each event A_v is mutually independent of all other events A_u except at most $(2\Delta)^4 = d$ of them. For every vertex v , we have $e\Pr(A_v)(d+1) \leq e\frac{16\Delta^4+1}{\Delta^5}$, which is smaller than 1 if Δ is large enough. Hence by the Lovász Local Lemma (Lemma 2.1) none of the events A_v occurs with non-zero probability, which concludes the proof. \square

4 Hitting every maximum biclique with an acyclic set of vertices

Let G be a graph with maximum degree $\Delta(G) \leq 3k - 1$. Haxell [19] proved that, if the vertices of G can be partitioned into r cliques C_1, \dots, C_r of size $\geq 2k$, then G contains an independent set S of size r . In particular, S intersects each clique C_i .

Using this result, Rabern [29] proved that every graph G satisfying $\omega(G) \geq \frac{3}{4}(\Delta(G) + 1)$ contains an independent set intersecting every maximum clique of G . This was later improved to $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$ by King [22], which is best possible. Christofides, Edwards, and King [10] later characterised exactly the graphs G satisfying $\omega(G) = \frac{2}{3}(\Delta(G) + 1)$ that do not admit an independent set intersecting every maximum cliques. Combined with King's earlier result, they thus obtained the following. Given two graphs G and G' , we denote by $G \circ G'$ the lexicographic product of G by G' , which is the graph with vertex-set $V(G) \times V(G')$ and edge-set $\{(u, u')(v, v') \mid uv \in E(G) \text{ or } u = v \text{ and } u'v' \in E(G')\}$.

Theorem 4.1 (Christofides *et al.* [10]). *Let G be a connected graph with $\omega(G) \geq \frac{2}{3}(\Delta(G) + 1)$. Then G has an independent set I such that $\omega(G - I) = \omega(G) - 1$, unless $G = C_n \circ K_p$ with $n \geq 5$ odd and $p = \frac{1}{2}\omega(G)$.*

The goal of this section is to prove the following generalisation of Theorem 4.1. Note that, in the proof of Theorem 1.6, we will only use Theorem 4.2 with $\overleftrightarrow{\omega}(D)$ being larger than $\frac{2}{3}(\Delta + 1)$. We find the characterisation of the equality case of independent interest and we thus also include the case $\overleftrightarrow{\omega}(D)$ being equal to $\frac{2}{3}(\Delta + 1)$.

Theorem 4.2. *Let D be a connected digraph and Δ be an integer with $\Delta_{\max}(D) \leq \Delta$ and $\overleftrightarrow{\omega}(D) \geq \frac{2}{3}(\Delta + 1)$. Then D has an acyclic set of vertices I such that $\overleftrightarrow{\omega}(D - I) = \overleftrightarrow{\omega}(D) - 1$, unless $D = C_n \circ K_p$ with $n \geq 5$ odd and $p = \frac{1}{2}\overleftrightarrow{\omega}(D) = \frac{1}{3}(\Delta + 1)$.*

We first prove the following result, which generalises to digraphs a more precise version of Haxell's result obtained by King [22, Theorem 5].

Theorem 4.3. *Let k be a positive integer and D be a digraph with vertices partitioned into bicliques C_1, \dots, C_r . If for every $i \in [r]$ and every $v \in C_i$, v has at most k out-neighbours outside C_i and at most $|C_i| - k$ in-neighbours outside C_i , then D contains an acyclic set of size r .*

Given a digraph D whose vertices are partitioned into independent sets V_1, \dots, V_r , an *acyclic system of representative* (with respect to V_1, \dots, V_r), or *ASR* for conciseness, is an acyclic set of vertices X such that, for every $i \in [r]$, $|V_i \cap X| = 1$. If we relax this condition to $|V_i \cap X| \leq 1$, then X is a *partial ASR*. Theorem 4.3 can be restated in terms of ASR as follows.

Theorem 4.4. *Let k be a positive integer and D be a digraph with vertices partitioned into independent sets V_1, \dots, V_r . If every vertex $v \in V_i$ satisfies $d^+(v) \leq k$ and $d^-(v) \leq |V_i| - k$, then D admits an ASR.*

Following the analogue of Haxell's idea [19], Theorem 4.4 has the following consequence about list-dicolouring that we find of independent interest.

Corollary 4.5. *Let D be a digraph, L be a list assignment of D , and k be an integer. If for every vertex $v \in V(D)$ and for every colour $c \in L(v)$, we have*

$$|N^+(v) \cap \{u \mid c \in L(u)\}| \leq k \quad \text{and} \quad |N^-(v) \cap \{u \mid c \in L(u)\}| \leq |L(v)| - k,$$

then D is L -dicolourable.

Proof. Let H be the auxiliary digraph where $V(H) = \{(v, c) \mid v \in V(D), c \in L(v)\}$ which contains an arc from (u, c_1) to (v, c_2) whenever $c_1 = c_2$ and $v \in N^+(u)$. Let us label with v_1, \dots, v_r the vertices of D , and for every $i \in [r]$ let $V_i = \{(v_i, c) \mid c \in L(v_i)\}$. By definition, V_1, \dots, V_r is a partition of $V(H)$ into independent sets. The conditions on L imply that every vertex $v \in V_i$ has out-degree at most k and in-degree at most $|V_i| - k$. Hence, by Theorem 4.4, H contains an ASR $X = \{(v_1, c_1), \dots, (v_r, c_r)\}$. Let α be the colouring of D which associates to each vertex v_i colour c_i . We claim that α is an L -dicolouring of D . We have $c_i \in L(v_i)$ by definition. Assume for a contradiction that D , coloured with α , contains a monochromatic directed cycle v_1, \dots, v_ℓ . Let c be the colour of this cycle, then $(v_1, c), \dots, (v_\ell, c)$ is a directed cycle in $H[X]$, a contradiction. \square

Proof of Theorem 4.4. For any subset of indices $I \subseteq [r]$ we denote by V_I the set of vertices $\bigcup_{i \in I} V_i$. Given a set of vertices $X \subseteq V(D)$, we denote by $\iota(X)$ the set of indices $\{i \in [r] \mid V_i \cap X \neq \emptyset\}$. For a single vertex v , we denote by $\iota(v)$ the integer i such that $v \in V_i$.

We proceed by induction on $r \geq 1$. The result being trivial when $r = 1$, we assume $r \geq 2$. Let us fix any vertex $x_1 \in V_r$, we will show as a stronger result that there exists an ASR containing x_1 . Assume for a contradiction that no such ASR exists.

Let $I \subseteq [r - 1]$ and $X, Y \subseteq (V_I \cup \{x_1\})$. We say that (I, X, Y) is a *good triplet* if each of the following properties holds:

1. X and Y are two disjoint acyclic sets with $x_1 \in X$;
2. Y is an ASR of $D[V_I]$ (with respect to $(V_i)_{i \in I}$);
3. every vertex in Y has exactly one in-neighbour in X and every vertex in X has at least one out-neighbour in Y , in particular $|X| \leq |Y|$; and
4. every vertex $v \in V_I \cup \{x_1\}$ has an in-neighbour in X or an out-neighbour in Y .

We first justify that there exists no good triplet in D .

Claim 4.6. *D admits no good triplet (I, X, Y) .*

Proof of claim. Assume for a contradiction that (I, X, Y) is a good triplet of D . The assumption on the degrees in D implies in particular $d^+(x) \leq k$ for every vertex $x \in X$ and $d^-(y) \leq |V_{\iota(y)}| - k$ for every vertex $y \in Y$. Observe that $|X| \leq |Y| = |I|$ by 3 and 2. We thus have

$$\sum_{x \in X} d^+(x) + \sum_{y \in Y} d^-(y) \leq k|I| + \sum_{i \in I} (|V_i| - k) = |V_I|.$$

On the other hand, we know by 4 that $(V_I \cup \{x_1\}) \subseteq (N^+(X) \cup N^-(Y))$. This implies

$$\sum_{x \in X} d^+(x) + \sum_{y \in Y} d^-(y) \geq |N^+(X) \cup N^-(Y)| \geq |V_I| + 1,$$

yielding the contradiction. \diamond

Our goal is now to show that D admits an infinite sequence $(Y_i)_{i \geq 1}$ of partial ASR, such that each Y_i is a proper subset of Y_{i+1} . This will yield the contradiction since such a sequence has length at most r .

Let R_1 be an ASR of $D[V_{[r-1]}]$ (with respect to V_1, \dots, V_{r-1}) for which the size of $Y_1 = Y'_1 = N^+(x_1) \cap R_1$ is minimum. The existence of R_1 is guaranteed by induction on r . Observe that $Y_1 \neq \emptyset$ for otherwise $R_1 \cup \{x_1\}$ is an ASR of D containing x_1 . We let X_1 be the singleton $\{x_1\}$. We will recursively build $(R_{i+1}, X_{i+1}, Y_{i+1}, Y'_{i+1})$ from (R_i, X_i, Y_i, Y'_i) in such a way that, for every $i \geq 1$, each of the following properties holds.

1. $X_i = \{x_1, \dots, x_i\}$ is acyclic. If $i > 1$ then $x_i \in V_{\iota(Y_{i-1})}$.
2. R_i is an ASR of $D[V_{[r-1]}]$ such that for every $1 \leq j \leq i-1$, $R_i \cap N^+(x_j) = Y'_j$. Subject to this property, $Y'_i = N^+(x_i) \cap R_i$ has minimum size.
3. $Y_i = \bigsqcup_{1 \leq j \leq i} Y'_j$ and $Y'_i \neq \emptyset$.

Observe that (R_1, X_1, Y_1, Y'_1) satisfies 1, 2, and 3. Also, the existence of such an infinite sequence $(R_i, X_i, Y_i, Y'_i)_{i \geq 1}$ implies the desired infinite sequence $(Y_i)_{i \geq 1}$ of partial ASR, yielding the contradiction.

Hence we assume that for some $i \geq 1$, there exists $(R_i, X_i, Y_i, Y'_i)_{1 \leq j \leq i}$ such that, for each $j \in [i]$, 1, 2, and 3 hold. We construct $(R_{i+1}, X_{i+1}, Y_{i+1}, Y'_{i+1})$ as follows.

- We let x_{i+1} be a vertex in $V_{\iota(Y_i)} \setminus (X_i \cup Y_i)$ such that

$$N^-(x_{i+1}) \cap X_i = \emptyset \quad \text{and} \quad N^+(x_{i+1}) \cap Y_i = \emptyset. \quad (\star)$$

If such a vertex x_{i+1} exists then $X_{i+1} = \{x_1, \dots, x_{i+1}\}$ is acyclic as x_{i+1} is a source in $D[X_{i+1}]$ and $X_i = \{x_1, \dots, x_i\}$ is acyclic by recurrence, so 1 holds.

Let us justify the existence of x_{i+1} . If no such vertex exists, then X_i and Y_i are two disjoint acyclic sets such that $x_1 \in X_i$, with Y_i being an ASR of $V_{\iota(Y_i)}$. Combining conditions 2 and 3 we obtain that $(N^+(x_1), \dots, N^+(x_i))$ partitions Y_i into non-empty parts, so in particular every vertex in Y_i has exactly one in-neighbour in X_i and every vertex in X_i has at least one out-neighbour in Y_i . Hence, if no vertex in $V_{\iota(Y_i)} \setminus (X_i \cup Y_i)$ satisfies (\star) , we deduce that $(\iota(Y_i), X_i, Y_i)$ is a good triplet, a contradiction to Claim 4.6.

- We let R_{i+1} be an ASR of $D[V_{[r-1]}]$ such that for every $1 \leq j \leq i$, $R_{i+1} \cap N^+(x_j) = Y'_j$, and subject to this property, $Y'_{i+1} = N^+(x_{i+1}) \cap R_{i+1}$ has minimum size. The existence of R_{i+1} is guaranteed as R_i is such an ASR. Hence 2 is maintained.
- It remains to show that 3 is maintained, that is $Y'_{i+1} \neq \emptyset$ and $Y'_{i+1} \cap Y_i = \emptyset$. First we have $Y'_{i+1} \cap Y_i = \emptyset$ by (\star) . Now assume for a contradiction that $Y'_{i+1} = \emptyset$, which together with (\star) implies that x_{i+1} has no out-neighbour in R_{i+1} .

By recurrence we know that (Y'_1, \dots, Y'_i) partitions Y_i . As x_{i+1} belongs to $V_{\iota(Y_i)} \setminus (X_i \cup Y_i)$ there exists a unique $j \leq i$ such that $\iota(x_{i+1}) \in \iota(Y'_j)$. As Y'_j is a partial ASR, there exists a unique $y_j \in Y'_j \cap V_{\iota(x_{i+1})}$. Consider the set \tilde{R}_j defined as

$$\tilde{R}_j = R_{i+1} \setminus \{y_j\} \cup \{x_{i+1}\}.$$

Observe that \tilde{R}_j is necessarily acyclic as x_{i+1} has no out-neighbour in R_{i+1} and R_{i+1} is an ASR of $D[V_{[r-1]}]$. Also, since $y_j \in V_{\iota(x_{i+1})}$, we obtain that \tilde{R}_j is an ASR of $D[V_{[r-1]}]$.

Since x_{i+1} has no in-neighbour in X_i by (\star) , for every $1 \leq \ell < j$, we have $\tilde{R}_j \cap N^+(x_\ell) = R_{i+1} \cap N^+(x_\ell) = Y'_\ell$. We finally have $\tilde{R}_j \cap N^+(x_j) = (R_{i+1} \cap N^+(x_j)) \setminus \{y_j\} = Y'_j \setminus \{y_j\}$, a contradiction to the choice of R_j .

Hence 3 is maintained and the proof follows. \square

With Theorem 4.3 in hand, we are now going to prove Theorem 4.2. We need two additional intermediate results. Let G be a (di)graph. We denote by $\mathcal{C}(G)$ the set of all maximum (bi)cliques of G . Given a collection $\mathcal{C} \subseteq \mathcal{C}(G)$ of maximum (bi)cliques of G , we denote by $\text{IG}(\mathcal{C})$ the intersection graph of \mathcal{C} , *i.e.* the undirected graph with vertex-set \mathcal{C} which contains an edge between C_1 and C_2 whenever $C_1 \cap C_2 \neq \emptyset$. We first need the following result due to Hajnal [16].

Lemma 4.7 (Hajnal [16]). *Let G be a graph and \mathcal{C} be a collection of maximum cliques of G , then*

$$|\bigcap \mathcal{C}| + |\bigcup \mathcal{C}| \geq 2\omega(G).$$

The second intermediate result we need is an extension of a result of Kostochka [24] due to Christofides, Edwards, and King [10]. The statement we use slightly differs from the original one (see [31, Lemma 6.12] for a proof of the result in the following form).

Lemma 4.8 (Christofides *et al.* [10]). *Let G be a graph with $\omega(G) \geq \frac{2}{3}(\Delta(G) + 1)$, let $\mathcal{C} \subseteq \mathcal{C}(G)$ such that $\text{IG}(\mathcal{C})$ is connected, let $X = \bigcap \mathcal{C}$ and let $Y = \bigcup \mathcal{C}$. If $X = \emptyset$ then $\omega(G) = \frac{2}{3}(\Delta(G) + 1)$ and Y has a partition $\{Q_i \mid i \in [n]\}$ into $n \geq 4$ cliques of G each of cardinality $\frac{1}{2}\omega(G)$ such that $\mathcal{C} \setminus \{Q_1 \cup Q_n\} = \{Q_i \cup Q_{i+1} \mid i \in [n-1]\}$.*

We are now ready to prove Theorem 4.2, let us first restate it for convenience.

Theorem 4.2. *Let D be a connected digraph and Δ be an integer with $\Delta_{\max}(D) \leq \Delta$ and $\overleftrightarrow{\omega}(D) \geq \frac{2}{3}(\Delta+1)$. Then D has an acyclic set of vertices I such that $\overleftrightarrow{\omega}(D-I) = \overleftrightarrow{\omega}(D) - 1$, unless $D = \overleftarrow{C}_n \circ \overrightarrow{K}_p$ with $n \geq 5$ odd and $p = \frac{1}{2}\overleftrightarrow{\omega}(D) = \frac{1}{3}(\Delta + 1)$.*

Proof. Let D be a minimum counterexample, which means that for some integer Δ , $\Delta_{\max}(D) \leq \Delta$, $\overleftrightarrow{\omega}(D) \geq \frac{2}{3}(\Delta+1)$, D is distinct from the obstruction $\overleftarrow{C}_n \circ \overrightarrow{K}_p$ and D does not contain any acyclic set of vertices I intersecting every maximum biclique of D . We first show that every vertex v belongs to a maximum biclique of D . Assume that this is not the case for some vertex v and let D' be $D - v$. Observe that $\overleftrightarrow{\omega}(D') = \overleftrightarrow{\omega}(D) \geq \frac{2}{3}(\Delta+1) \geq \Delta_{\max}(D')$. By minimality of D , either D' has an acyclic set I such that $\overleftrightarrow{\omega}(D' - I) = \overleftrightarrow{\omega}(D') - 1$ or D' has a connected component $H = \overleftarrow{C}_n \circ \overrightarrow{K}_p$ with $n \geq 5$ and $p = \frac{1}{2}\overleftrightarrow{\omega}(D)$. In the former case, as v does not belong to a maximum clique of D , I is also an acyclic set of D such that $\overleftrightarrow{\omega}(D - I) = \overleftrightarrow{\omega}(D) - 1$, a contradiction to D being a counterexample. In the latter case, every vertex $x \in V(H)$ satisfies $d_H^+(x) = d_H^-(x) = \Delta$. By connectivity of D ,

some vertex $x \in V(H)$ is adjacent to v , implying that $\max(d_D^+(x), d_D^-(x)) > \Delta$, a contradiction to $\Delta_{\max}(D) \leq \Delta$.

Let \mathcal{H} be $\text{IG}(\mathcal{C}(G))$. We denote by $\mathcal{H}_1, \dots, \mathcal{H}_r$ the connected components of \mathcal{H} .

Claim 4.9. *There exists a connected component \mathcal{H}_i of \mathcal{H} such that $\bigcap_{C \in V(\mathcal{H}_i)} C = \emptyset$.*

Proof of claim. Assume for a contradiction that $X_i = \bigcap_{C \in V(\mathcal{H}_i)} C \neq \emptyset$ holds for every $i \in [r]$. Let D' be the subdigraph of D induced by $\bigcup_{i \in [r]} X_i$.

We claim that we can apply Theorem 4.3 to D' , the vertices of D' being partitioned into bicliques X_1, \dots, X_r , with $k = \lfloor \frac{1}{3}(\Delta + 1) \rfloor$. Let i be any index of $[r]$ and v_i be any vertex in X_i . Let us justify that the out-degree of v_i outside X_i in D' is at most k , and that its in-degree outside X_i is at most $|X_i| - k$.

Observe that v_i has at least $\vec{\omega}(D) - 1$ out-neighbours in $Y_i = \bigcup_{C \in V(\mathcal{H}_i)} C$, so the out-degree of v_i outside X_i in D' is at most $\Delta - \vec{\omega}(D) + 1 \leq \frac{1}{3}(\Delta + 1)$. Since it is an integer, the out-degree of v_i outside X_i in D' is thus at most k . On the other hand, v_i has at most $\Delta - |Y_i| + 1$ in-neighbours outside X_i in D' . By Lemma 4.7 applied to $S(D)$ we have $|Y_i| \geq 2\vec{\omega}(D) - |X_i| \geq \frac{4}{3}(\Delta + 1) - |X_i|$. Hence v_i has at most $|X_i| - \frac{1}{3}(\Delta + 1) \leq |X_i| - k$ in-neighbours outside X_i in D' .

Therefore, we can apply Theorem 4.3 to D' , and conclude that D' admits an acyclic set I such that $\vec{\omega}(D') = \vec{\omega}(D - I)$. By construction of D' , I is also an acyclic set of D such that $\vec{\omega}(D) = \vec{\omega}(D - I)$, a contradiction. \diamond

Henceforth, by Claim 4.9, we assume without loss of generality that $\bigcap_{C \in V(\mathcal{H}_r)} C = \emptyset$. By Lemma 4.8 applied to $S(D)$ with $\mathcal{C} = V(\mathcal{H}_r)$, we obtain $\vec{\omega}(D) = \omega(S(D)) = \frac{2}{3}(\Delta + 1)$ and $Y = \bigcup_{C \in \mathcal{C}} C$ admits a partition Q_1, \dots, Q_n into $n \geq 4$ bicliques of D each of cardinality $\frac{1}{2}\vec{\omega}(D)$ such that $\mathcal{C} \setminus \{Q_1 \cup Q_n\} = \{Q_i \cup Q_{i+1} \mid i \in [n-1]\}$.

If $Q_1 \cup Q_n \in \mathcal{C}$, then $D[Y]$ is Δ -diregular, so necessarily $Y = V(D)$ as D is connected, and D is exactly $\overleftarrow{C}_n \circ \overrightarrow{K}_p$. If n is odd, this is a contradiction to D being a counterexample. If n is even, one can arbitrarily pick one representative of each Q_i with $i \geq 1$ odd to obtain an acyclic set intersecting every maximum biclique of D , a contradiction.

Hence $\mathcal{C} = \{Q_i \cup Q_{i+1} \mid i \in [n-1]\}$. Let D' be the digraph obtained from D by removing $\bigcup_{i=2}^{n-1} Q_i$ and adding every possible arcs between Q_1 and Q_n . It is straightforward that $\Delta_{\max}(D) \geq \Delta_{\max}(D')$ and that D' is smaller than D .

Note that $\vec{\omega}(D') \geq \vec{\omega}(D)$ as $Q_1 \cup Q_n$ is a biclique of D' of size $\vec{\omega}(D)$. We claim that equality holds, so assume for a contradiction that D' contains a biclique C' of size $\vec{\omega}(D) + 1$. Let $q_1 = |C' \cap Q_1|$ and $q_n = |C' \cap Q_n|$. Observe that $q_1 > 0$ and $q_n > 0$ as C' is not a biclique of D . Let us fix $v \in C' \setminus (Q_1 \cup Q_n)$, which exists since $|Q_1 \cup Q_n| < |C'|$. Recall that v belongs to a maximum biclique C of D , which is necessarily disjoint from $Q_1 \cup Q_n$ (otherwise C would belong to \mathcal{C}). If $q_1 + q_n > \frac{1}{3}(\Delta + 1)$, then $d_D^+(v) \geq q_1 + q_n + \vec{\omega}(D) - 1 > \Delta$, a contradiction. Hence $q_1 + q_n \leq \frac{1}{3}(\Delta + 1)$, which implies $|C' \setminus (Q_1 \cup Q_n)| = \vec{\omega}(D) + 1 - q_1 - q_n \geq \vec{\omega}(D) + 1 - \frac{1}{3}(\Delta + 1)$. Let v_1 be any vertex in $C' \cap Q_1$, then $d_D^+(v_1) \geq |C' \setminus (Q_1 \cup Q_n)| + |Q_1 \cup Q_2| - 1 \geq 2\vec{\omega}(D) - \frac{1}{3}(\Delta + 1) \geq \Delta + 1$, a contradiction.

This shows that $\vec{\omega}(D') = \vec{\omega}(D)$. Note also that D' is connected as D is, and vertices in $\bigcup_{i=2}^{n-1} Q_i$ do not have neighbours outside Y . Since D is a minimum counterexample, either $D' = \overleftarrow{C}_k \circ \overrightarrow{K}_p$ or D' has an acyclic set I such that $\vec{\omega}(D' - I) = \vec{\omega}(D') - 1$. In the former case, we obtain that $D = \overleftarrow{C}_{k+n-2} \circ \overrightarrow{K}_p$, a contradiction to D being a counter example: either $k + n - 2$ is odd or we can easily find an acyclic set intersecting every maximum biclique. Assume the latter case holds.

As I intersects every maximum biclique of D' , in particular it intersects $Q_1 \cup Q_n$. We assume by symmetry that it intersects Q_1 . For every $i \in [n]$, let v_i be an arbitrary vertex of Q_i . If $2 \leq i \leq n-1$, observe that $N^+(v_i) = N^-(v_i) = (Q_{i-1} \cup Q_i \cup Q_{i+1}) \setminus \{v_i\}$.

If n is even, $I \cup \{v_{2i+1} \mid i \in [\frac{n-2}{2}]\}$ is an acyclic set intersecting every maximum biclique of D . If n is odd, then $I \setminus Q_1 \cup \{v_{2i} \mid i \in [\frac{n-1}{2}]\}$ is an acyclic set intersecting every maximum biclique of D . Note that this holds as vertices in Q_1 belong to exactly one biclique, namely $Q_1 \cup Q_2$. Both cases yield a contradiction. \square

5 Handling vertices with a dense neighbourhood

In the short proof of Theorem 1.2 due to King and Reed [23], Theorem 3.1 handles the case of graphs in which every vertex has a sparse neighbourhood. The following result handles the remaining case, that is graphs admitting a vertex with a dense neighbourhood.

Theorem 5.1 (King and Reed [23]). *Let a and ε be constants satisfying $0 < \varepsilon < \frac{1}{6} - 2\sqrt{a}$. Let G be a graph with maximum degree Δ satisfying $\omega(G) \leq \frac{2}{3}(\Delta+1)$ and let v be such that $m(v) > (1-a)\binom{n}{2}$. Then*

$$\chi(G) \leq \max(\chi(G-v), (1-\varepsilon)(\Delta+1)).$$

The goal of this section is to prove Theorem 1.9, which is the direct analogue of Theorem 5.1 for digraphs. We first recall it here for convenience.

Theorem 1.9. *Let a and ε be constants satisfying $0 < \varepsilon \leq \frac{1}{6} - 4\sqrt{a}$. There exists $\Delta(a) \geq 1$ such that the following holds. Let D be a digraph with maximum degree $\Delta_{\max}(D) = \Delta \geq \Delta(a)$ satisfying $\vec{\omega}(D) \leq \frac{2}{3}(\Delta+1)$ and let $v \in V(D)$ be such that $\max(m^-(v), m^+(v)) > (1-a)\Delta(\Delta-1)$. Then*

$$\vec{\chi}(D) \leq \max(\vec{\chi}(D-v), (1-\varepsilon)(\Delta+1)).$$

We will use the following classical result due to Erdős, Rubin, and Taylor [14].

Lemma 5.2 (Erdős et al. [14]). *Let G be a graph obtained from K_n by removing a matching of size p . Then G is $(n-p)$ -choosable.*

This directly gives the following.

Lemma 5.3. *Let D be a digraph obtained from \overleftrightarrow{K}_n by removing a matching of size p . Then D is $(n-p)$ -dichoosable.*

Proof. Let L be any $(n-p)$ -list assignment of D . By Lemma 5.2, there exists an L -colouring α of $S(D)$, because $S(D)$ is obtained from K_n by removing a matching of size p . As D is obtained from $\overleftrightarrow{S}(D)$ by adding an acyclic set of arcs, α must be an L -dicolouring of D . \square

Proof of Theorem 1.9. We take $\Delta(a) = \max\left(\frac{1-a}{\sqrt{a-a}}, \frac{1-a}{a}\right)$. By directional duality we assume without loss of generality that $m^+(v) \geq (1-a)\Delta(\Delta-1)$ and let $N = N^+(v) \cup \{v\}$ and $\bar{N} = V(D) \setminus N$. We assume that $|N| = \Delta+1$ for otherwise we add new vertices dominated by v . We partition N

into three parts as follows:

$$\begin{aligned} N_1 &= \left\{ u \in N \mid |N^+(u) \cap \bar{N}| \geq \frac{\Delta}{2} \right\}, \\ N_2 &= \left\{ u \in (N \setminus N_1) \mid |N^+(u) \cap (\bar{N} \cup N_1)| \geq 2\sqrt{a}\Delta \right\}, \text{ and} \\ N_3 &= N \setminus (N_1 \cup N_2). \end{aligned}$$

Observe first that

$$\Delta^2 \geq \sum_{u \in N \setminus \{v\}} d^+(u) = m^+(v) + |A(N, \bar{N})|.$$

Since $m^+(v) > (1-a)\Delta(\Delta-1)$, we obtain

$$|A(N, \bar{N})| < \Delta^2 - (1-a)\Delta(\Delta-1) = a\Delta^2 + (1-a)\Delta.$$

By definition of N_1 , this implies

$$\frac{\Delta}{2}|N_1| < a\Delta^2 + (1-a)\Delta.$$

Using the fact that $\Delta \geq \Delta(a) \geq \frac{1-a}{\sqrt{a}-a}$, we thus have $|N_1| < 2\sqrt{a}\Delta$. By definition, $v \notin N_1$ as $N^+(v) \subseteq N$. We also have $v \notin N_2$ as $N^+(v) \cap (\bar{N} \cup N_1) = N_1$, and $|N_1| < 2\sqrt{a}\Delta$. We will now bound the size of N_2 . Recall that every vertex $u \in N_1$ satisfies $2|N^+(u) \cap \bar{N}| \geq \Delta \geq d^-(u)$. We thus have the following.

$$\begin{aligned} |A(N_2 \cup N_3, \bar{N} \cup N_1)| &= |A(N, \bar{N})| + |A(N_2 \cup N_3, N_1)| - |A(N_1, \bar{N})| \\ &\leq |A(N, \bar{N})| + \sum_{u \in N_1} d^-(u) - \sum_{u \in N_1} |N^+(u) \cap \bar{N}| \\ &\leq |A(N, \bar{N})| + \sum_{u \in N_1} |N^+(u) \cap \bar{N}| \\ &= |A(N, \bar{N})| + |A(N_1, \bar{N})| \\ &\leq 2|A(N, \bar{N})| \\ &< 2a\Delta^2 + 2(1-a)\Delta. \end{aligned}$$

By definition of N_2 , this implies

$$2\sqrt{a}\Delta|N_2| < 2a\Delta^2 + 2(1-a)\Delta,$$

which together with $\Delta \geq \Delta(a) \geq \frac{1-a}{\sqrt{a}-a}$ implies $|N_2| < 2\sqrt{a}\Delta$.

Let $k = \max(\bar{\chi}(D-v), \lfloor (1-\varepsilon)(\Delta+1) \rfloor)$. Since $v \in N_3$, $D - N_3$ is k -dicolourable, as it is a subdigraph of $D - v$. Let α be such a k -dicolouring and L be the list assignment of $D[N_3]$ defined as follows

$$L(u) = [k] \setminus \alpha(N^+(u) \setminus N_3).$$

We are now going to prove that $D[N_3]$ is L -dicolourable, implying that D has dichromatic number at most k (it is straightforward to check that any L -dicolouring of $D[N_3]$ extends α into a dicolouring

of D). First note that, by construction, every vertex $u \in N_3$ satisfies $|N^+(u) \setminus N_3| < 2\sqrt{a}\Delta + |N_2| < 4\sqrt{a}\Delta$. Hence every vertex $u \in N_3$ satisfies

$$|L(u)| > k - 4\sqrt{a}(\Delta + 1) \geq (1 - \varepsilon)(\Delta + 1) - 1 - 4\sqrt{a}(\Delta + 1) \geq \frac{5}{6}(\Delta + 1) - 1.$$

As $|L(u)|$ is an integer, we thus have $|L(u)| \geq \lfloor \frac{5}{6}(\Delta + 1) \rfloor$. Let M be a maximum matching in $\bar{D}[N_3]$ (the complementary of $D[N_3]$). Let X be the vertices of N_3 that are not covered M . By definition, X is a biclique of D so $|X| \leq \frac{2}{3}(\Delta + 1)$. Note that $|N_3| = 2|M| + |X|$. If $|M| \leq \frac{1}{6}(\Delta + 1)$ then $|M| + |X| \leq \frac{5}{6}(\Delta + 1)$. If $|M| \geq \frac{1}{6}(\Delta + 1)$ then $|M| + |X| = |N_3| - |M| \leq \frac{5}{6}(\Delta + 1)$.

In both cases, we obtain $|M| + |X| \leq \frac{5}{6}(\Delta + 1)$. Since $D[N_3]$ is a subdigraph of a digraph obtained from $\overleftarrow{K}_{|N_3|}$ by removing a matching of size $|M|$, $D[N_3]$ is $(|M| + |X|)$ -dichoosable by Lemma 5.3. Therefore $D[N_3]$ must be $\lfloor \frac{5}{6}(\Delta + 1) \rfloor$ -dichoosable, so in particular it is L -dicolourable, implying the result. \square

6 Proof of the main result

This section is devoted to the proof of our main result, namely Theorem 1.6. We need the following intermediate result proved by the second author [28]. Recall that an oriented graph is a digraph without any digon.

Theorem 6.1 (Picasarri-Arrieta [28]). *Let D be an oriented graph, then $\vec{\chi}(D) \leq \max(2, \Delta_{\min}(D))$.*

We are now going to prove Theorem 1.6. Let us first restate it.

Theorem 1.6. *There exists $\varepsilon > 0$ such that every digraph D satisfies*

$$\vec{\chi}(D) \leq \lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \vec{\omega}(D) \rceil.$$

Proof. Let Δ_1 be as in Corollary 3.2 and, for a real number $a > 0$, let $\Delta(a)$ be as in Theorem 1.9. We define a , ε_0 , Δ_2 , ε , and γ_ε as follows:

- $a = \frac{1}{600}$,
- $\varepsilon_0 = \frac{1}{6} - 4\sqrt{a} > 0$,
- $\Delta_2 = \min \left\{ \Delta \in \mathbb{N} \mid a > \frac{\log^3 \Delta}{\Delta - 1} \right\}$,
- $\varepsilon = \min \left(\frac{1}{\Delta_1}, \frac{1}{\Delta_2}, \frac{1}{\Delta(a)}, \frac{1}{3}\varepsilon_0, \frac{a}{16e^7} \right)$, and
- $\gamma_\varepsilon = \frac{\varepsilon}{1 - \varepsilon} \leq 2\varepsilon$.

We claim that the statement holds for this specific value of ε . Assume this is not the case, and let D be a counterexample with minimum order. For the sake of conciseness let us denote $\tilde{\Delta}(D)$ by $\tilde{\Delta}$, $\Delta_{\max}(D)$ by Δ_{\max} , $\vec{\omega}(D)$ by $\vec{\omega}$, and let $k = \lceil (1 - \varepsilon)(\tilde{\Delta} + 1) + \varepsilon \vec{\omega} \rceil$. As D is a minimum counterexample, it satisfies $\vec{\chi}(D) > k$ and for every vertex $v \in V$ we have $\vec{\chi}(D - v) \leq k$. This directly implies that every vertex v of D satisfies $\min(d^-(v), d^+(v)) \geq k$, for otherwise a k -dicolouring of D can be obtained from any k -dicolouring of $D - v$ by choosing for v a colour that is not appearing either in its in-neighbourhood or in its out-neighbourhood. In particular, this implies that $\tilde{\Delta}$ must be close to Δ_{\max} , as we justify in the following claim.

Claim 6.2. $\Delta_{\max} \leq (1 + \gamma_\varepsilon)\tilde{\Delta}$.

Proof of claim. Assume this is not the case, so there exists a vertex v such that $\max(d^+(v), d^-(v)) > (1 + \gamma_\varepsilon)\tilde{\Delta}$. By directional duality, we assume without loss of generality that $d^+(v) \geq d^-(v)$. Therefore, we have $\tilde{\Delta}^2 \geq d^+(v) \cdot d^-(v) > (1 + \gamma_\varepsilon)\tilde{\Delta} \cdot d^-(v)$, which implies $d^-(v) < \frac{1}{1+\gamma_\varepsilon}\tilde{\Delta} = (1 - \varepsilon)\tilde{\Delta} \leq k$, a contradiction. \diamond

With Theorem 4.2 in hands, we can also justify that there exists a significant gap between $\vec{\omega}$ and Δ_{\max} .

Claim 6.3. $\vec{\omega} \leq \frac{2}{3}(\Delta_{\max} + 1)$.

Proof of claim. Assume for a contradiction that $\vec{\omega} > \frac{2}{3}(\Delta_{\max} + 1)$. By Theorem 4.2, there exists an acyclic set of vertices $I \subseteq V(D)$ which intersects every maximum biclique of D . Among all such sets I , we choose one with maximum size. In particular, this implies that every vertex in $V(D) \setminus I$ has at least one in-neighbour and one out-neighbour in I . Let D' be $D - I$. As I is acyclic and intersects every maximum clique of D , we have $\vec{\omega}(D') = \vec{\omega} - 1$. The following shows that we also have $\tilde{\Delta}(D') \leq \tilde{\Delta} - 1$.

$$\begin{aligned} \tilde{\Delta}(D') &= \max \left\{ \sqrt{d_{D'}^+(v) \cdot d_{D'}^-(v)} \mid v \in V(D') \right\} \\ &\leq \max \left\{ \sqrt{(d_D^+(v) - 1)(d_D^-(v) - 1)} \mid v \in V(D') \right\} \\ &\leq \max \left\{ \sqrt{d_D^+(v)d_D^-(v)} - 1 \mid v \in V(D') \right\} \\ &\leq \tilde{\Delta} - 1, \end{aligned}$$

where in the first inequality we used the maximality of I and in the second inequality we used that $\sqrt{(p-1)(q-1)} \leq \sqrt{pq} - 1$ holds for every pair of real numbers $p, q \geq 1$. We briefly justify it here for completeness.

$$\begin{aligned} &0 \leq (\sqrt{p} - \sqrt{q})^2 \\ \Rightarrow & -p - q \leq -2\sqrt{pq} \\ \Rightarrow & pq - p - q + 1 \leq pq - 2\sqrt{pq} + 1 \\ \Rightarrow & (p-1)(q-1) \leq (\sqrt{pq} - 1)^2 \\ \Rightarrow & \sqrt{(p-1)(q-1)} \leq \sqrt{pq} - 1 \end{aligned}$$

Since I is acyclic, we have $\vec{\chi}(D) \leq \vec{\chi}(D') + 1$, as any dicolouring of D' can be extended to D by colouring I with an additional colour. Together with the minimality of D we obtain

$$\begin{aligned} \vec{\chi}(D) &\leq \left[(1 - \varepsilon)(\tilde{\Delta}(D') + 1) + \varepsilon\vec{\omega}(D') \right] + 1 \\ &\leq \left[(1 - \varepsilon)\tilde{\Delta} + \varepsilon(\vec{\omega} - 1) \right] + 1 \\ &= \left[(1 - \varepsilon)(\tilde{\Delta} + 1) - (1 - \varepsilon) + \varepsilon\vec{\omega} - \varepsilon + 1 \right] = k, \end{aligned}$$

a contradiction. \diamond

Our last claim guarantees that $\tilde{\Delta}$ is large enough.

Claim 6.4. $\Delta_{\max} \geq \tilde{\Delta} \geq \max(\Delta_1, \Delta_2, \Delta(a))$.

Proof of claim. It is clear from the definitions that $\Delta_{\max} \geq \tilde{\Delta}$. Assume for a contradiction that $\tilde{\Delta} < \max(\Delta_1, \Delta_2, \Delta(a))$. Since $\varepsilon \leq \min\left(\frac{1}{\Delta_1}, \frac{1}{\Delta_2}, \frac{1}{\Delta(a)}\right)$ we have $\varepsilon < \frac{1}{\tilde{\Delta}}$, which implies:

$$k = \left\lceil \tilde{\Delta} + 1 + \varepsilon(\vec{\omega} - \tilde{\Delta} - 1) \right\rceil \geq \left\lceil \tilde{\Delta} + \frac{1}{\tilde{\Delta}}(\vec{\omega} - 1) \right\rceil.$$

Using the facts that $\vec{\chi}(D) > k$ and that $\vec{\chi}(D) \leq \lceil \tilde{\Delta} + 1 \rceil$, we deduce

$$\lceil \tilde{\Delta} + 1 \rceil > \left\lceil \tilde{\Delta} + \frac{1}{\tilde{\Delta}}(\vec{\omega} - 1) \right\rceil.$$

This is a contradiction unless $\vec{\omega} = 1$ and $\tilde{\Delta}$ is an integer. Henceforth we assume that $\vec{\omega} = 1$, *i.e.* D is an oriented graph, and $\tilde{\Delta} \in \mathbb{N}$. In particular, we have $k \geq \tilde{\Delta}$. As D is an oriented graph, by Theorem 6.1 we must have $\Delta_{\min}(D) \leq 1$, for otherwise $\vec{\chi}(D) \leq \Delta_{\min}(D) \leq \tilde{\Delta} \leq k$, a contradiction. If $\tilde{\Delta} = 1$ then $k = \lceil 2 - \varepsilon \rceil = 2$. If $\tilde{\Delta} = 0$ then $k = 1$. Both cases yield a contradiction, which concludes the proof of the claim. \diamond

We now distinguish two cases, depending on the sparseness of D .

Case 1 *There exists a vertex v such that $\max(m^-(v), m^+(v)) \geq (1-a)\Delta(\Delta-1)$.*

By Claim 6.3 we have $\vec{\omega} \leq \frac{2}{3}(\Delta_{\max} + 1)$. As $\varepsilon_0 = \frac{1}{6} - 4\sqrt{a}$, by Theorem 1.9 we have

$$\vec{\chi}(D) \leq \max(\vec{\chi}(D-v), (1-\varepsilon_0)(\Delta_{\max} + 1)).$$

Recall that $\vec{\chi}(D-v) \leq k$ as D is a minimum counterexample. Since $\vec{\chi}(D) > k$, we thus necessarily have

$$\vec{\chi}(D) \leq (1-\varepsilon_0)(\Delta_{\max} + 1) \leq (1-\varepsilon_0)(1+\gamma_\varepsilon)(\tilde{\Delta} + 1) \leq (1-\varepsilon)(\tilde{\Delta} + 1),$$

where in the second inequality we used Claim 6.2, and in the last inequality we used $\varepsilon \leq \varepsilon_0 - \gamma_\varepsilon$, which holds because $\gamma_\varepsilon \leq 2\varepsilon$ and $\varepsilon \leq \frac{1}{3}\varepsilon_0$. This yields the contradiction as $(1-\varepsilon)(\tilde{\Delta} + 1) \leq k$.

Case 2 *For every vertex $v \in V(D)$ we have $\max(m^-(v), m^+(v)) \leq (1-a)\Delta(\Delta-1)$.*

A fortiori every vertex satisfies $\min(m^-(v), m^+(v)) \leq (1-a)\Delta(\Delta-1)$. By Claim 6.4 we know that $\Delta_{\max} \geq \max(\Delta_1, \Delta_2)$. Hence, by definition of Δ_2 , we have $a > \frac{\log^3 \Delta}{\Delta-1}$. We can thus apply Corollary 3.2, which together with Claim 6.2 gives

$$\vec{\chi}(D) \leq \left(1 - \frac{a}{5e^7}\right) (\Delta_{\max} + 1) \leq \left(1 - \frac{a}{5e^7}\right) (1 + \gamma_\varepsilon)(\tilde{\Delta} + 1) \leq (1-\varepsilon)(\tilde{\Delta} + 1).$$

where in the last inequality we used $\varepsilon \leq \frac{a}{5e^7} - \gamma_\varepsilon$, which holds because $\gamma_\varepsilon \leq 2\varepsilon$ and $\varepsilon \leq \frac{a}{15e^7}$. This contradicts $\vec{\chi}(D) > k$ and concludes the proof. \square

7 Further research: analogues for Δ_{\min}

In this work we proved an analogue of Reed's result for digraphs by taking the maximum degree of a digraph to be $\tilde{\Delta}$. A natural question then arises: can we strengthen this result by replacing $\tilde{\Delta}$ with Δ_{\min} ? The answer to this question turns out to be negative. Moreover, for every fixed real numbers $\varepsilon, \varepsilon'$ with $0 < \varepsilon' < 2\varepsilon$, there exists a digraph D with

$$\vec{\chi}(D) > \lceil (1 - \varepsilon)(\Delta_{\min}(D) + 1) + \varepsilon' \Delta_{\min}(D) \rceil.$$

This is because the k -DICOLOURABILITY problem is NP-complete for every fixed $k \geq 2$, even when restricted to digraphs D satisfying $\Delta_{\min}(D) = k$ and $\vec{\omega}(D) = \lceil \frac{k+1}{2} \rceil$, as shown in [28]. In particular, the reduction shows that there exist such digraphs satisfying $\vec{\chi}(D) = k + 1$. Hence, if we choose k to be at least $\frac{2-2\varepsilon+2\varepsilon'}{2\varepsilon-\varepsilon'}$, we deduce the existence of digraphs D with $\Delta_{\min}(D) = k$ and $\vec{\chi}(D) = k + 1$, while $\lceil (1 - \varepsilon)(k + 1) + \varepsilon'k \rceil \leq k$.

However, we believe that an analogue of our result holds with Δ_{\min} instead of $\tilde{\Delta}$ if we also adapt the notion of clique number. Given a digraph D , a *directed clique* of D is a set of vertices X that can be partitioned into (X_1, X_2) such that both $D[X_1]$ and $D[X_2]$ are complete digraphs, and D contains every possible arc uv with $u \in X_1$ and $v \in X_2$. The *directed clique number* of D , denoted by $\vec{\omega}(D)$, is the size of the largest directed clique of D . We propose the following conjecture.

Conjecture 7.1. *There exists $\varepsilon > 0$ such that every digraph D satisfies*

$$\vec{\chi}(D) \leq \lceil (1 - \varepsilon)\Delta_{\min}(D) + \varepsilon\vec{\omega}(D) \rceil.$$

In particular, the conjecture above implies the following slightly weaker one, which if true shows that the condition $\varepsilon' < 2\varepsilon$ mentioned above is indeed necessary.

Conjecture 7.2. *There exists $\varepsilon > 0$ such that every digraph D satisfies*

$$\vec{\chi}(D) \leq \lceil (1 - \varepsilon)\Delta_{\min}(D) + 2\varepsilon\vec{\omega}(D) \rceil.$$

We are inclined to believe that these two conjectures are true even for $\varepsilon = \frac{1}{2}$. One direction for proving Conjecture 7.1 would be to prove an analogue of Theorem 1.6 using the maximum out-degree Δ^+ instead of $\tilde{\Delta}$, that is to prove the existence of $\varepsilon > 0$ such that every digraph D satisfies

$$\vec{\chi}(D) \leq \lceil (1 - \varepsilon)(\Delta^+(D) + 1) + \varepsilon\vec{\omega}(D) \rceil,$$

where $\Delta^+(D) = \max_{v \in V(D)}(d^+(v))$. If such a result holds, then the following proposition, the proof of which uses a trick introduced in [28], implies Conjecture 7.1.

Proposition 7.3. *For every digraph D , there exists a digraph H such that $\vec{\chi}(D) \leq \vec{\chi}(H)$, $\Delta^+(H) \leq \Delta_{\min}(D)$ and $\vec{\omega}(H) \leq \vec{\omega}(D)$.*

Proof. Let us fix a digraph D , and let (X, Y) be the partition of $V(D)$ where $X = \{x \in V(D) \mid d_D^+(x) \leq \Delta_{\min}(D)\}$ and $Y = V(D) \setminus X$. In particular, every vertex $y \in Y$ satisfies $d^-(y) \leq \Delta_{\min}(D)$.

Let D' be the digraph obtained from D by removing every arc from Y to X and replacing every arc from X to Y by a digon. Let H be the digraph obtained from D' by reversing every arc of $D'[Y]$. It is straightforward to check that, by construction, $\Delta^+(H) \leq \Delta_{\min}(D')$ and $\vec{\omega}(H) \leq \vec{\omega}(D)$.

We now justify that $\vec{\chi}(H) \geq \vec{\chi}(D)$. Assume that this is not the case, and let ϕ be a $(\vec{\chi}(D) - 1)$ -dicolouring of H . By definition, D coloured with ϕ must contain a monochromatic directed cycle

$C = v_1, \dots, v_\ell, v_1$. Observe that $V(C) \not\subseteq X$, otherwise C is a directed cycle of H . In a similar way, $V(C) \not\subseteq Y$, for otherwise $v_1, v_\ell, \dots, v_2, v_1$ is a directed cycle of H . Hence C contains an arc xy from X to Y , but $\{xy, yx\}$ is a digon of H , a contradiction. \square

It is of interest that already the case of oriented graphs is open.

Problem 7.4. Show the existence of $\varepsilon > 0$ such that every oriented graph D with sufficiently large maximum out-degree $\Delta^+(D)$ satisfies $\bar{\chi}(D) \leq (1 - \varepsilon)\Delta^+(D)$.

The hardness of this problem mainly comes from the fact that the maximum in-degree of a digraph can be arbitrarily larger than its maximum out-degree. Hence, the proof of Theorem 1.7 cannot be directly adapted as we cannot bound the degree of dependence of bad events in order to apply the Lovász Local Lemma.

References

- [1] P. Aboulker and G. Aubian. Four proofs of the Directed Brooks' Theorem. *Discrete Mathematics*, page 113193, 2022.
- [2] P. Aboulker, G. Aubian, and P. Charbit. Digraph colouring and arc-connectivity. *preprint arXiv:2304.04690*, 2023.
- [3] P. Aboulker, G. Aubian, P. Charbit, and S. Thomassé. (P6, triangle)-free digraphs have bounded dichromatic number. *preprint arXiv:2212.02272*, 2022.
- [4] P. Aboulker, P. Charbit, and R. Naserasr. Extension of Gyárfás-Sumner conjecture to digraphs. *The Electronic Journal of Combinatorics*, 28(P2.27), 2021.
- [5] N. Alon and J. H. Spencer. *The probabilistic method*. Wiley Series in Discrete Mathematics and Optimization. Wiley-Blackwell, Hoboken, NJ, 3 edition, Aug. 2008.
- [6] S. D. Andres and W. Hochstättler. Perfect digraphs. *Journal of Graph Theory*, 79(1):21–29, 2015.
- [7] J. Bang-Jensen and G. Z. Gutin. *Digraphs: Theory, Algorithms and Applications*. Springer-Verlag, London, 2nd edition, 2009.
- [8] D. Bokal, G. Fijavz, M. Juvan, P. M. Kayll, and B. Mohar. The circular chromatic number of a digraph. *Journal of Graph Theory*, 46(3):227–240, 2004.
- [9] M. Bonamy, T. Perrett, and L. Postle. Colouring graphs with sparse neighbourhoods: Bounds and applications. *Journal of Combinatorial Theory, Series B*, 155:278–317, 2022.
- [10] D. Christofides, K. Edwards, and A. D. King. A note on hitting maximum and maximal cliques with a stable set. *Journal of Graph Theory*, 73(3):354–360, 2013.
- [11] L. Cook, T. Masařík, M. Pilipczuk, A. Reinald, and U. S. Souza. Proving a directed analogue of the Gyárfás-Sumner conjecture for orientations of P_4 . *The Electronic Journal of Combinatorics*, 30(P3.36), 2023.

- [12] P. Erdős. Problems and results in number theory and graph theory. In *Proceedings of the ninth Manitoba Conference on Numerical Mathematics and Computing*, pages 3–21, 1979.
- [13] P. Erdős and L. Lovász. Problems and results on 3-chromatic hypergraphs and some related questions. *Infinite and Finite Sets*, 10(2):609–627, 1975.
- [14] P. Erdős, A. L. Rubin, and H. Taylor. Choosability in graphs. *Congr. Numer.*, 26(4):125–157, 1979.
- [15] D. Gonçalves, L. Picasarri-Arrieta, and A. Reinald. Brooks-type colourings of digraphs in linear time. *preprint arXiv:2405.05222*, 2024.
- [16] A. Hajnal. A theorem on k -saturated graphs. *Canadian Journal of Mathematics*, 17:720–724, 1965.
- [17] A. Harutyunyan and B. Mohar. Gallai’s theorem for list coloring of digraphs. *SIAM Journal on Discrete Mathematics*, 25(1):170–180, 2011.
- [18] A. Harutyunyan and B. Mohar. Strengthened Brooks’ theorem for digraphs of girth at least three. *The Electronic Journal of Combinatorics*, 18(P195), 2011.
- [19] P. E. Haxell. A note on vertex list colouring. *Combinatorics, Probability and Computing*, 10(4):345–347, 2001.
- [20] E. Hurley, R. de Joannis de Verclos, and R. J. Kang. An improved procedure for colouring graphs of bounded local density. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 135–148. SIAM, 2021.
- [21] T. Kelly and L. Postle. A local epsilon version of Reed’s Conjecture. *Journal of Combinatorial Theory, Series B*, 141:181–222, 2020.
- [22] A. D. King. Hitting all maximum cliques with a stable set using lopsided independent transversals. *Journal of Graph Theory*, 67(4):300–305, 2011.
- [23] A. D. King and B. A. Reed. A short proof that χ can be bounded ε away from $\Delta + 1$ toward ω . *Journal of Graph Theory*, 81(1):30–34, Jan. 2015.
- [24] A. V. Kostochka. Degree, density and chromatic number of graphs. *Metody Diskret. Analiz.*, 35:45–70, 1980.
- [25] B. Mohar. Circular colorings of edge-weighted graphs. *Journal of Graph Theory*, 43(2):107–116, 2003.
- [26] M. Molloy and B. A. Reed. *Graph colouring and the probabilistic method*. Algorithms and Combinatorics. Springer, Berlin, Germany, Nov. 2001.
- [27] V. Neumann-Lara. The dichromatic number of a digraph. *Journal of Combinatorial Theory, Series B*, 33:265–270, 1982.
- [28] L. Picasarri-Arrieta. Strengthening the Directed Brooks’ Theorem for oriented graphs and consequences on digraph redicolouring. *Journal of Graph Theory*, 106(1):5–22, 2024.

- [29] L. Rabern. On hitting all maximum cliques with an independent set. *Journal of Graph Theory*, 66(1):32–37, 2011.
- [30] B. A. Reed. ω , Δ , and χ . *Journal of Graph Theory*, 27(4):177–212, 1998.
- [31] M. Stiebitz, T. Schweser, and B. Toft. *Brooks' Theorem: Graph Coloring and Critical Graphs*. Springer Nature Switzerland, 2024.
- [32] M. Talagrand. Concentration of measure and isoperimetric inequalities in product spaces. *Publications Mathématiques de l'Institut des Hautes Etudes Scientifiques*, 81:73–205, 1995.