

Universality of the mean-field equations of networks of Hopfield-like neurons

Olivier C Faugeras, Etienne Tanré

To cite this version:

Olivier C Faugeras, Etienne Tanré. Universality of the mean-field equations of networks of Hopfieldlike neurons. $2024.$ hal- $04677247v2$

HAL Id: hal-04677247 <https://inria.hal.science/hal-04677247v2>

Preprint submitted on 26 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

UNIVERSALITY OF THE MEAN-FIELD EQUATIONS OF NETWORKS OF HOPFIELD-LIKE NEURONS

OLIVIER FAUGERAS[∗] AND ETIENNE TANRE´†

CONTENTS

^{*}UNIVERSITÉ CÔTE D'AZUR, CNRS, LJAD

[†]UNIVERSITÉ CÔTE D'AzUR, INRIA, CNRS, LJAD

Date: August 23, 2024.

Abstract. We revisit the problem of characterising the mean-field limit of a network of Hopfield-like neurons. Building on the previous work of [13, 1, 14] and [9] we establish for a large class of networks of Hopfield-like neurons, i.e. rate neurons, the mean-field equations on a time interval [0, T], $T > 0$, of the thermodynamic limit of these networks, i.e. the limit when the number of neurons goes to infinity. Unlike all previous work, except [9], we do not assume that the synaptic weights describing the connections between the neurons are i.i.d. as zero-mean Gaussians. The limit equations are stochastic and very simply described in terms of two functions, a "correlation" function noted $K_Q(t, s)$ and a "mean" function noted $m_Q(t)$. The "noise" part of the equations is a linear function of the Brownian motion, which is obtained by solving a Volterra equation of the second kind whose resolving kernel is expressed as a function of K_Q . We give a constructive proof of the uniqueness of the limit equations. We use the corresponding algorithm for an effective computation of the functions K_Q and m_Q , given the weights distribution. Several numerical experiments are reported.

Keywords: Large deviations; Girsanov transform; Hopfield Neural network; Interacting Particle System.

MSC2020 AMS classification: 60F10, 92B20, 60B10

1. INTRODUCTION

We revisit the problem of characterizing the limit of a network of Hopfield neurons. Hopfield [15] defined a large class of neuronal networks and characterized some of their computational properties [16, 17], i.e. their ability to perform computations.

Inspired by his work, Sompolinsky and co-workers studied the thermodynamic limit of these networks when the interaction term is linear [7] using the dynamic mean-field theory developed in [25] for symmetric spin glasses. The method they use is a functional integral formalism developed in particle physics and produces the self-consistent mean-field equations of the network. This was later extended to the case of a nonlinear interaction term, the nonlinearity being an odd sigmoidal function [24]. Using the same formalism the authors established the self-consistent mean-field equations of the network and the dynamics of its solutions which featured a chaotic behavior for some values of the network parameters. More recently, a derivation of the equations using the path integral formalism was provided by the same authors [8].

A little later the problem was picked up again independently by mathematicians, although in a different context. Ben Arous and Guionnet applied large deviation techniques to study the thermodynamic limit of a network of spins interacting linearly through i.i.d. centered Gaussian weights. The intrinsic spin dynamics (without interactions) is a stochastic differential equation, unlike in the case of the work of Sompolinsky and collaborators where the only source of uncertainty lies in the interaction weights. Ben Arous and Guionnet proved that the annealed

(averaged) law of the empirical measure satisfies a large deviation principle and that the good rate function of this large deviation principle achieves its minimum value at a unique non Markovian measure [13, 1, 14]. They also proved averaged propagation of chaos results. The advance provided by the work of Ben Arous and Guionnet is, beyond the crucial fact that they set the problem on firm mathematical ground, that they provide a description of the limit dynamics in a much more precise way than all previous authors.

Moynot and Samuelides [19] adapted their work to the case of a network of Hopfield neurons with a nonlinear interaction term, the nonlinearity being a sigmoidal function, and proved similar results in the case of discrete time. The intrinsic neural dynamics is the gradient of a quadratic potential. They did not provide a description of the limit dynamics akin to the one of Ben Arous and Guionnet.

Even more recently, van Meegen, Kühn and Helias, in [27] nicely unified the Large Deviation approach of Ben Arous and Guionnet with the field-theoretical one of Sompolinsky and collaborators by showing that the Large Deviation rate function is in effect equal to the effective action arising in the field theory. They did not deal correctly with the limit dynamics though in the sense that they did not state in the limit equations the dependency of the effective interaction term with the Brownian motion.

Adapting the work of Ben Arous and Guionnet, Cabana and Touboul, in a series of papers [4, 5, 3], have proposed a number of generalisations to more complicated setups. They applied the theory of Large Deviations and showed the existence of a unique probability measure where the rate function is zero, they identified the corresponding measure but did not provide a pathwise description of this probability measure. In effect their description, like that of all previous authors, is based upon a certain process called the effective interaction process which is dependent of the Brownian motion under the limit law but this dependency is not analysed.

In [9] the authors proved that the result in [13, 1, 14] is in fact universal, i.e. that their limit probability measure is independent of the distribution of the weights under some mild assumptions.

In this article we show how to go beyond the "self-consistent" approach and provide a complete description of the effective interaction term as a function of the Brownian motion under the limit law. Thanks to the result in [9] our results also hold for a much larger class of synaptic weights distributions than the Gaussian case.

The paper is organised as follows. In Section 2 we present the classes of spin and neuron models that we analyse and the network equations that govern their activity. We then state the problem at hand, namely the characterisation of these network equations when the network size grows arbitrarily large, the thermodynamic limit. In Section 3 we recall the two main results which inspired our work and briefly state our three main results. In Section 4 we state and prove our

first two main results, the stochastic equations of the dynamics of the thermodynamic limit of the networks and the characterisation of the diffusion part of these equations. Section 5 is a pause to analyse the special case when the interaction between the neurons is constant, albeit random, not depending on the states of the neurons. Section 6 contains the proof of our third main result, the uniqueness of the thermodynamic limit. Being constructive this proof opens up the door to numerical experiments many of them presented in Section 7.

2. The models

We consider a filtered probability space $(\Omega, \mathcal{G}, (\mathcal{G}_t), p)$ equipped with a sequence of one-dimensional Brownian motions $(B_t^i)_{t\geq 0, i\in\mathbb{N}}$.

The object of interest is a fully connected network of Hopfield-like neurons, in short Hopfield neurons. A Hopfield neuron is an abstraction of a real neuron where the state of the neuron is described by the values of its membrane potential. The dynamics of an isolated Hopfield neuron is driven by a certain stochastic differential equation where the "noise" is assumed to account for the probabilistic effects present in various biological actors within the neuron, such as the ion channels. Our view of a Hopfield neuron is rather large since it encompasses the spin-glass model of Ben Arous and Guionnet [13, 1, 14]. When we will need to differentiate between the two models we will refer to the H-model for the biological one and to the S-model for the spin-glass model. This is described in Section 2.1. Hopfield neurons can interact, one says that they are connected. The biological support of these interactions is the notion of synaptic connection. In Hopfield neurons, the synaptic connection between a neuron i and a neuron i is extremely simplified and boils down to the product of two numbers, the synaptic weight $J_{i \leftarrow j}$ that represents the maximum strength of the influence of j over i , and a function of the membrane potential X^j of neuron j which represents its activity, its firing rate. This is described in Section 2.2.

2.1. Single neuron.

Consider a Hopfield neuron with membrane potential Y_t . Its dynamics is described by a stochastic differential equation

(1)
$$
\begin{cases} dY_t &= g(Y_t)dt + \lambda dB_t \\ \text{Law of } Y_0 &= \mu_0, \end{cases}
$$

where λ is a nonzero real parameter. The function g is chosen such that (1) has a unique strong solution which does not explode in finite time. In the cases studied in this paper we have $q(y) = -\nabla U(y)$.

Remark 1.

H-model: $U(y) = \frac{\alpha}{2}y^2$, i.e $g(y) = -\alpha y$, with $\alpha > 0$. This is the H-model where the membrane potential is an Orstein-Uhlenbeck process

$$
Y_t = Y_0 \exp(-\alpha t) + \lambda \int_0^t \exp(-\alpha (t-s))dB_s.
$$

S-model: U is a confining potential. This is the S-model studied by Ben Arous and Guionnet [1]. The probability law μ_0 has a compact support $[-A, A]$ for some $A > 1$ which does not put any mass on the boundary $\{-A, A\}$. The function U is C^2 on the interval $]-A, A[$, and U tends to infinity when $|x| \to A$ sufficiently fast to ensure that

$$
\lim_{\substack{|x| \to A \\ <}} k_U(x) = +\infty,
$$

where

(2)
$$
k_U(x) = 2 \int_0^x \exp 2U(y) \left(\int_0^y \exp -2U(z) dz \right) dy.
$$

An example of such a function U is

(3)
$$
U(y) = -\log(A^2 - y^2).
$$

It follows that

(4)
$$
g(y) = -\frac{2y}{A^2 - y^2}.
$$

Assuming that the support of the initial condition μ_0 is included in $(-A, A)$, it is recalled in [1] that the solution does not explode in finite time i.e. that if $T_{\varepsilon} = \inf\{t \mid |Y_t| \geq A - \varepsilon\}$, then for all $T, p(T_{\varepsilon} \leq T) \leq \frac{\exp T}{1 + k_{\text{tr}}(A)}$ $\frac{\exp T}{1+k_U(A-\varepsilon)}$ so that $p(\lim_{\varepsilon\downarrow 0}T_{\varepsilon}=+\infty)=1$. More details about explosion times can be found in the classical books by Rogers and Williams [23], Borodin [2], and Peskir and Shiryaev [21].

Figure 1 shows the functions $U(x)$ and $q(x)$ when $A = 2$.

FIGURE 1. S-model $(A = 2)$. Left: the function $U(x)$ on the interval $(-2, 2)$. Right: the function $g(x) = -\nabla U(x)$

Let us note P the law of the solution to (1). The previous remark shows that in the case $g(y) = -\alpha y$, P is a probability measure on the set \mathcal{C}_T of continuous functions on $[0, T]$ with values in R while in the second case it is a probability measure on the set \mathcal{C}_T^A of continuous functions on $[0,T]$ with values in $[-A, A]$. In both cases we note \mathcal{F}_t the σ -algebra generated by $(Y_s, 0 \le s \le t)$, and F the corresponding filtration. In the sequel we note $\mathcal C$ either $\mathcal C_T$ or $\mathcal C_T^A$ and P is a probability measure on $(C, \mathcal{F}_T, \mathbb{F})$.

2.2. Network of neurons.

We consider N Hopfield neurons and connect each one of them to all others (and to itself). The strengths of the connections are random, described by the synaptic weights. We introduce a second probability space $(\Theta, \mathcal{A}, \gamma)$ and i.i.d. random variables $(J_{i \leftarrow j})_{i,j=1,\cdots,N}$ on Θ which are, under γ , i.i.d. as

(5)
$$
J_{i \leftarrow j} \simeq \mathcal{N}\left(\frac{J}{N}, \frac{\sigma^2}{N}\right), i, j = 1, \cdots, N.
$$

We note J the $N \times N$ matrix of the $J_{i \leftarrow j}$ s and $\mathcal E$ the expected value with respect to the probability γ .

Remark 2. The hypothesis that the weights distribution is Gaussian is lifted in Section 3.2.

We consider a continuous function $f : \mathbb{R} \to \mathbb{R}$ which converts the values of the membrane potential into a quantity called the "activity" of the neuron. A classical example of such activity is the firing rate, i.e. the number of action potentials produced per second.

The network equations are

(6)
$$
\begin{cases} dX_t^i = \left(g(X_t^i) + \sum_{j=1}^N J_{i \leftarrow j} f(X_t^j) \right) dt + \lambda dB_t^i \quad \text{for } i = 1, \cdots, N \\ \text{Law of } X_0 = \mu_0^{\otimes N}, \end{cases}
$$

where the $(B_t^i)_{t\geq 0}$ are N independent Brownian motions on $(\Gamma, \mathcal{G}, (\mathcal{G}_t), p)$.

The S-model studied by Ben Arous and Guionnet features $f(y) = y$ while the $J_{i \leftarrow j}$ s are i.i.d. as $\mathcal{N}(0, \frac{\sigma^2}{N})$ $\frac{\sigma^2}{N}$, the so-called "soft spin glass" case. We extend this model to the case with non centered weights $J_{i \leftarrow j}$ in Section 4.

The H-model corresponds to the case $g(y) = -\alpha y$, f a sigmoidal function and the $J_{i\leftarrow j}$ are not necessarily centered. For future reference we assume that in this case

(7)
$$
|f(x)| \le C \quad \forall x \in \mathbb{R}
$$

Key to the approach in this paper is to relate the solution to N independent copies of (1) with that of (6). This relation is obtained through the application of the Cameron-Martin [6] and Girsanov [12] Theorems which we recall here for future reference.

Theorem 3 (Cameron-Martin). Let B be an N-dimensional Brownian motion on $(\Omega, \mathcal{G}, (\mathcal{G}_t), p)$. Let $\Phi = (\Phi^1, \cdots, \Phi^N)$ be in $\mathcal{L}^2([0,T]; \mathbb{R}^N)$. Define

$$
\zeta_0^T(\Phi, \omega) = \int_0^T \Phi(t) \, d\mathbf{B}_t - \frac{1}{2} \int_0^T \|\Phi(t, \omega)\|^2 \, dt,
$$

$$
\tilde{\mathbf{B}}_t(\omega) = \mathbf{B}_t(\omega) - \int_0^t \Phi(s, \omega) \, ds,
$$

$$
d\tilde{p}(\omega) = \exp\left\{\zeta_0^T(\Phi, \omega)\right\} dp(\omega)
$$

Then, if

$$
\tilde{p}(\Omega) = 1
$$

B is a new N-dimensional Brownian motion defined on $(\Omega, \mathcal{G}, \tilde{p})$ w.r.t. to the same filtration (\mathcal{G}_t) . Moreover a sufficient condition for (8) to hold is

$$
\mathbb{E}_p\left[\exp\left(\int_0^T \|\boldsymbol{\Phi}(t,\omega)\|^2 dt\right)\right] < \infty
$$

Remark 4. We recall that the quantity $\frac{d\tilde{p}}{dp}$ is called the Radon-Nikodym derivative of the probability measure \tilde{p} w.r.t. the probability measure p.

We also recall that $\mathcal{L}^2([0,T];\mathbb{R}^N)$ denotes the set of \mathbb{R}^N -valued \mathcal{G}_t -adapted processes $(X_t)_{0 \le t \le T}$ such that $\int_0^T ||X_t||^2 dt < \infty$ a.s..

Theorem 5 (Girsanov). Consider B an N-dimensional Brownian motion on $(\Omega, \mathcal{G}, (\mathcal{G}_t), p)$. Let $(X_t)_{0 \leq t \leq T}$ be the N-dimensional Itô process given by

$$
\mathbf{X}_t = \mathbf{X}_0 + \int_0^t \mathbf{a}(s) \, d\mathbf{B}_s + \int_0^t \mathbf{b}(s) \, ds
$$

where $\mathbf{a}, \mathbf{b} \in \mathcal{L}^2([0,T];\mathbb{R}^N)$ Let $\mathbf{\Phi} = (\Phi^1, \cdots, \Phi^N)$ be in $\mathcal{L}^2([0,T];\mathbb{R}^N)$. Let $\tilde{\mathbf{B}}$ and \tilde{p} be defined as in Theorem 3. If $\tilde{p}(\Omega) = 1$ then \mathbf{X}_t is still an Itô process on the probability space $(\Omega, \mathcal{G}, \tilde{p})$ w.r.t. the same filtration (\mathcal{G}_t) and satisfies

$$
\mathbf{X}_t = \mathbf{X}_0 + \int_0^t \mathbf{a}(s) d\tilde{\mathbf{B}}_s + \int_0^t \left(\mathbf{b}(s) + \mathbf{a}(s) \cdot \mathbf{\Phi}(s)\right) ds.
$$

As mentioned above, we consider a network of N non-interacting neurons, the state of each one being described by (1):

(9)
$$
\begin{cases} dY_t^i = g(Y_t^i)dt + \lambda dB_t^i & 1 \le i \le N \\ \text{Law of } Y_0 = \mu_0^{\otimes N}. \end{cases}
$$

The law of its solution is the probability measure $P^{\otimes N}$ on \mathcal{C}^N .

We relate (9) and (6) through Theorems 3 and 5.

Lemma 6. The system (6) has a unique weak solution. Its law, noted $P^{N}(\mathbf{J})$, is absolutely continuous with respect to the law $P^{\otimes N}$ of the uncoupled system. The Radon-Nikodym derivative is (10)

$$
\frac{dP^N(\mathbf{J})}{dP^{\otimes N}} = \exp\left\{\sum_i \frac{1}{\lambda} \int_0^T \sum_j J_{i \leftarrow j} f(Y_t^j) dB_t^i - \frac{1}{2\lambda^2} \int_0^T \left(\sum_j J_{i \leftarrow j} f(Y_t^j)\right)^2 dt\right),\right\}
$$

where the Y^j s are solutions to (9).

Proof. We apply Theorems 3 and 5. Let $\Phi \in \mathcal{L}^2([0,T];\mathcal{C}^N)$ and define

$$
\zeta_0^T = \sum_{i=1}^N \int_0^T \Phi^i(t) \, dB_t^i - \frac{1}{2} \int_0^T \Phi^i(t)^2 \, dt
$$

The Cameron-Martin Theorem 3 states that if

(11)
$$
\mathbb{E}_{P^{\otimes N}}\left[e^{\int_0^T \|\Phi(t)\|^2 dt}\right] < \infty,
$$

we can define for all $t \in [0, T]$

$$
\tilde{B}_t^i = B_t^i - \int_0^t \Phi^i(s) \, ds,
$$

and

$$
dP^{N}(\mathbf{J}) = \exp\left\{\zeta_0^{T}\right\} dP^{\otimes N},
$$

which are such that $P^{N}(\mathbf{J})$ is a probability measure on \mathcal{C}^{N} and $(\tilde{B}^{i})_{i}$ an Ndimensional Brownian motion under $P^{N}(\mathbf{J})$. Moreover the Girsanov Theorem states that the Y^i s satisfy also

$$
Y_t^i = Y_0 + \int_0^t \left(g(Y_s^i) + \lambda \Phi^i(s) \right) ds + \lambda \tilde{B}_t^i, i = 1, \cdots, N.
$$

The choice of

$$
\Phi^{i}(t) = \frac{1}{\lambda} \sum_{j=1}^{N} J_{i \leftarrow j} f(Y_{t}^{j}), \ i = 1, \cdots, N
$$

shows that, if (11) is satisfied, the Y^i s, which we rewrite X^i s, are solutions to (6) with the Brownian motion \tilde{B} under $P^N(\mathbf{J})$ and that the Radon-Nikodym derivative of $P^{N}(\mathbf{J})$ w.r.t. to $P^{\otimes N}$ is given by (10).

Condition (11) is clearly satisfied for the S-model since the $Yⁱ$ s take values in $]-A, A[$ and for the H-model since f is assumed to be bounded, see (7). This proves that (6) has a unique weak solution whose law is $P^{N}(\mathbf{J})$.

2.3. Statement of the problem.

The goal is to describe the asymptotic behaviour of the system (6) as N increases. Let X^i be the solution of (6), we denote $\hat{\mu}_N$ the associated empirical measure

$$
\hat{\mu}_N = \frac{1}{N} \sum_{i=1}^N \delta_{X^i}.
$$

The measure $\hat{\mu}_N$ is a probability measure on the set of trajectories C. Let us give an example of how it acts on test functions: Consider the function φ defined by

$$
\forall Y \in \mathcal{C}, \quad \varphi(Y) = f(Y_t)f(Y_s).
$$

The integral of φ w.r.t. $\hat{\mu}_N$ yields

(12)
$$
\langle \varphi, \hat{\mu}_N \rangle = \int \varphi(Y) d\hat{\mu}_N(Y) = \frac{1}{N} \sum_{i=1}^N f(X_t^i) f(X_s^i).
$$

3. Results

3.1. The results of Ben Arous and Guionnet [1, 14].

This is the case of the S-model where the weights are centered i.e. $J = 0$ in (5), and $\lambda = 1$. Note that our parameter σ , the standard deviation of the synaptic weights, is the same as their parameter β , the inverse temperature. Key to their approach is the averaging of the law $P^{N}(\mathbf{J})$ with respect to **J** and replacing the initial problem by that of studying the limit behaviour of the empirical measure under the averaged law Q^N given by

$$
Q^N = \int_{\Theta} P^N(\mathbf{J}(\theta)) d\gamma(\theta) = \mathcal{E}[P^N(\mathbf{J})]
$$

A summary of the essential results is as follows

- Existence and uniqueness of an averaged (annealed) limit: There exists a unique probability law Q on C such that the law of the empirical measure $\hat{\mu}_N$ under Q^N converges to δ_Q (Theorem 2.1 in [14]).
- Averaged (annealed) propagation of chaos: Q^N propagates chaos, or is Q-chaotic, i.e. given $m \geq 1$ and m continuous bounded functions $(\varphi_j)_{j=1,\dots,m}$ defined on C

$$
\lim_{N \to \infty} \mathbb{E}_{Q^N} \bigg[\prod_{j=1}^m \varphi_j(X^j) \bigg] = \prod_{j=1}^m \mathbb{E}_Q \bigg[\varphi_j(X) \bigg].
$$

This is not very useful in practice because the use of Q^N in the left hand side requires to average over the weights J but they also prove

Existence and uniqueness of a quenched limit: The law of the empirical measure of the quenched system, i.e. the law of $\hat{\mu}_N$ under $P^N(\mathbf{J})$, converges to δ_Q for almost all **J**s (Theorem 2.7 in [1]). This of course does not imply a quenched propagation of chaos since the neurons are not exchangeable for almost all weights but they also prove

Quenched propagation of chaos: If the function g in (1) and (6) is the gradient of an even function¹ and the initial law μ_0 is symmetric, then for any set of m continuous bounded functions $(\varphi_j)_{j=1,\dots,m}$ defined on C

$$
\mathbb{E}_{P^N(\mathbf{J})}\bigg[\prod_{j=1}^m \varphi_j(X^j)\bigg] \xrightarrow[N \to \infty]{p} \prod_{j=1}^m \mathbb{E}_Q\bigg[\varphi_j(X)\bigg].
$$

This means that for all $\varepsilon > 0$

$$
\lim_{N \to \infty} \gamma \left(\theta : \left| \mathbb{E}_{P^N(\mathbf{J})} \left[\prod_{j=1}^m \varphi_j(X^j) \right] - \prod_{j=1}^m \mathbb{E}_Q \left[\varphi_j(X) \right] \right| > \varepsilon \right) = 0.
$$

Characterisation of the limit law: The limit law Q is characterised by its Radon-Nikodym derivative w.r.t. to the law P

(13)
$$
\frac{dQ}{dP} = \int \exp\left\{ \int_0^T G_t^Q \, dB_t - \frac{1}{2} \int_0^T (G_t^Q)^2 \, dt \right\} \, d\gamma_Q,
$$

where G^Q is under γ_Q a centered Gaussian process with covariance function $K_Q(t,s) = \sigma^2 \mathbb{E}_Q[f(X_t)f(X_s)]$. Moreover this limit law is unique and solution of the following stochastic system

(14)
$$
\begin{cases} X_t &= X_0 + \int_0^t g(X_s) ds + B_t \\ B_t &= W_t + \int_0^t \int_0^s \tilde{K}_Q^s(s, u) dB_u ds \\ \text{Law of } X &= Q, \quad Q_{\mathcal{F}_0} = \mu_0 \\ K_Q(t, s) &= \sigma^2 \mathbb{E}_Q[f(X_t)f(X_s)], \end{cases}
$$

where W_t is a Brownian motion under Q .

The deterministic function \tilde{K}_{Q}^{t} is defined as follows: Let (G_t^Q) $_{t}^{Q})_{t\in[0,T]}$ be a centered Gaussian process with covariance K_Q , one defines

$$
\tilde{K}_Q^t(s, u) = \frac{\int_{\Theta} \exp\left\{-\frac{1}{2} \int_0^t (G_v^Q(\theta))^2 dv\right\} G_s^Q(\theta) G_u^Q(\theta) d\gamma_Q(\theta)}{\int_{\Theta} \exp\left\{-\frac{1}{2} \int_0^t (G_v^Q(\theta))^2 dv\right\} d\gamma_Q(\theta)} \quad \forall 0 \le s, u \le t,
$$

where G^Q is as before, under γ_Q , a centered Gaussian process with covariance K_Q . Note that this is equivalent to changing γ_Q into $\tilde{\gamma}_Q$ defined by

$$
d\tilde{\gamma}_Q^t(\theta) = \frac{\exp\left(-\frac{1}{2} \int_0^t (G_v^Q(\theta))^2 dv\right)}{\int_{\Theta} \exp\left(-\frac{1}{2} \int_0^t (G_v^Q(\theta))^2 dv\right) d\gamma_Q(\theta)} d\gamma_Q(\theta),
$$

¹Note that this condition is satisfied in the S- and H-models.

so that

$$
\tilde{K}_Q^t(s,u)=\int_{\Theta}G_s^Q(\theta)G_u^Q(\theta)\,d\tilde{\gamma}_Q^t(\theta).
$$

Equations (14) look a bit mysterious and our next goals are a) to generalise them to the non-zero mean weights case, i.e. $J \neq 0$ in (5), and b) to explore the relationship between the Brownian motions B_t and W_t as well as that between K_Q and \tilde{K}_{Q}^{t} .

3.2. The result of Dembo, Lubetzky and Zeitouni [9].

This is also the case of the S-model in which the weights are still assumed zero mean but without the assumption that their distribution is Gaussian. They prove that under the following mild assumptions

(15)
$$
\begin{cases} \mathcal{E}[J_{i \leftarrow j}] = 0, \ \mathcal{E}[J_{i \leftarrow j}^2] = \frac{\sigma^2}{N}, \\ \exists \varepsilon_0 > 0 \text{ such that } \sup_{i,j,N} \left\{ \mathcal{E}[e^{\varepsilon_0 \sqrt{N}|J_{i \leftarrow j}|}] \right\} < \infty, \end{cases}
$$

the empirical measure $\hat{\mu}_N$ converges weakly almost surely in the interactions **J** toward the **same** measure δ_Q as in the Gaussian case.

3.3. Statement of the results.

We have obtained three main results. The first one is a characterization of the dynamics of the trajectories of the limit measure Q , this is Theorem 7. The equations (19) of this dynamics contain the Brownian motion under P and the Brownian motion under Q which are related by an integral equation. Our second contribution is to solve this equation to obtain a simple relation between these two Brownian motions featuring functions of the covariance K_Q , this is Proposition 14. The third contribution is a proof of the uniqueness of the limit measure Q , this is Theorem 26. It is a constructive Picard-like method that can be used to effectively construct Q and simulate its trajectories. Our proof is different from the one in [1] which is valid for the S-model for zero-mean weights $(J = 0)$ and carries over to the nonzero-mean case. Note that, because of [9], our results are independent of the distribution of the $J_{i\leftarrow j}$ s, provided they satisfy the assumptions (15).

4. Non zero-mean weights

It is more relevant to neuroscience to consider the case where the weights are distributed as (5) with $J \neq 0$. This complicates quite a bit the limit dynamics as we show now.

If Q^N is the average of the law $P^N(\mathbf{J})$ given by (16) w.r.t. the weights and P the law of the uncoupled neuron one finds, using the independence property of the weights that

(16)
$$
\frac{dQ^N}{dP^{\otimes N}} = \int \prod_{i=1}^N \exp\left\{ \int_0^T \left(\frac{1}{\lambda} \sum_{j=1}^N J_{i \leftarrow j}(\theta) f(Y_t^j) \right) dB_t^i - \frac{1}{2} \int_0^T \left(\frac{1}{\lambda} \sum_{j=1}^N J_{i \leftarrow j}(\theta) f(Y_t^j) \right)^2 dt \right\} d\gamma(\theta),
$$

where the Y^j s in the above formula are the solutions of the uncoupled system

(17)
$$
dY_t^j = g(Y_t^j)dt + \lambda dB_t^j \qquad j = 1, \cdots, N
$$

and are therefore not functions of the $J_{i \leftarrow j}$ s.

If one defines

$$
G_t^i(\theta) = \sum_{j=1}^N J_{i \leftarrow j}(\theta) f(Y_t^j),
$$

one observes that, conditionally on $\mathcal{F}^{B^1,\cdots,B^N}_t$ and Y^1_0,\cdots,Y^N_0 , the G^i s, $i=1,\cdots,N$ are Gaussian processes whose mean and covariance are easily calculated using equations similar to (12):

$$
\mathcal{E}[G_t^i] = \frac{J}{N} \sum_{j=1}^N f(Y_t^j) = J \int f(Y_t) d\hat{\mu}_N(Y),
$$

and

$$
\mathcal{E}[G_t^i G_s^i] = \sigma^2 \int f(Y_t) f(Y_s) d\hat{\mu}_N(Y) + J^2 \int f(Y_t) d\hat{\mu}_N(Y) \times \int f(Y_s) d\hat{\mu}_N(Y),
$$

where $\hat{\mu}_N(Y)$ is the empirical measure

$$
\hat{\mu}_N(Y) = \frac{1}{N} \sum_{j=1}^N \delta_{Y^j}.
$$

Note that because the weights $J_{i \leftarrow j}$ are independent, the processes (G_t^i) and (G_t^j) $_{t}^{j}),$ $i \neq j$, are conditionally independent given $\mathcal{F}^{B^1,\cdots,B^N}_t$ and Y^1_0,\cdots,Y^N_0 .

Under γ , the law of each G_t^i is $\gamma_{\hat{\mu}_N}$, meaning that G_t^i , conditionally on $\mathcal{F}_t^{B^1,\dots,B^N}$ Let γ , the law of each G_t is $\gamma_{\hat{\mu}_N}$, inealling that G_t , conditionally on \mathcal{F}_t
 $\gamma_{\hat{\mu}_N}$ is a Caussian process with mean m_s , $(t) = I \int f(V) d\hat{\mu}_s$ and Y_0^1, \dots, Y_0^N , is a Gaussian process with mean $m_{\hat{\mu}_N}(t) = J \int f(Y_t) d\hat{\mu}_N(Y)$ J $\frac{J}{N}\sum_j f(Y^j_t)$ $\binom{7}{t}$, and covariance

$$
K_{\hat{\mu}_N}(t,s) = \sigma^2 \int f(Y_t) f(Y_s) d\hat{\mu}_N(Y) = \frac{\sigma^2}{N} \sum_{j=1}^N f(Y_t^j) f(Y_s^j).
$$

Having noticed this, it is possible to generalise the approach in [1, 14] and obtain that the limit law Q is characterised by the following modification of (13)

(18)
$$
\frac{dQ}{dP} = \int \exp\left\{\frac{1}{\lambda} \int_0^T G_t^Q dB_t - \frac{1}{2\lambda^2} \int_0^T (G_t^Q)^2 dt\right\} d\gamma_Q,
$$

where B is a P Brownian motion and, conditionally on \mathcal{F}_t^B and μ_0 , G_t^Q t_i^Q is under γ_Q a Gaussian process with mean $m_Q(t) := J \mathbb{E}^Q[f(Y_t)]$ and covariance $K_Q(t, s) :=$ $\sigma^2 \mathbb{E}^Q[f(Y_t)f(Y_s)].$

The main result of this section is the mean field dynamics:

Theorem 7. The law Q is that of the solutions of the following system

(19)
$$
\begin{cases} X_t & = X_0 + \int_0^t g(X_s) ds + J \int_0^t \mathbb{E}^Q[f(X_s)] ds + C_t \\ C_t & = \lambda W_t + \int_0^t \int_0^s \frac{1}{\lambda^2} \tilde{K}_Q^{s,\lambda}(s, u) dC_u ds \\ K_Q^{t,\lambda}(t, s) & = \sigma^2 \mathbb{E}^Q[f(X_t)f(X_s)] \\ \text{Law of } X & = Q, \quad Q_{|\mathcal{F}_0} = \mu_0 \end{cases}
$$

where W is a Q-Brownian motion and $\tilde{K}_Q^{t,\lambda}$ is given by (27).

Remark 8. The proof of Theorem 7 shows that the P-Brownian motion B is obtained by

(20)
$$
B_t = \frac{1}{\lambda} \left[\int_0^t m_Q(s) \, ds + C_t \right]
$$

Remark 9. It is sufficient to prove Theorem 7 in the particular case where $\lambda = 1$. Indeed, for any λ , we set $\bar{X}_t^i = \frac{1}{\lambda} X_t^i$ and the particle system \bar{X}^i evolves with the dynamics (6) with $\lambda = 1$, $\bar{f}(x) = \frac{1}{\lambda} f(\lambda x)$ and $\bar{g}(x) = \frac{1}{\lambda} g(\lambda x)$. Thus, in the following, we state the intermediate results for a general $\hat{\lambda}$ but for the sake of clarity we write the details only for $\lambda = 1$. We also simplify the notation $K^{t,\lambda}$ in K^t .

Before we can give the proof we need to take a short detour and discuss the interpretation of covariance functions such as K_Q as linear operators acting on functions in $L^2([0,t])$, $0 \le t \le T$ and the solutions to Volterra equations of the second kind.

4.1. Covariance functions and linear operators.

We consider the set of correlation functions K (symmetric positive kernels on $[0, T] \times [0, T]$, such that

(21)
$$
\int_0^T K(s,s) ds < \infty.
$$

Note that this is true for the correlation functions arising from any probability measure on $\mathcal C$ through the fourth equation in (14) because for the S-model where $f(y) = y$ the X_t s are bounded by $\pm A$ and for the H-model f is bounded, see (7).

Given such a correlation function K we associate to it the linear operator \bar{K}^t on $L^2([0,t])$ by

(22)
$$
(\bar{K}^t \cdot \psi)(s) = \int_0^t K(s, u)\psi(u) du \quad 0 \le s \le t, \forall \psi \in L^2([0, t]).
$$

In this context K is called the kernel of the operator \bar{K}^t .

Such an operator is called a Trace-Class operator (because of (21)) and the corresponding norm is noted

(23)
$$
\|\bar{K}^t\|_{t,1} = \int_0^t |K(s,s)| ds, \ 0 < t \leq T.
$$

It is known, e.g. [22], that the set of Trace-Class operators on $L^2([0,t])$ is a subset of the set of Hilbert-Schmidt operators equipped with the norm

(24)
$$
\|\bar{K}^t\|_{t,2} = \left(\int_0^t \int_0^t K(s,u)^2 ds du\right)^{1/2}, \ 0 < t \leq T
$$

and that the norms satisfy

(25)
$$
\|\bar{K}^t\|_{t,2} \le \|\bar{K}^t\|_{t,1}, \ 0 < t \le T.
$$

Given two such operators \bar{K}_1^t and \bar{K}_2^t their composition $\bar{K}_{12}^t = \bar{K}_1^t \circ \bar{K}_2^t$ is defined by

(26)
$$
K_{12}^{t}(s, u) = \int_{0}^{t} K_{1}^{t}(s, v) K_{2}^{t}(v, u) dv \quad \forall s, u \in [0, t]
$$

Given such a correlation operator \bar{K}^t we can define on $L^2([0,t])$ another operator, noted $\bar{\tilde{K}}^{t,\lambda}$ by

(27)
$$
\bar{\tilde{K}}^{t,\lambda} = \bar{K}^t \circ \left(\mathrm{Id} + \frac{1}{\lambda^2} \bar{K}^t \right)^{-1} = \left(\mathrm{Id} + \frac{1}{\lambda^2} \bar{K}^t \right)^{-1} \circ \bar{K}^t,
$$

 λ being a non zero real.

Remark 10. The reader may be puzzled by the presence of the upper index t in \bar{K}^t and $\bar{\tilde{K}}^{t,\lambda}$, This is meant to indicate that these operators act on functions of $L^2([0,t])$. This would be unnecessary if the operators acting on functions of $L^2([0, s])$, $0 < s < t$ were the restrictions of those acting on functions of $L^2([0, t])$. This is true for \overline{K} but it is *not* true in general for $\overline{\overline{K}}$.

Remark 11. Note that because K is a correlation function, \bar{K}^t has positive eigenvalues, hence Id + $\frac{1}{\lambda^2} \bar{K}^t$ is invertible, and its inverse $(\mathrm{Id} + \frac{1}{\lambda^2} \bar{K}^t)^{-1}$ is contracting.

The relation between \bar{K}^t and $\bar{\tilde{K}}^{t,\lambda}$ is given by the following Lemma (see Lemma A1 in [1]):

Lemma 12. Let $(G_t)_{0 \leq t \leq T}$ be a centered Gaussian process with correlation function K under the law ν . Under the law $\tilde{\nu}^t$ defined by

$$
d\tilde{\nu}^t(\theta) = \frac{\exp\{-\frac{1}{2\lambda^2} \int_0^t G_s^2(\theta) ds\}}{\int \exp\{-\frac{1}{2\lambda^2} \int_0^t G_s^2(\theta) ds\} d\nu} d\nu(\theta)
$$

 G_t is a centered Gaussian process with correlation function $\tilde{K}^{t,\lambda}$ defined by (27). In other words, \forall 0 \leq s, $u \leq t \leq T$

$$
\int_{\Theta} G_s(\theta) G_u(\theta) d\nu(\theta) = K(s, u) \text{ and } \int_{\Theta} G_s(\theta) G_u(\theta) d\tilde{\nu}^t(\theta) = \tilde{K}^{t, \lambda}(s, u).
$$

We collect some useful results about the norms of the operators \bar{K}_{Q}^{t} and $\bar{\tilde{K}}_{Q}^{t,\lambda}$ in the following lemma.

Lemma 13. We have

(28)
$$
\|\bar{K}_Q^t\|_{t,1}, \|\bar{K}_Q^{t,\lambda}\|_{t,1} \leq \begin{cases} \sigma^2 A^2 T & \text{S-model} \\ \sigma^2 C^2 T & \text{H-model} \end{cases} 0 \leq t \leq T,
$$

where A is the parameter in the S-model and C is the parameter in the H-model defined by (7).

Proof. According to Remark 9, we assume $\lambda = 1$. The first inequality in (28) is obtained from the definition of the Trace-Class norm:

$$
\|\bar{K}_Q^t\|_{t,1} = \sigma^2 \int_0^t \mathbb{E}^Q[f(X_s)^2] \, ds
$$

and the facts that $f = Id$ and $X_t^2 \leq A^2$ for the S-model, and $|f|$ is bounded by C for the H-model.

The second inequality in (28) is a direct consequence of Lemma A2 in [1]. \Box

4.2. A Volterra equation.

On the road to relating the processes B_t and W_t we have to spend some time on characterising the solutions to a Volterra equation of the second kind.

Our main result is the following Proposition.

Proposition 14. Consider a function $c \in L^2([0,T])$, a kernel $(\kappa(t,s))_{0 \leq s,t \leq T}$ such that $\kappa(t,s) = 0$ if $t < s$ and

(29)
$$
\forall t \in [0, T], \quad \int_0^t \kappa^2(t, s) \, ds \left(= \int_0^T \kappa^2(t, s) \, ds \right) \le M,
$$

and a stochastic process W.

The equation

(30)
$$
B_t = W_t + \int_0^t c(s) \, ds + \int_0^t \left(\int_0^s \kappa(s, u) \, dB_u \right) \, ds \, 0 \le t \le T,
$$

has a unique solution given by

(31)
$$
B_t = W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H(s, u) \widetilde{W}_u du \right) ds + \int_0^t c(s) ds + \int_0^t \left(\int_0^s H(s, u) c(u) du \right) ds
$$

where

(32)
$$
\widetilde{W}_t = \int_0^t \kappa(t, s) dW_s.
$$

The function H is defined from κ by

(33)
$$
\bar{H} = (\text{Id} - \bar{\kappa})^{-1} - \text{Id}.
$$

Moreover we have

(34)
$$
|H(s, u)| \leq Mh(MT) \quad 0 \leq s, u \leq t,
$$

where $h(x)$, $x \geq 0$, is the sum of the absolutely convergent series

$$
h(x) = \sum_{n \ge 0} \frac{x^n}{\sqrt{n!}}
$$

Proof. Necessary condition:

Consider a solution (B_t) to (30), one denotes $\tilde{B}_t = \int_0^t \kappa(t, s) dB_s$. Thus, (30) can be written

(35)
$$
B_t = W_t + \int_0^t c(s) \, ds + \int_0^t \tilde{B}_s \, ds.
$$

Differentiating both sides of (35), multiplying by $\kappa(t, s)$, and integrating between 0 and t yields

(36)
$$
\tilde{B}_t = \int_0^t \kappa(t, s) dW_s + \int_0^t \kappa(t, s) c(s) ds + \int_0^t \kappa(t, s) \tilde{B}_s ds
$$

$$
=:\widetilde{W}_t + \tilde{c}(t) + \int_0^t \kappa(t, s) \tilde{B}_s ds,
$$

Now (36) is a classical linear Volterra equation of the second kind in the unknown \tilde{B}_t , except for the fact that \widetilde{W}_t is a random process instead of a deterministic function in $L^2([0,T])$. A close look at the proof in this case (which is found, e.g.

in [26, pp.10-15]) shows that it carries over to our case and that the unique solution to (36) can be expressed as

(37)
$$
\tilde{B}_t = \widetilde{W}_t + \tilde{c}(t) + \int_0^t H(t,s) \left(\widetilde{W}_s + \tilde{c}(s)\right) ds,
$$

where the resolvent kernel H is the sum of the absolutely convergent series

(38)
$$
H(t,s) = \sum_{n\geq 0} (\kappa(t,s))^{(n+1)}.
$$

The function $\kappa(t,s)^{(n)}$ is the kernel of the operator $\bar{\kappa}$ iterated n times. We notice that

$$
\bar{H} = (\mathrm{Id} - \bar{\kappa})^{-1} - \mathrm{Id}
$$

Remark 15. Note that under the assumption on κ in Proposition 14, one can simplify the composition defined by (26) into

$$
\kappa^{(2)}(s, u) = \int_u^s \kappa(s, v)\kappa(v, u) dv \quad \forall u \le s \in [0, t].
$$

In other words, the composition does not depend on t for such kernels.

Combining (35) and (37) we have obtained the necessary condition

$$
B_t = W_t + \int_0^t \widetilde{W}_s ds + \int_0^t c(s) ds + \int_0^t \left(\int_0^s H(s, u) (\widetilde{W}_u + \widetilde{c}(u)) du \right) ds.
$$

Now it is known from the theory of Volterra equations that the resolvent kernel H_Q and the kernel κ satisfy the relation

$$
H(s, v) = \kappa(s, v) + \int_v^s \kappa(s, u) H(u, v) du
$$

= $\kappa(s, v) + \int_v^s H(s, u) \kappa(u, v) du$,

or in operator form

(39)
$$
\bar{H} = \bar{\kappa} + \bar{\kappa} \circ \bar{H} = \bar{\kappa} + \bar{H} \circ \bar{\kappa}.
$$

This implies that, according to (36),

$$
\int_0^t \tilde{c}(s) ds + \int_0^t \left(\int_0^s H(s, u) \tilde{c}(u) du \right) ds = \int_0^t \left(\int_0^s H(s, u) c(u) du \right) ds
$$

and therefore

$$
B_t = W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H(s, u) \widetilde{W}_u du \right) ds + \int_0^t c(s) ds + \int_0^t \left(\int_0^s H(s, u) c(u) du \right) ds.
$$

The last identity and (32) prove that the process B_t is adapted to \mathbb{F}^W . Sufficient condition:

We show that, conversely, if B_t is given by (31) it satisfies (30). Indeed, (31) commands that

$$
dB_u = dW_u + (\widetilde{W}_u + c(u))du + \left(\int_0^u H(u, v)(\widetilde{W}_v + c(v)) dv\right) du.
$$

Multiplying both sides of this equation by $\kappa(s, u)$ and integrating w.r.t. u between 0 and s , we obtain, using (36) and (32)

(40)
$$
\int_0^s \kappa(s, u) dB_u = \widetilde{W}_s + \int_0^s \kappa(s, u) (\widetilde{W}_u + c(u)) du + \int_0^s \kappa(s, u) \left(\int_0^u H(u, v) (\widetilde{W}_v + c(v)) dv \right) du,
$$

or, in operator form,

$$
\int_0^s \kappa(s, u) dB_u = \widetilde{W}_s + (\bar{\kappa} + \bar{\kappa} \circ \bar{H}) \cdot (\widetilde{W} + c)(s)
$$

$$
= \widetilde{W}_s + \bar{H} \cdot (\widetilde{W} + c)(s),
$$

by (39). We have obtained

$$
\int_0^s \kappa(s, u) dB_u = \widetilde{W}_s + \int_0^s H(s, u) \widetilde{W}_u du + \int_0^s H(s, u) c(u) du.
$$

Integrating both sides of this equation between 0 and t and adding $W_t + \int_0^t c(s) ds$ to both sides yields

$$
W_t + \int_0^t c(s) ds + \int_0^t \left(\int_0^s \kappa(s, u) dB_u \right) ds = W_t + \int_0^t \widetilde{W}_s ds
$$

+
$$
\int_0^t \left(\int_0^s H(s, u) \widetilde{W}_u du \right) ds + \int_0^t c(s) ds + \int_0^t \left(\int_0^s H(s, u) c(u) du \right) ds,
$$

and since the right hand side of this equation is equal to B_t by (31) this indicates that (30) is satisfied, as claimed.

To finish the proof we must show (34). It is an easy consequence of the proof found in [26, pp. 10-15], of (25) and assumption (29). \Box

4.3. Proof of Theorem 7.

We now prove Theorem 7. Using Remark 9 we write the proof in the case $\lambda = 1$.

Proof. We start from (18) which we reproduce here in the case $\lambda = 1$. Note that we have dropped for simplicity the upper index Q.

(41)
$$
\frac{dQ}{dP} = \int \exp\left\{ \int_0^T G_t \, dB_t - \int_0^T G_t^2 \, dt \right\} d\gamma_Q
$$

The plan is to rewrite dQ/dP in such a way that we can apply Girsanov's Theorem, Theorem 5, and read off the dynamics from the new expression.

We write

$$
G_t = m_Q(t) + R_t,
$$

where R_t is, under γ_Q , a zero mean Gaussian process with correlation function $K_Q(t, s)$ and (see Lemma 12), under $\tilde{\gamma}_Q^t$, a zero mean Gaussian process with correlation function $\tilde{K}_{Q}^{t}(t, s)$, where $\tilde{\gamma}_{Q}^{t}$ is defined by

(42)
$$
d\tilde{\gamma}_Q^t = \frac{\exp\left\{-\int_0^t R_s^2 ds\right\}}{\int \exp\left\{-\int_0^t R_s^2 ds\right\} d\gamma_Q} d\gamma_Q := \Lambda_t d\gamma_Q
$$

We write

(43)
$$
\frac{dQ}{dP} = \exp\left\{ \int_0^T m_Q(t) \, dB_t - \int_0^T m_Q^2(t) \, dt \right\} \times \int \exp\left\{ \int_0^T R_t \, dB_t - \int_0^T R_t m_Q(t) \, dt - \frac{1}{2} \int_0^T R_t^2 \, dt \right\} d\gamma_Q.
$$

The first factor in the right hand side does not pose any problem. The second does because of the presence of R_t . The following parts of the proof are meant to get rid of R_t .

We rewrite the second factor of the right hand side as follows, using (42) ,

$$
\int \exp\left\{\int_0^T R_t dB_t - \int_0^T R_t m_Q(t) dt - \frac{1}{2} \int_0^T R_t^2 dt \right\} d\gamma_Q =
$$
\n
$$
\int \exp\left\{-\frac{1}{2} \int_0^T R_t^2 dt \right\} d\gamma_Q \times \int \exp\left\{\int_0^T R_t dB_t - \int_0^T R_t m_Q(t) dt \right\} d\tilde{\gamma}_Q^T
$$

Dealing with the term $\int \exp \left\{-\frac{1}{2}\right\}$ 2 $\int_0^T R_t^2 dt \bigg\} d\gamma_Q$: We define

$$
h(t) = \log \int \exp \left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} d\gamma_Q.
$$

We have

$$
h'(t) = -\frac{1}{2} \int R_t^2 \frac{\exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} d\gamma_Q}{\int \exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} d\gamma_Q},
$$

and by (42) and the definition of \tilde{K}_{Q}^{t}

$$
h'(t) = -\frac{1}{2} \int R_t^2 d\tilde{\gamma}_Q = -\frac{1}{2} \tilde{K}_Q^t(t, t)
$$

Since $h(0) = 0$ we conclude that

$$
h(t) = -\frac{1}{2} \int_0^t \tilde{K}_Q^s(s, s) ds
$$

and hence that

(44)
$$
\int \exp\left\{-\frac{1}{2}\int_0^T R_t^2 dt\right\} d\gamma_Q = \exp\left\{-\frac{1}{2}\int_0^T \tilde{K}_Q^t(t,t) dt\right\}
$$

 $\textbf{Dealing with the term } \int \exp \left\{ \int_0^T R_t \, dB_t - \int_0^T R_t m_Q(t) \, dt \right\} \, d\tilde{\gamma}_Q^T;$

Conditionally to B_t , $\int_0^T R_t dB_t$ is, under $\tilde{\gamma}_{Q}^T$, a centered Gaussian process with variance $\int_0^T \int_0^T \tilde{K}_Q^T(t,s) dB_s dB_t$. Similarly $\int_0^T R_t m_Q(t) dt$ is, under $\tilde{\gamma}_Q^T$, a centered Gaussian process with variance $\int_0^T \int_0^T \tilde{K}_Q^T(t,s) m_Q(t) m_Q(s) ds dt$. Their sum is also, conditionally to B_t , a centered Gaussian process and therefore, by Gaussian calculus

(45)
$$
\int \exp \left\{ \int_0^T R_t \, dB_t - \int_0^T R_t m_Q(t) \, dt \right\} d\tilde{\gamma}_Q^T = \\ \exp \left\{ \frac{1}{2} \Big(\int_0^T \int_0^T \tilde{K}_Q^T(t, s) \, dB_s \, dB_t - 2 \mathbb{E}_{\tilde{\gamma}_Q^T} \Big[\int_0^T R_t \, dB_t \times \int_0^T R_t m_Q(t) \, dt \Big] + \int_0^T \int_0^T \tilde{K}_Q^T(t, s) m_Q(t) m_Q(s) \, ds \, dt \Big) \right\} = \\ \exp \left\{ \frac{1}{2} \int_0^T \int_0^T \tilde{K}_Q^T(t, s) \, dB_s \, dB_t - \int_0^T \int_0^T \tilde{K}_Q^T(t, s) m_Q(s) \, ds \, dB_t + \frac{1}{2} \int_0^T \int_0^T \tilde{K}_Q^T(t, s) m_Q(t) m_Q(s) \, ds \, dt \right\}.
$$

Combining (43)-(45) we end up with the following formula for dQ/dP :

(46)
\n
$$
\frac{dQ}{dP} = \exp\left\{\int_0^T m_Q(t) \, dB_t - \frac{1}{2} \int_0^T m_Q(t)^2 \, dt\right\} \times \exp\left\{-\frac{1}{2} \int_0^T \tilde{K}_Q^t(t, t) \, dt\right\} \times \exp\left\{\frac{1}{2} \int_0^T \int_0^T \tilde{K}_Q^T(t, s) \, dB_s \, dB_t - \int_0^T \int_0^T \tilde{K}_Q^T(t, s) m_Q(s) \, ds \, dB_t + \frac{1}{2} \int_0^T \int_0^T \tilde{K}_Q^T(t, s) m_Q(t) m_Q(s) \, ds \, dt\right\}
$$

This formula does not allow us yet to use Girsanov Theorem, Theorem 5. Therefore we need to massage it in order to make this possible.

We apply Itô formula three times, one for each of the last three terms of the product in the right hand side of (46). We first note that, according to (42),

(47)
$$
\frac{d\Lambda_t}{dt} = \frac{1}{2}\Lambda_t \left(\tilde{K}_Q^t(t,t) - R_t^2 \right).
$$

Indeed,

$$
\frac{d\Lambda_t}{dt} = -\frac{1}{2} \left[R_t^2 \Lambda_t - \exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} \frac{\int R_t^2 \exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} d\gamma_Q}{\left(\int \exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} d\gamma_Q\right)^2} \right]
$$

\n
$$
= \frac{1}{2} \Lambda_t \left[\int R_t^2 \frac{\exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\}}{\int \exp\left\{-\frac{1}{2} \int_0^t R_s^2 ds\right\} d\gamma_Q} d\gamma_Q - R_t^2 \right]
$$

\n
$$
= \frac{1}{2} \Lambda_t \left(\int R_t^2 d\tilde{\gamma}_Q - R_t^2 \right)
$$

\n
$$
= \frac{1}{2} \Lambda_t \left(\tilde{K}_Q^t(t, t) - R_t^2 \right).
$$

Second we note that, by definition,

(48)
$$
\tilde{K}_Q^T(t,s) = \int R_t R_s \Lambda_T d\gamma_Q.
$$

Dealing with the term $\exp(\frac{1}{2}\int_0^T\int_0^T\tilde{K}_Q^T(t,s)\,dB_s\,dB_t)$: We prove that

(49)
$$
\exp\left\{\frac{1}{2}\int_0^T \int_0^T \tilde{K}_Q^T(t,s) dB_s dB_t\right\} = \exp\left\{\frac{1}{2}\int_0^T \tilde{K}_Q^t(t,t) dt\right\} \times \exp\left\{\int_0^T \int_0^t \tilde{K}_Q^t(t,s) dB_s dB_t\right\} \times \exp\left\{-\frac{1}{2}\int_0^T \left(\int_0^t \tilde{K}_Q^t(t,s) dB_s\right)^2 dt\right\}.
$$

Indeed, it follows from (48) that

(50)
$$
\int_0^T \int_0^T \tilde{K}_Q^T(t,s) dB_s dB_t = \int \left(\int_0^T R_t dB_t \right)^2 \Lambda_T d\gamma_Q
$$

$$
= \int \int_0^T d \left(\left(\int_0^t R_s dB_s \right)^2 \Lambda_t \right) d\gamma_Q
$$

Because Λ_t is deterministic, Itô product rule indicates that

$$
d\left(\left(\int_0^t R_s dB_s\right)^2 \Lambda_t\right) = \left(\int_0^t R_s dB_s\right)^2 d\Lambda_t + \Lambda_t d\left(\int_0^t R_s dB_s\right)^2
$$

There remains to write Itô's formula for $\left(\int_0^t R_s dB_s\right)^2$

(51)
$$
\left(\int_0^t R_s dB_s\right)^2 = 2 \int_0^t R_s \left(\int_0^s R_u dB_u\right) dB_s + \int_0^t R_s^2 ds,
$$

to remember Isserli's theorem

(52)
$$
\int R_t^2 R_s R_u d\tilde{\gamma}_Q^t = \tilde{K}_Q^t(t, t) \tilde{K}_Q^t(s, u) + 2 \tilde{K}_Q^t(t, s) \tilde{K}_Q^t(t, u) \quad s, u \le t
$$

and to combine (48) , (47) , (50) , (51) and (52) to obtain

$$
\int_0^T \int_0^T \tilde{K}_Q^T(t,s) dB_s dB_t = 2 \int_0^T \int_0^t \tilde{K}_Q^t(t,s) dB_s dB_t + \int_0^T \tilde{K}_Q^t(t,t) dt - \int_0^T \left(\int_0^t \tilde{K}_Q^t(t,s) dB_s \right)^2 dt.
$$

Dealing with the term $\exp\left\{\frac{1}{2}\right\}$ $\frac{1}{2} \int_0^T \int_0^T \tilde{K}^T_Q(t,s) m_Q(s) m_Q(t)\, ds\, dt\Big\}$: Similar arguments show that

(53)
$$
\exp\left\{\frac{1}{2}\int_0^T \int_0^T \tilde{K}_Q^T(t,s)m_Q(s)m_Q(t) ds dt\right\} = \exp\left\{\int_0^T m_Q(t) \left(\int_0^t \tilde{K}_Q^t(t,s)m_Q(s) ds\right) dt\right\} \times \exp\left\{-\frac{1}{2}\int_0^T \left(\int_0^t \tilde{K}_Q^t(t,s)m_Q(s) ds\right)^2 dt\right\}.
$$

Indeed, by (48), we have

$$
\int_0^T \int_0^T \tilde{K}_Q^T(t,s) m_Q(s) m_Q(t) ds dt = \int \left(\int_0^T \int_0^T R_t R_s m_Q(t) m_Q(s) ds dt \right) \Lambda_T d\gamma_Q
$$

=
$$
\int \left(\int_0^T R_t m_Q(t) dt \right)^2 \Lambda_T d\gamma_Q
$$

We apply Itô's formula to $\left(\int_0^T R_t m_Q(t) dt\right)^2 \Lambda_T$. By (47) we have:

$$
\left(\int_0^T R_t m_Q(t) dt\right)^2 \Lambda_T = 2 \int_0^T R_t m_Q(t) \left(\int_0^t R_s m_Q(s) ds\right) \Lambda_t dt
$$

+
$$
\frac{1}{2} \int_0^T \left(\int_0^t R_s m_Q(s) ds\right)^2 \Lambda_t \left(\tilde{K}_Q^t(t, t) - R_t^2\right) dt.
$$

Integrating both sides of the previous equation w.r.t. to γ_Q and making use of (48) yields

$$
\int \left(\int_0^T R_t m_Q(t) dt\right)^2 \Lambda_T d\gamma_Q = 2 \int_0^T m_Q(t) \left(\int_0^t \tilde{K}_Q^t(t, s) m_Q(s) ds\right) dt \n+ \frac{1}{2} \int_0^T \tilde{K}_Q^t(t, t) \left(\int_0^t \int_0^t \tilde{K}_Q^t(s, u) m_Q(s) m_Q(u) ds du\right) dt \n- \frac{1}{2} \int_0^T \left(R_t^2 \left(\int_0^t R_s m_Q(s) ds\right)^2 \Lambda_t\right) d\gamma_Q dt.
$$

Using (52) again we obtain

$$
\int \left(\int_0^T R_t m_Q(t) dt\right)^2 \Lambda_T d\gamma_Q = 2 \int_0^T m_Q(t) \left(\int_0^t \tilde{K}_Q^t(t,s) m_Q(s) ds\right) dt
$$

$$
- \frac{1}{2} \int_0^T \left(\int_0^t \tilde{K}_Q^t(t,s) m_Q(s) ds\right)^2 dt,
$$

i.e. (53).

 $\textbf{Dealing with the term } \exp\left\{-\int_0^T\int_0^T\tilde{K}_Q^T(t,s)m_Q(s)\,ds\,dB_t\right\}\!;$ Similar arguments show that

(54)
$$
\exp\left\{-\int_0^T \int_0^T \tilde{K}_Q^T(t,s)m_Q(s) ds dB_t\right\}
$$

\n
$$
= \exp\left\{-\int_0^T \left(\int_0^t \tilde{K}_Q^t(t,s)m_Q(s) ds\right) dB_t\right\} \times
$$

\n
$$
\exp\left\{-\int_0^T m_Q(t) \left(\int_0^t \tilde{K}_Q^t(t,s) dB_s\right) dt\right\} \times
$$

\n
$$
\exp\left\{\int_0^T \left(\int_0^t \tilde{K}_Q^t(t,s) dB_s\right) \times \left(\int_0^t \tilde{K}_Q^t(t,s)m_Q(s) ds\right) dt\right\}
$$

Indeed, by (48)

$$
\int_0^T \int_0^T \tilde{K}_Q^T(t,s) m_Q(s) ds dB_t = \int \left(\int_0^T \int_0^T R_t R_s m_Q(s) ds dB_t \right) \Lambda_T d\gamma_Q
$$

=
$$
\int \left(\int_0^T R_t dB_t \right) \left(\int_0^T R_s m_Q(s) ds \right) \Lambda_T d\gamma_Q
$$

We apply Itô's formula to the product $\left(\int_0^T R_t dB_t\right) \left(\int_0^T R_s m_Q(s) ds\right) \Lambda_T$ using (47):

$$
\left(\int_0^T R_t dB_t\right) \left(\int_0^T R_s m_Q(s) ds\right) \Lambda_T = \int_0^T \left(\int_0^t R_s m_Q(s) ds\right) R_t \Lambda_t dB_t
$$

+
$$
\int_0^T \left(\int_0^t R_s dB_s\right) R_t m_Q(t) \Lambda_t dt
$$

+
$$
\frac{1}{2} \int_0^T \left(\int_0^t R_s dB_s\right) \left(\int_0^t R_u m_Q(u) du\right) \left(\tilde{K}_Q^t(t,t) - R_t^2\right) \Lambda_t dt
$$

We integrate both sides of the previous equation w.r.t. γ_Q and use (48) and (52) to obtain

$$
\int \left(\int_0^T R_t dB_t \right) \left(\int_0^T R_s m_Q(s) ds \right) \Lambda_T d\gamma_Q = \int_0^T \left(\int_0^t \tilde{K}_Q(t, s) m_Q(s) ds \right) dB_t + \int_0^T m_Q(t) \left(\int_0^t \tilde{K}_Q(t, s) dB_s \right) dt - \int_0^T \left(\int_0^t \tilde{K}_Q(t, s) dB_s \right) \left(\int_0^t \tilde{K}_Q^t(t, s) m_Q(s) ds \right) dt
$$

Collecting (46) , (49) , (53) and (54) , and defining

$$
\tilde{B}_t = \int_0^t \tilde{K}_Q^t(t, s) dB_s \qquad \tilde{m}_Q(t) = \int_0^t \tilde{K}_Q^t(t, s) m_Q(s) ds,
$$

we obtain

(55)
\n
$$
\frac{dQ}{dP} = \exp\left\{ \int_0^T (m_Q(t) - \tilde{m}_Q(t) + \tilde{B}_t) \, dB_t - \frac{1}{2} \int_0^T (m_Q(t) - \tilde{m}_Q(t) + \tilde{B}_t)^2 \, dt \right\}
$$

We are almost done. We next define

(56)
$$
c_Q(t) := m_Q(t) - \tilde{m}_Q(t) = \left(\text{Id} - \tilde{K}_Q^t\right) \cdot m_Q(t)
$$

By Girsanov's Theorem, Theorem 5, we obtain that

$$
W_t = B_t - \int_0^t (c_Q(s) + \tilde{B}_s) ds
$$

is a Brownian motion under Q. Therefore we have

$$
B_t = W_t + \int_0^t c_Q(s) \, ds + \int_0^t \tilde{B}_s \, ds.
$$

Using the definition of \tilde{B} , we obtain the Volterra equation

$$
B_t = W_t + \int_0^t c_Q(s) \, ds + \int_0^t \int_0^s \tilde{K}^s(s, u) \, dB_u \, ds.
$$

and we apply Proposition 14 with $\kappa(t,s) = \tilde{K}^s(s, u) \mathbb{1}_{u \leq s}$ to solve it.

(57)
$$
B_t = W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds
$$

 $+ \int_0^t c_Q(s) ds + \int_0^t \left(\int_0^s H_Q(s, u) c_Q(u) du \right) ds.$

We now use (56) and (39) and obtain

$$
B_t = W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds
$$

+
$$
\int_0^t m_Q(s) ds + \int_0^t \left(\int_0^s H_Q(s, u) m_Q(u) du \right) ds
$$

-
$$
\int_0^t \int_0^s \widetilde{K}^s(s, u) m_Q(u) du ds
$$

-
$$
\int_0^t \left(\int_0^s H_Q(s, u) \int_0^u \widetilde{K}^u(u, v) m_Q(v) dv du \right) ds.
$$

and hence

(58)
$$
B_t = W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds + \int_0^t m_Q(s) ds.
$$

We apply Proposition 14 with $c \equiv 0$ and observe that the process $W_t + \int_0^t \widetilde{W}_s ds +$ $\int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds$ is the solution C_t to the Volterra equation

$$
C_t = W_t + \int_0^t \int_0^s \tilde{K}_Q^s(s, u) dC_u ds.
$$

By Girsanov's Theorem 5, and the fact $m_Q(s) = J \mathbb{E}^Q[f(X_s)]$, this yields the equations (19) of the dynamics. \Box

Remark 16. Note that as a side result we have proved (20) in Remark 8. Indeed, for a general λ , we recall the Volterra type equation

$$
C_t = \lambda W_t + \int_0^t \int_0^s \frac{1}{\lambda^2} \tilde{K}_Q^{s,\lambda}(s, u) dC_u ds.
$$

The process $\frac{C_t}{\lambda}$ solves a Volterra equation of type (30) with $c \equiv 0$ and $\kappa(t, s) =$ $\frac{1}{\lambda^2} \tilde{K}^{t,\lambda}(t,s) \mathbb{1}_{0 \leq s \leq t}.$

Remark 17. This result is a precise description of the somewhat mysterious "effective interaction term" used in the papers $[27]$ and $[4]²$. Indeed, the equation (58)

$$
B_t = \frac{1}{\lambda} \left(\lambda W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds + \int_0^t m_Q(s) ds \right),
$$

allows us to rewrite the meanfield dynamics (19) as

$$
\begin{cases}\nX_t &= X_0 + \int_0^t g(X_s) ds + J \int_0^t \mathbb{E}^Q[f(X_s)] ds + \lambda \int_0^t \widetilde{W}_s ds \\
&+ \lambda \int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds + \lambda W_t \\
\widetilde{W}_t &= \frac{1}{\lambda^2} \int_0^t \widetilde{K}_Q^{t, \lambda}(t, s) dW_s \\
H_Q(t, s) &= \sum_{n \ge 0} \left(\frac{1}{\lambda^2} \widetilde{K}_Q^{t, \lambda}(t, s) \mathbb{1}_{s \le t} \right)^{(n+1)} \\
K_Q^{t, \lambda}(t, s) &= \sigma^2 \mathbb{E}^Q[f(X_t)f(X_s)] \\
\text{Law of } X &= Q, \quad Q_{|\mathcal{F}_0} = \mu_0\n\end{cases}
$$

where W is a Q-Brownian motion and $\tilde{K}_Q^{t,\lambda}$ is given by (27) as a function of $K_Q^{t,\lambda}$. The dependency of the effective interaction term $J\mathbb{E}^Q[f(X_t)] + \lambda \widetilde{W}_t +$ $\lambda \int_0^t H_Q(t, s) \widetilde{W}_s ds$ upon the Brownian W and the law Q is clear. It is a Gaussian process with mean $J\mathbb{E}^Q[f(X_t)]$. Its covariance can be computed using standard Itô calculus as a function of $\tilde{K}_Q^{\tilde{t},\lambda}$.

²This term was derived by an incorrect application of Girsanov's Theorem, Theorem 5, in [4]

5. THE SPECIAL CASE $f = 1$

We now study the special case where the function f in (6) is constant and equal for example to 1. This means that each neuron j contributes exactly $J_{i \leftarrow j}$ to the activity of neuron *i*, independently of the value X_t^j of its membrane potential. It can be seen as a special case of the H-model when the sigmoidal function f is flat. It may appear as a toy example but it is useful as it shows the abstract concepts of the previous sections at work. We note Q_1 the corresponding limit law of the network.

5.1. Annealed mean-field dynamics.

The third equation in (19) reads in this case

$$
K_{Q_1}^t(u,s) = \sigma^2, \quad \forall u, \ s \le t \le T
$$

We prove the following Lemma:

Lemma 18. We have:

(59)
$$
\tilde{K}_{Q_1}^{t,\lambda}(u,s) = \frac{\sigma^2 \lambda^2}{\lambda^2 + \sigma^2 t}, \quad \forall u, s \le t \le T
$$

Proof. Given $\varphi \in L^2([0,T])$ we have by (22)

(60)
$$
\bar{K}_{Q_1}^t \cdot \varphi(s) = \sigma^2 \int_0^t \varphi(u) \, du
$$

Next we identify the kernel of $(\mathrm{Id} + \frac{1}{\lambda^2} \bar{K}_{Q_1}^t)^{-1}$. Given $\varphi \in L^2([0,T])$, denote by ψ its image by $(\mathrm{Id} + \frac{1}{\lambda^2} \bar{K}_{Q_1}^t)^{-1}$. By (60) we have

$$
\varphi(s) = (\mathrm{Id} + \frac{1}{\lambda^2} \bar{K}_{Q_1}^t) \cdot \psi(s) = \psi(s) + \frac{\sigma^2}{\lambda^2} \int_0^t \psi(u) \, du \, \forall s \le t.
$$

Integrating both sides between 0 and t yields

(61)
$$
\int_0^t \varphi(s) \, ds = \int_0^t \psi(s) \, ds + \frac{\sigma^2}{\lambda^2} t \int_0^t \psi(u) \, du = \left(1 + \frac{\sigma^2}{\lambda^2} t\right) \int_0^t \psi(s) \, ds.
$$

We conclude that for all $\varphi \in L^2([0,T])$, by (60) and (61)

$$
\left(\bar{K}_{Q_1}^t \circ (\text{Id} + \frac{1}{\lambda^2} \bar{K}_{Q_1}^t)^{-1}\right) \cdot \varphi(t) = \bar{K}_{Q_1}^t \cdot \psi(t) = \sigma^2 \int_0^t \psi(s) \, ds = \sigma^2 \frac{1}{1 + \frac{\sigma^2}{\lambda^2} t} \int_0^t \varphi(s) \, ds,
$$

which proves the Lemma. \Box

Remark 19. This is an illustration of the fact, mentioned in Remark 10, that $\bar{\tilde{K}}^{s,\lambda}$ is not the restriction of $\bar{\tilde{K}}^{t,\lambda}, s < t \leq T$, to $L^2([0, s])$. Indeed, the kernel $\tilde{K}^{t,\lambda}(u, v)$ is constant and equal to $\frac{\sigma^2 \lambda^2}{\lambda^2 + \sigma^2}$ $\frac{\sigma^2 \lambda^2}{\lambda^2 + \sigma^2 t}$ on $[0, s] \times [0, s]$. while $\tilde{K}^{s,\lambda}(u, v)$ is also constant but equal to $\frac{\sigma^2 \lambda^2}{\lambda^2 + \sigma^2}$ $\overline{\lambda^2+\sigma^2s}$.

Lemma 20. We have:

(62)
$$
H_{Q_1}(t,s) = \frac{\sigma^2}{\lambda^2 + \sigma^2 s} \mathbb{1}_{s \leq t}
$$

Proof. By definition

$$
H_{Q_1}(t,s) = \sum_{n\geq 0} \left(\frac{1}{\lambda^2} \tilde{K}^{t,\lambda}_{Q_1}(t,s) \mathbb{1}_{s\leq t}\right)^{(n+1)}
$$

We obtain successively

$$
\left(\frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(t,s)\mathbb{1}_{s\leq t}\right)^{(1)} = \frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(t,s)\mathbb{1}_{s\leq t} = \frac{\sigma^2}{\lambda^2 + \sigma^2 t}\mathbb{1}_{s\leq t}
$$
\n
$$
\left(\frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(t,s)\mathbb{1}_{s\leq t}\right)^{(2)} = \frac{1}{\lambda^4}\int_s^t \tilde{K}_{Q_1}^{t,\lambda}(t,u)\tilde{K}_{Q_1}^{u,\lambda}(u,s) du = \frac{\sigma^2}{\lambda^2 + \sigma^2 t}\log\frac{\lambda^2 + \sigma^2 t}{\lambda^2 + \sigma^2 s}\mathbb{1}_{s\leq t}
$$
\n
$$
\left(\frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(t,s)\mathbb{1}_{s\leq t}\right)^{(3)} = \int_s^t \left(\frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(t,u)\right)^{(1)} \left(\frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(u,s)\right)^{(2)} du = \frac{\sigma^2}{\lambda^2 + \sigma^2 t}\frac{1}{2}\left(\log\frac{\lambda^2 + \sigma^2 t}{\lambda^2 + \sigma^2 s}\right)^2 \mathbb{1}_{s\leq t}
$$
\n
$$
\vdots
$$

$$
\left(\frac{1}{\lambda^2}\tilde{K}_{Q_1}^{t,\lambda}(t,s)\mathbb{1}_{s\leq t}\right)^{(n)} = \frac{\sigma^2}{\lambda^2 + \sigma^2 t} \frac{1}{(n-1)!} \left(\log \frac{\lambda^2 + \sigma^2 t}{\lambda^2 + \sigma^2 s}\right)^{(n-1)} \mathbb{1}_{s\leq t},
$$

and therefore, by (38) , we obtain the Lemma. \Box

Remark 21. One finally verifies (39) in the case $\kappa(t,s) = \frac{1}{\lambda^2} \tilde{K}^{t,\lambda}_{Q_1}(t,s) \mathbb{1}_{s \leq t}$:

$$
\frac{1}{\lambda^2} \tilde{K}_{Q_1}^{t,\lambda}(t,s) \mathbb{1}_{s \le t} - H_{Q_1}(t,s) = \frac{\sigma^2}{\lambda^2 + \sigma^2 t} - \frac{\sigma^2}{\lambda^2 + \sigma^2 s} \n= -\int_s^t \frac{1}{\lambda^2} \tilde{K}_{Q_1}^{t,\lambda}(t,u) H_{Q_1}(u,s) du,
$$

i.e.

$$
\frac{1}{\lambda^2} \bar{\tilde{K}}_{Q_1}^{t,\lambda} + \frac{1}{\lambda^2} \bar{\tilde{K}}_{Q_1}^{t,\lambda} \circ \bar{H}_{Q_1} = \bar{H}_{Q_1}.
$$

Let us now write the equations (19) of the dynamics is this case. We clearly have

$$
m_Q(t)=J,
$$

and by the second equation in (19)

$$
C_t = \lambda W_t + \sigma^2 \int_0^t \frac{C_s}{\lambda^2 + \sigma^2 s} ds
$$

We can then compute C_t as a function of the Q_1 Brownian motion W_t by solving the previous linear SDE obtaining

(63)
$$
C_t = \lambda \left(\lambda^2 + \sigma^2 t\right) \int_0^t \frac{dW_s}{\lambda^2 + \sigma^2 s}
$$

and write the corresponding dynamics as

(64)
$$
X_t = X_0 + \int_0^t g(X_s) ds + Jt + \lambda \left(\lambda^2 + \sigma^2 t\right) \int_0^t \frac{dW_s}{\lambda^2 + \sigma^2 s}
$$

or equivalently in differential form

(65)
$$
dX_t = \left(g(X_t) + J + \lambda \sigma^2 \int_0^t \frac{dW_s}{\lambda^2 + \sigma^2 s}\right) dt + \lambda dW_t
$$

Remark 22. The "effective interaction term" of Remark 17, $J + \lambda \sigma^2 \int_0^t$ dW_s $\frac{dW_s}{\lambda^2+\sigma^2s}$, has mean J and variance $\frac{t\sigma^4}{\lambda^2+\sigma^2t}$ at time t as can be easily verified. Note that this is different from the variance reported in previous work such as $[11]$ [Equations (25)] or [4][Section 3.3]. It shows the advantage of the approach developed in this article of working out the exact dependency of the effective interaction term with the Brownian.

5.2. Quenched mean-field dynamics.

The first equation in (6) writes in this case

(66)
$$
dX_t^i = \left(-X_t^i + S_i\right) dt + \lambda dB_t^i \quad 1 \le i \le N,
$$

where

$$
S_i = \sum_{j=1}^{N} J_{i \leftarrow j}.
$$

Since the $J_{i \leftarrow j}$ s are i.i.d. as $\mathcal{N}(J/N, \sigma^2/N)$, the S_i s are i.i.d. as $\mathcal{N}(J, \sigma^2)$. The propagation of chaos is obvious in this case since the N equations (66) are independent and the quenched mean-field dynamics is also obvious.

(67)
$$
dX_t = (-X_t + G) dt + \lambda dB_t,
$$

$$
G \simeq \mathcal{N}(J, \sigma^2), \text{ independent of } B
$$

5.3. Comparison between the quenched and annealed dynamics.

We know from the results in [1, 14] that these two dynamics should be identical. It is unclear though that the solutions to (65) and (67) have the same laws. In effect they do, as shown in the next Proposition.

Proposition 23. The solutions to (65) and (67) are two Gaussian processes with identical means and covariance functions, assuming identical Gaussian distributed initial conditions.

Proof. Gaussianity is clear. We show that the two processes have the same mean and variance. The solution to the quenched mean-field dynamics (67) is readily computed as

(68)
$$
X_t = X_0 e^{-t} + G(1 - e^{-t}) + e^{-t} \lambda \int_0^t e^s dW_s
$$

where $G \simeq \mathcal{N}(J, \sigma^2)$, X_0 et W are assumed to be independent. Hence, taking the expected value

(69)
$$
\mathbb{E}(X_t) = J + (\mathbb{E}(X_0) - J)e^{-t},
$$

and

$$
Var(X_t) = Var(G)(1 - e^{-t})^2 + Var(X_0)e^{-2t} + \lambda^2 \frac{1 - e^{-2t}}{2},
$$

and therefore

(70)
$$
\text{Var}(X_t) = \text{Var}(G) + \frac{\lambda^2}{2} - 2\text{Var}(G)e^{-t} + \left(\text{Var}(G) + \text{Var}(X_0) - \frac{\lambda^2}{2}\right)e^{-2t}
$$

Assuming $g(x) = -x$ we write (65) as (using the notation Y_t instead of X_t for sake of clarity)

(71)
$$
dY_t = \left(-Y_t + J + \lambda \sigma^2 \int_0^t \frac{dW_s}{\lambda^2 + \sigma^2 s}\right) dt + \lambda dW_t
$$

so that, taking the expected value of both sides

(72)
$$
d\mathbb{E}(Y_t) = (-\mathbb{E}(Y_t) + J)dt,
$$

and

$$
d(Y_t - \mathbb{E}(Y_t)) = \left(-(Y_t - \mathbb{E}(Y_t)) + \lambda \sigma^2 \int_0^t \frac{dW_s}{\lambda^2 + \sigma^2 s} \right) dt + \lambda dW_t.
$$

Equation (72) is readily integrated yielding

(73)
$$
\mathbb{E}(Y_t) = J + (\mathbb{E}(X_0) - J)e^{-t}
$$

This equation is identical to (69) : the two processes X and Y have the same mean. An application of Itô's formula to $(Y_t - \mathbb{E}(Y_t))^2$ yields

$$
d(Y_t - \mathbb{E}(Y_t))^2 = -2(Y_t - \mathbb{E}(Y_t))^2 dt + 2\lambda \sigma^2 (Y_t - \mathbb{E}(Y_t)) \int_0^t \frac{dW_s}{\lambda^2 + \sigma^2 s} dt
$$

+ 2\lambda (Y_t - \mathbb{E}(Y_t))dW_t + \lambda^2 dt.

Define $v(t) := \mathbb{E}[(Y_t - \mathbb{E}(Y_t))^2]$. Taking the expected value of both sides of the previous equation we obtain

(74)
$$
v(t) = v(0) - 2 \int_0^t v(s) ds + \lambda^2 t + 2\lambda \sigma^2 \int_0^t \mathbb{E}\left[(Y_s - \mathbb{E}(Y_s)) \int_0^s \frac{dW_\theta}{\lambda^2 + \sigma^2 \theta} \right] ds
$$

Define Z_t by

(75)
$$
dZ_t = \frac{dW_t}{\lambda^2 + \sigma^2 t}.
$$

So, we have

$$
d(Y_t - \mathbb{E}(Y_t)) = (-(Y_t - \mathbb{E}(Y_t)) + \lambda \sigma^2 Z_t)dt + \lambda dW_t.
$$

We apply Itô's product formula to $Z_t(Y_t - \mathbb{E}(Y_t))$:

$$
d(Z_t(Y_t - \mathbb{E}(Y_t))) =
$$

$$
(-Z_t(Y_t - \mathbb{E}(Y_t)) + \lambda \sigma^2 Z_t^2)dt + \lambda Z_t dW_t + \frac{\lambda}{\lambda^2 + \sigma^2 t}dt + \frac{Y_t - \mathbb{E}(Y_t)}{\lambda^2 + \sigma^2 t}dW_t.
$$

According to (75) we have

$$
\mathbb{E}[Z_t^2] = \frac{t}{\lambda^2(\lambda^2 + \sigma^2 t)}.
$$

Integrate between 0 and t, use $Z_0 = 0$, and take the expected value to obtain

$$
\mathbb{E}[Z_t(Y_t - \mathbb{E}(Y_t))] = \int_0^t - \mathbb{E}[Z_s(Y_s - \mathbb{E}(Y_s))] + \frac{1}{\lambda}ds.
$$

That is

$$
\mathbb{E}[Z_t(Y_t - \mathbb{E}(Y_t))] = \frac{1 - e^{-t}}{\lambda}.
$$

Thus (74) writes

$$
v(t) = v(0) - 2 \int_0^t v(s) \, ds + \lambda^2 t + 2\sigma^2 \int_0^t (1 - e^{-s}) \, ds.
$$

This yields

$$
v(t) = \sigma^2 + \frac{\lambda^2}{2} - 2\sigma^2 e^{-t} + (\sigma^2 + v(0) - \frac{\lambda^2}{2})e^{-2t},
$$

which is identical to (70) .

6. Uniqueness of Q

We prove the uniqueness of the measure Q in (19). Such a proof has been provided in [1] for the S-model in the case $J = 0$ and is not constructive. Our proof is valid for the H-model as well as the S-model for any value of J. It is based upon a fixed-point approach and is constructive. In effect it is the basis for a numerical algorithm described in Section 7.

To simplify notations we assume $\lambda = 1$ and rewrite the system (19) in this case using the results of the previous sections. We will be using several times in our proof Proposition 14 for different values of the kernel κ .

According to (31) in the case $c \equiv 0$ and $\kappa = \tilde{K}_{Q}^{t}$, (19) is equivalent to

(76)
$$
X_t = X_0 + \int_0^t g(X_s) ds + J \int_0^t \mathbb{E}^Q[f(X_s)] ds + W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H_Q(s, u) \widetilde{W}_u du \right) ds
$$

 W_t is Q -Brownian motion, \widetilde{W}_t is given by (32) where the correlation function \tilde{K}_Q^t is linked to $K_Q(t,s) = \sigma^2 \mathbb{E}_Q[f(X_t)f(X_s)]$ by (27). Finally the function H_Q is given by (33).

We consider the family of Polish spaces, Cartesian products of $L^2([0,t])$ with \mathcal{K}_t , the set of Trace-Class operators on $L^2([0,t])$, for $0 < t \leq T$. The space $L^2([0,t]) \times \mathcal{K}_t$ is endowed with the norm:

(77)
$$
\|(m,\bar{K})\|_{t}^{2} = \|m\|_{L^{2}([0,t])}^{2} + \|\bar{K}\|_{t,2}^{2} \quad \forall m \in L^{2}([0,t]), \ \bar{K} \in \mathcal{K}_{t}
$$

Given \bar{K} in \mathcal{K}_t we associate with it the elements $\bar{\tilde{K}}^t$ of \mathcal{K}_t by (27) (in the case $\lambda = 1$) and \bar{H}^t by (33) with $\kappa(t,s) = \tilde{K}^t(t,s) \mathbb{1}_{t \geq s}$. For (m,\bar{K}) in $L^2([0,t]) \times \mathcal{K}_t$ we consider the process $X^{m,\bar{K}}$ solution of the SDE

(78)
$$
X_t^{m,\bar{K}} = X_0 + \int_0^t g(X_s^{m,\bar{K}}) ds + J \int_0^t m(s) ds + W_t + \int_0^t \widetilde{W}_s ds + \int_0^t \left(\int_0^s H(s, u) \widetilde{W}_u du \right) ds
$$

where \widetilde{W} is given by (32) with $\kappa(t,s) = \widetilde{K}^t(t,s) \mathbb{1}_{t \geq s}$. We define the mapping Λ from $L^2([0,t]) \times \mathcal{K}_t$ as follows

(79)
$$
\Lambda(m,\bar{K}) = (\mathbb{E}\left[f(X_t^{m,\bar{K}})\right], \sigma^2 \mathbb{E}\left[f(X_t^{m,\bar{K}})f(X_s^{m,\bar{K}})\right])
$$

where $X_t^{m,\bar{K}}$ $t^{m,K}$ is the solution to (78).

Lemma 24. The mapping Λ maps $L^2([0,t]) \times \mathcal{K}_t$ to itself for all $0 < t \leq T$.

Proof. Clear from the definition (79) since f (respectively $X^{m,\bar{K}}$) is bounded for the H-model (respectively for the S-Model). \Box

We use the metric induced by the norm (77) :

(80)
$$
d_t((m_1, \bar{K}_1), (m_2, \bar{K}_2))^2 = ||m_1 - m_2||^2_{L^2([0,t])} + ||\bar{K}_1 - \bar{K}_2||^2_{t,2}
$$

Observe that if X is a solution to (76), then the corresponding pair (m_Q, \bar{K}_Q) is a fixed point of Λ and conversely if (m, K) is such a fixed point of Λ , (78) defines a solution to (76) on $[0, T]$. Our problem is now translated into a fixed point problem for which we prove the following contraction lemma

Lemma 25. For $t \leq T$, $\bar{K}_1, \bar{K}_2 \in \mathcal{K}_T, m_1, m_2 \in L^2([0, T])$, we have

(81)
$$
d_t(\Lambda((m_1, \bar{K}_1)), \Lambda((m_2, \bar{K}_2)))^2 \leq c_T \int_0^t d_s((m_1, \bar{K}_1), (m_2, \bar{K}_2))^2 ds.
$$

where d_t is defined by (80) and c_T is a constant depending only on T.

Proof. Let (m_1, \bar{K}_1) and (m_2, \bar{K}_2) be two elements of $L^2([0,T]) \times \mathcal{K}_T$, (p_1, \bar{G}_1) and (p_2, \bar{G}_2) their images by Λ , see (79). In order to simplify the reading and w.l.o.g. we assume $J = \sigma = 1$. Let $(X_t^1)_{0 \le t \le T}$ and $(X_t^2)_{0 \le t \le T}$ be the solutions to the corresponding equations (78). We write

$$
(82)
$$

$$
X_t^1 - X_t^2 = \int_0^t (g(X_s^1) - g(X_s^2)) ds + \int_0^t (m_1(s) - m_2(s)) ds
$$

+
$$
\int_0^t \int_0^s (\tilde{K}_1^s(s, u) - \tilde{K}_2^s(s, u)) dW_u ds
$$

+
$$
\int_0^t \left(\int_0^s H_1(s, u) \int_0^u \tilde{K}_1^u(u, v) dW_v du - \int_0^s H_2(s, u) \int_0^u \tilde{K}_2^u(u, v) dW_v du \right) ds
$$

By (79) and Cauchy-Schwarz we write

$$
(p_1(t) - p_2(t))^2 = \left(\mathbb{E}[f(X_t^1) - f(X_t^2)]\right)^2 \le \mathbb{E}[(f(X_t^1) - f(X_t^2))^2]
$$

Using the fact that f is Lipschitz continuous we conclude that

(83)
$$
(p_1(t) - p_2(t))^2 \leq C \mathbb{E}[(X_t^1 - X_t^2)^2]
$$

for some positive constant C.

By (79) and Cauchy-Schwarz again we write

$$
(G_1(t,s) - G_2(t,s))^2 = \left(\mathbb{E}[f(X_t^1)f(X_s^1) - f(X_t^2)f(X_s^2)] \right)^2
$$

\n
$$
\leq \left(\mathbb{E}[|f(X_t^1)| |f(X_s^1) - f(X_s^2)| + |f(X_s^2)| |f(X_t^1) - f(X_t^2)|] \right)^2
$$

\n
$$
\leq \left(\mathbb{E}\left[\sqrt{f(X_t^1)^2 + f(X_s^2)^2} \sqrt{(f(X_s^1) - f(X_s^2))^2 + (f(X_t^1) - f(X_t^2))^2} \right] \right)^2
$$

\n
$$
\leq \mathbb{E}\left[f(X_t^1)^2 + f(X_s^2)^2\right] \times \mathbb{E}\left[(f(X_s^1) - f(X_s^2))^2 + (f(X_t^1) - f(X_t^2))^2\right]
$$

Using the facts that f is bounded and Lipschitz continuous for the H-model, or is the identity and $|X_t^1|, |X_s^2|$ are a.s. bounded by A for the S-model we can write

(84)
$$
(G_1(t,s) - G_2(t,s))^2 \leq C \mathbb{E}[(X_t^1 - X_t^2)^2] + C \mathbb{E}[(X_s^1 - X_s^2)^2]
$$

for some other positive constant C.

We thus have to upperbound $\mathbb{E}[(X_t^1 - X_t^2)^2]$.

The application of Ito's formula to $(X_t^1 - X_t^2)^2$ yields

$$
\mathbb{E}[(X_t^1 - X_t^2)^2] = 2\mathbb{E}\left[\int_0^t (X_s^1 - X_s^2)(g(X_s^1) - g(X_s^2)) ds\right] +
$$

\n
$$
2\mathbb{E}\left[\int_0^t (m_1(s) - m_2(s))(X_s^1 - X_s^2) ds\right] +
$$

\n
$$
2\mathbb{E}\left[\int_0^t (\widetilde{W}_1(s) - \widetilde{W}_2(s)) ds\right] +
$$

\n
$$
2\mathbb{E}\left[\int_0^t (X_s^1 - X_s^2) \left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right) ds\right]
$$

The right hand side of this equation is the sum of the expectations of four terms. The third term being equal to

$$
\int_0^t \int_0^s (\tilde{K}_1^t(s, u) - \tilde{K}_2^t(s, u)) dW_u ds,
$$

its expectation is equal to 0.

We consider next the first term $2\mathbb{E}\left[\int_0^t (X_s^1 - X_s^2)(g(X_s^1) - g(X_s^2)) ds\right]$. In the case of the H-Model, $g(x) = -\alpha x$, $\alpha > 0$, and g is a decreasing function. g is also a decreasing function in the case of the S-model, as shown in Figure 1. Therefore in both cases $2\mathbb{E}\left[\int_0^t (X_s^1 - X_s^2)(g(X_s^1) - g(X_s^2)) ds\right] \leq 0.$

We have obtained

(85)
\n
$$
\mathbb{E}[(X_t^1 - X_t^2)^2] \le 2\mathbb{E}\left[\int_0^t (m_1(s) - m_2(s))(X_s^1 - X_s^2) ds\right] +
$$
\n
$$
2\mathbb{E}\left[\int_0^t (X_s^1 - X_s^2) \left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right) ds\right]
$$
\nFrom $2\mathbb{E}\left[\int_0^t (m_1(s) - m_2(s))(X_s^1 - X_s^2) ds\right]$.

 $\textbf{Term}\hspace{0.1cm} 2\mathbb{E}\left[\int_{0}^{t}(m_{1}(s)-m_{2}(s))(X_{s}^{1}-X_{s}^{2})\,ds\right]$ By the inequality

(86)
$$
ab \leq \frac{1}{2}(a^2 + b^2)
$$

true for any two real numbers a and b we obtain

$$
(87) \quad 2\mathbb{E}\left[\int_0^t (m_1(s) - m_2(s))(X_s^1 - X_s^2) \, ds\right]
$$

$$
\leq \int_0^t (m_1(s) - m_2(s))^2 \, ds + \int_0^t \mathbb{E}\left[(X_s^1 - X_s^2)^2\right] \, ds
$$

$$
= ||m_1 - m_2||_{L^2([0,t])}^2 + \int_0^t \mathbb{E}\left[(X_s^1 - X_s^2)^2\right] \, ds
$$

Term $2\mathbb{E}\left[\int_0^t (X^1_s-X^2_s)\left(\int_0^s (H_1(s,u)\widetilde{W}_1(s,u)-H_2(s,u)\widetilde{W}_2(s,u))\,du\right)\,ds\right]$ By the same inequality we have

$$
2\mathbb{E}\left[\int_0^t (X_s^1 - X_s^2) \left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right) ds\right] \le
$$

$$
\int_0^t \mathbb{E}\left[\left(X_s^1 - X_s^2\right)^2\right] ds +
$$

$$
\int_0^t \mathbb{E}\left[\left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right)^2\right] ds
$$

We proceed by upperbounding the second term in the right hand side of this inequality.

First note that

$$
H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u) =
$$

\n
$$
H_1(s, u)\left(\int_0^u \tilde{K}_1^u(u, v) dW_v\right) - H_2(s, u)\left(\int_0^u \tilde{K}_2^u(u, v) dW_v\right) =
$$

\n
$$
H_1(s, u)\left(\int_0^u (\tilde{K}_1^u(u, v) - \tilde{K}_2^u(u, v)) dW_v\right) + (H_1(s, u) - H_2(s, u))\left(\int_0^u \tilde{K}_2^u(u, v) dW_v\right),
$$

so that, by (86) and Cauchy-Schwarz,

$$
\left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right)^2 \le
$$

$$
2\int_0^s H_1(s, u)^2 du \times \int_0^s \left(\int_0^u (\tilde{K}_1^u(u, v) - \tilde{K}_2^u(u, v)) dW_v\right)^2 du +
$$

$$
2\int_0^s (H_1(s, u) - H_2(s, u))^2 du \times \int_0^s \left(\int_0^u \tilde{K}_2^u(u, v) dW_v\right)^2 du.
$$

We take the expected value of both sides of this inequality and apply twice Ito's isometry formula so that

$$
\mathbb{E}\left[\left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right)^2\right] \le
$$

$$
2\int_0^s H_1(s, u)^2 du \times \int_0^s \int_0^u (\tilde{K}_1^u(u, v) - \tilde{K}_2^u(u, v))^2 dv du +
$$

$$
2\int_0^s (H_1(s, u) - H_2(s, u))^2 du \times \int_0^s \int_0^u \tilde{K}_2^u(u, v)^2 dv du
$$

By (34) $(H_1)^2$ can be upperbounded by a constant depending only on T. Moreover by (24) $\int_0^s \int_0^u \tilde{K}_2^u(u, v)^2 dv du \leq \|\tilde{\tilde{K}}_2\|_{t,2}^2$. By definition of $\tilde{\tilde{K}}_2^t$, and using the fact

that $(\mathrm{Id} + \bar{K}_2^t)^{-1}$ is contracting (see Remark 11), we write

$$
\|\tilde{K}_2^t\|_{t,2} = \|\tilde{K}_2^t \circ (\text{Id} + \bar{K}_2^t)^{-1}\|_{t,2} \le \|\tilde{K}_2^t\|_{t,2} \, \|(\text{Id} + \bar{K}_2^t)^{-1}\|_{t,2} \le \|\tilde{K}_2^t\|_{t,2} \le C_T
$$

for some positive constant C_T by (25) and (28). To sum up, we have found that there exists a positive constant C_T such that

(88)
$$
\int_0^t \mathbb{E}\left[\left(\int_0^s (H_1(s, u)\widetilde{W}_1(s, u) - H_2(s, u)\widetilde{W}_2(s, u)) du\right)^2\right] ds \le
$$

$$
C_T \left(\int_0^t \int_0^s \int_0^u (\tilde{K}_1^u(u, v) - \tilde{K}_2^u(u, v))^2 dv du ds + \int_0^t \int_0^s (H_1(s, u) - H_2(s, u))^2 du ds\right)
$$

The term $\int_0^s \int_0^u (\tilde{K}_1^u(u,v) - \tilde{K}_2^u(u,v))^2 dv du$ is less than $\|\tilde{K}_1^t - \tilde{K}_2^t\|_{t,2}^2$ and

$$
\begin{split} \bar{K}_1^t - \bar{K}_2^t &= \bar{K}_1^t \circ (\text{Id} + \bar{K}_1^t)^{-1} - \bar{K}_2^t \circ (\text{Id} + \bar{K}_2^t)^{-1} \\ &= (\text{Id} + \bar{K}_1^t)^{-1} \circ \bar{K}_1^t - \bar{K}_2^t \circ (\text{Id} + \bar{K}_2^t)^{-1} \\ &= (\text{Id} + \bar{K}_1^t)^{-1} \circ (\bar{K}_1^t \circ (\text{Id} + \bar{K}_2^t) - (\text{Id} + \bar{K}_1^t) \circ \bar{K}_2^t) \circ (\text{Id} + \bar{K}_2^t)^{-1} \\ &= (\text{Id} + \bar{K}_1^t)^{-1} \circ (\bar{K}_1^t - \bar{K}_2^t) \circ (\text{Id} + \bar{K}_2^t)^{-1} .\end{split}
$$

We take the norm of the two sides of this equality and, using again the fact that the operators $(\mathrm{Id} + \bar{K}_1^t)^{-1}$ and $(\mathrm{Id} + \bar{K}_2^t)^{-1}$ are contracting (Remark 11), we obtain

(89)
$$
\|\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t\|_{t,2} \le \|\bar{K}_1^t - \bar{K}_2^t\|_{t,2},
$$

and therefore

(90)
$$
\int_0^t \int_0^s \int_0^u (\tilde{K}_1^u(u,v) - \tilde{K}_2^u(u,v))^2 dv du ds \leq T ||\bar{K}_1 - \bar{K}_2||_{t,2}^2
$$

Consider now the obvious inequality

$$
\int_0^t \int_0^s (H_1(s, u) - H_2(s, u))^2 du ds \le C_T ||\bar{H}_1 - \bar{H}_2||_{t, 2}^2.
$$

According to (33) in the case $\kappa(t,s) = \tilde{K}^t(t,s) \mathbb{1}_{t \ge s}$

$$
\bar{H}_1 - \bar{H}_2 = (\text{Id} - \bar{\tilde{K}}_1^t)^{-1} - (\text{Id} - \bar{\tilde{K}}_2^t)^{-1}
$$

=
$$
(\text{Id} - \bar{\tilde{K}}_1^t)^{-1} \circ (\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t) \circ (\text{Id} - \bar{\tilde{K}}_2^t)^{-1},
$$

so that we have

$$
\begin{aligned}\n\|\bar{H}_1 - \bar{H}_2\|_{t,2} &\leq \|([\mathrm{Id} - \bar{\tilde{K}}_1^t)^{-1}\|_{t,2} \|\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t \|_{t,2} \, \| (\mathrm{Id} - \bar{\tilde{K}}_2^t)^{-1} \|_{t,2} \\
&= \|\bar{H}_1 + \mathrm{Id} \|_{t,2} \, \|\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t \|_{t,2} \, \|\bar{H}_2 + \mathrm{Id} \|_{t,2} \\
&\leq (\|\bar{H}_1\|_{t,2} + \|\mathrm{Id}\|_{t,2}) \|\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t \|_{t,2} (\|\bar{H}_2\|_{t,2} + \|\mathrm{Id}\|_{t,2}) \\
&= (\|\bar{H}_1\|_{t,2} + 1) \|\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t \|_{t,2} (\|\bar{H}_2\|_{t,2} + 1).\n\end{aligned}
$$

Because of (34) $\|\bar{H}_1\|_{t,2}$ and $\|\bar{H}_2\|_{t,2}$ are upper-bounded by a constant depending only on T and thus

$$
\|\bar{H}_1 - \bar{H}_2\|_{t,2} \le C_T \|\bar{\tilde{K}}_1^t - \bar{\tilde{K}}_2^t\|_{t,2},
$$

and, by (89),

(91)
$$
\|\bar{H}_1 - \bar{H}_2\|_{t,2} \leq C_T \|\bar{K}_1^t - \bar{K}_2^t\|_{t,2}
$$

for some positive constant C_T . We have obtained

(92)
$$
\int_0^t \int_0^s (H_1(s, u) - H_2(s, u))^2 du ds \leq C_T ||\bar{K}_1 - \bar{K}_2^t||_{t, 2}.
$$

Now by combining (85)-(92) we obtain

$$
\mathbb{E}[(X_t^1 - X_t^2)^2] \le C_T \left(\int_0^t \mathbb{E}[(X_s^1 - X_s^2)^2] ds + ||m_1 - m_2||_{L^2([0,t])}^2 + ||\bar{K}_1 - \bar{K}_2||_{t,2}^2 \right),
$$

so that Gronwall's Lemma commands that (93)

$$
\mathbb{E}[(X_t^1 - X_t^2)^2] \le C_T e^{TC_T} \left(\|m_1 - m_2\|_{L^2([0,t])}^2 + \|\bar{K}_1 - \bar{K}_2\|_{t,2}^2 \right) = D_T d_t((m_1, \bar{K}_1), (m_2, \bar{K}_2))
$$

for some positive constant D_T .

To conclude we have

$$
d_t(\Lambda((m_1,\bar{K}_1)),\Lambda((m_2,\bar{K}_2)))^2 = ||p_1 - p_2||^2_{L^2([0,t]} + ||\bar{G}_1 - \bar{G}_2||^2_{t,2},
$$

and, by (83) and (84),

$$
d_t(\Lambda((m_1, \bar{K}_1)), \Lambda((m_2, \bar{K}_2)))^2
$$

\n
$$
\leq C_T \left(\int_0^t \mathbb{E}[(X_s^1 - X_s^2)^2]ds + \int_0^t \int_0^t \mathbb{E}[(X_u^1 - X_u^2)^2]du ds \right).
$$

By (93)

$$
d_t(\Lambda((m_1, \bar{K}_1)), \Lambda((m_2, \bar{K}_2)))^2 \le c_T \int_0^t d_s((m_1, \bar{K}_1), (m_2, \bar{K}_2))^2 ds,
$$

for some positive constant c_T , which finishes the proof. \Box

We can now state the Theorem:

Theorem 26. The map Λ has a unique fixed point equal to (m_Q, \bar{K}_Q) in (19).

Proof. The existence and uniqueness of the fixed point are classically obtained by iterating Lemma 25 to obtain

$$
d_T(\Lambda^{k+1}((m,\bar{K})),\Lambda^k((m,\bar{K})))^2 \le \frac{c_T^k T^k}{k!} d_T(\Lambda((m,\bar{K})),(m,\bar{K}))^2.
$$

Existence of a fixed point arises from the fact that $\Lambda^k((m,\bar{K}))$ is a Cauchy sequence, hence convergent. Uniqueness is proved by assuming the existence of two fixed points $(m_1^{\infty}, \bar{K}_1^{\infty})$ and $(m_2^{\infty}, \bar{K}_2^{\infty})$ which must satisfy

$$
d_t((m_1^{\infty}, \bar{K}_1^{\infty}), (m_2^{\infty}, \bar{K}_2^{\infty}))^2 \leq c_T \int_0^t d_s((m_1^{\infty}, \bar{K}_1^{\infty}), (m_2^{\infty}, \bar{K}_2^{\infty}))^2 ds,
$$

and by Grönwall Lemma this commands that

$$
d_t((m_1^{\infty}, \bar{K}_1^{\infty}), (m_2^{\infty}, \bar{K}_2^{\infty})) = 0 \quad 0 \le t \le T
$$

□

This Lemma provides us with the following estimate of the rate of convergence to the fixed point $(m^{\infty}, \bar{K}^{\infty})$:

Corollary 27. Rate of convergence to the Fixed Point K^{∞}

$$
d_T((m^{\infty}, \bar{K}^{\infty}), \Lambda^{\ell}((m, \bar{K})))^2 \leq \sum_{j \geq \ell} d_T(\Lambda^{j+1}((m, \bar{K})), \Lambda^j((m, \bar{K})))^2
$$

$$
\simeq \frac{c_T^{\ell} T^{\ell}}{\ell!} d_T(\Lambda((m, \bar{K})), (m, \bar{K}))^2
$$

It also implies that $m_Q(t)$ is zero in the S-model case.

Corollary 28. If μ_0 is symmetric (which implies $\mathbb{E}[X_0] = 0$), the S-model satisfies $m_Q(t) = 0 \quad \forall \ 0 \le t \le T, \quad \forall J.$

Proof. In the S-model case, the function f is the identity, the function q is defined by (4) . Using the symmetry of the Brownian motion and the fact that f and q are odd, we immediately deduce that for any process X_t solution of (19) , $-X_t$ is also a solution. If in addition μ_0 is symmetric, the processes X_t and $-X_t$ start with the same law. By uniqueness of the solution, we deduce that for any $t \geq 0$, the law of X_t and $-X_t$ are equal. As a consequence, we have $\mathbb{E}[X_t] = 0$. □

7. Numerical simulations of the limit dynamics

We have implemented the fixed point algorithm Λ in the Julia language. We choose a time T, a time step Δt and we estimate a realisation of the solution on $[0, T]$ by discretising (19). For simplicity again we only consider the case $\lambda = 1$.

We note $\Delta \Phi_l$, $l = 0, \cdots, L-1$, $L = \lfloor \frac{1}{\Delta} \rfloor$ $\frac{T}{\Delta t}$, the difference $\Phi_{(l+1)\Delta t} - \Phi_{l\Delta t}$ for $\Phi \in \{W, B, X\}.$

At iteration $n \geq 1$ of the numerical approximation of Λ we are given $m^{(n-1)}(l\Delta t)$ and $\tilde{K}^{(n-1)}(l\Delta t, \overline{j}\Delta s)$, $j \leq l = 0, \cdots, L-1$, and we produce P independent

solutions of (19) noted $X_l^{n,p}$ $\ell_i^{n,p}, l = 0, \cdots, L, p = 1, \cdots, P$. These are obtained by drawing sequences of independent Brownian increments $\Delta W_l^{n,p} \simeq \mathcal{N}(0, \Delta t)$, $l = 0, \dots L - 1, p = 1, \dots, P, n \ge 1.$

Discretising (19) yields the equations

(94)
$$
\Delta X_l^{n,p} = g(X_l^{n,p})\Delta t + Jm^{(n-1)}(l\Delta t)\Delta t + \Delta B_l^{n,p}
$$

$$
\Delta B_l^{n,p} = \Delta W_l^{n,p} + \Delta t \sum_{j=0}^{l-1} \tilde{K}^{l\Delta t, (n-1)}(l\Delta t, j\Delta t)\Delta B_j^{n,p} \quad l = 0, \cdots L - 1,
$$

One then computes

$$
m^{(n)}(l\Delta t) = \frac{1}{P} \sum_{p} f(X_l^{n,p}) l = 0, \cdots, L-1
$$

$$
K^{(n)}(l\Delta t, j\Delta t) = \frac{1}{P} \sum_{p} f(X_l^{n,p}) f(X_j^{n,p}) j, l = 0, \cdots, L-1.
$$

The iteration of the map Λ produces a sequence of symmetric functions $K^{(n)}(t,s)$, $0 \leq s, t \leq T$ and $\tilde{K}^{t,(n)}(s,u), 0 \leq u \leq s \leq t$. In the discrete time setting these functions are approximated by the $L \times L$ symmetric matrices $K^{(n)}(l\Delta t, j\Delta t)$, $0 \leq$ $j, l \leq L-1$ and the L $i \times i$ symmetric matrices $\tilde{K}^{i\Delta t, (n)}(l\Delta t, j\Delta t)$, $0 \leq j, l \leq i-1$, $i = 1, \cdots, L$.

Equation (27) indicates that the *i*th symmetric $i \times i$ matrix $\tilde{K}^{i\Delta t,(n)}$ is obtained from the $i \times i$ sub-matrix $K^{i\Delta t, (n)}$ of the $L \times L$ matrix $K^{(n)}$ by

$$
\tilde{K}^{i\Delta t,(n)} = K^{i\Delta t,(n)} \left(\mathrm{Id}_i + \Delta t K^{i\Delta t,(n)} \right)^{-1} = \left(\mathrm{Id}_i + \Delta t K^{i\Delta t,(n)} \right)^{-1} K^{i\Delta t,(n)},
$$

where Id_i is the $i \times i$ identity matrix.

We then use $m^{(n)}$ and $\tilde{K}^{(n)}$ in (94) to compute the next approximation of Λ .

Remark 29. Note that since in the second equation of (94) only the lth row of $K^{l\Delta t,(n-1)}$ is needed, we do not have to compute $(\mathrm{Id}_l + \Delta t K^{l\Delta t,(n-1)})^{-1}$ but we rather solve for x the linear system

$$
\left(\mathrm{Id}_l + \Delta t K^{l\Delta t, (n-1)}\right)x = K^{l\Delta t, (n-1)}e_l,
$$

where e_l is the *l*th vector of the standard basis of \mathbb{R}^l . This can be achieved very efficiently by, for example, the conjugate gradient algorithm.

7.1. Numerical simulation results: the H-model.

We consider the case where the function f is given by

$$
f(x) = \frac{1}{2}(1 + \tanh(x))
$$

As shown in Figure 2 the minimum activity is 0, the maximum is 1. This is the function used in most works in mathematical neuroscience. We show some results of the estimations produced by the iterative method Λ as a set of composite figures.

FIGURE 2. The sigmoid function used in the numerical experiments for the H-model. Note that it varies between 0 and 1.

Each figure displays, on the first row, $K_Q(t, s)$, $\tilde{K}_Q^t(t, s)$, and the covariance of the solution X_t , Cov (X_t, X_s) , on the second row, the functions $K_Q(t, t)$, $\tilde{K}_Q^t(t, t)$, and $Cov(X_t, X_t)$; on the third row we display the function $m_Q(t)$, and two "typical" trajectories X_t^i and X_t^j as well as the corresponding activities $f(X_t^i)$ and $f(X_t^j)$ $\binom{J}{t}$ for various values of J and σ . Together with the estimation of K_Q and m_Q we obtain an estimation of the standard deviation of these estimates.

We start with an example where $J = 0.0$. Figure 3 shows the results as an array of plots. The function $m_O(t)$ is identically 0 by construction and thus not shown. The plot of $K_Q(t, t)$ in this figure shows three curves, $K_Q(t, t)$ in blue which falls between a red and a green curve. The red curve is $K_Q(t, t)$ plus twice the estimated standard deviation while the green is $K_Q(t, t)$ minus twice the estimated standard deviation. Both figures are for $\Delta t = 0.04$, the left one is for 100 000 trajectories while the right one is for 500 000 trajectories, yielding a more accurate estimation. In both cases we have iterated Λ ten times.

We now increase the excitation level and consider the case $J = 1$, shown in figure 4. The function $m_O(t)$ is non zero and converges when time increases to a constant value slightly below 0.7 and acts as a constant injected current resulting in increased values of the potential values X_t and activity values $f(X_t)$.

We next double the excitation level and go to $J = 2$. The function m_Q then saturates at 1.8 resulting in a further increase in the activity level shown as $f(X_t)$, its values getting much more often very close to the maximum value of 1. If we increase the excitation level to $J = 5$ we note that, as expected, the function $m_O(t)$ saturates at an even higher value, close to 5, resulting in activity levels very close to the maximum value of 1.

We now turn to negative values of J . Figure 5 has a similar structure to 4. Decreasing the values of J from -1 to -2.5 and to -5 decreases the injected current $m_O(t)$ resulting in a decrease of the neuronal acivity $f(X_t)$.

7.2. Numerical simulation results: the S-model.

FIGURE 3. H-model: $J = 0$, $\sigma = 1$. The computation is performed on the time interval [0, 8.0] with $\Delta t = 0.04$. The Monte Carlo method for estimating the fixed point of Λ uses 100 000 trajectories (Left) and 500 000 trajectories (Right). Λ is iterated 10 times. Note the improvement in the accuracy of the estimation of K_Q .

We next show some results of the estimations produced by the iterative method Λ for the S-model, also as a set of composite figures. Since we know from Corollary 28 that $m_Q(t)$ is zero if the initial condition is zero mean, we do not show it. Moreover, changing the value of J has no effect on K_Q hence we only show the case $J = 0$. Each figure displays, on the first row, from left to right two "typical" trajectories $X^{i}(t)$ and $X^{j}(t)$. The second row shows $K_{Q}(t, s)$ and $\tilde{K}_{Q}^{t}(t, s)$. Note that for the S-model $K_Q(t, s)$ is the correlation $\mathbb{E}_Q(X_t X_s)$ and thus we do not show $Cov(X_t, X_s)$ in this case. The third row shows the two functions $K_Q(t, t)$ and $\tilde{K}_Q^t(t,t)$.

As a general comment, this case is more difficult numerically than the H-model: If Δt is chosen too large, some of the generated trajectories exit the interval $|-A A$. To remedy this problem, following Ethier and Kurtz, [10] we change the definition (3) to

$$
U(x) = -k \log(A^2 - x^2),
$$

where in our work $k = 2, 4$. This change has the effect of having much less exiting trajectories.

FIGURE 4. H-model: From left to right and top to bottom, $J =$ 1, 2, 5 and $\sigma = 1$. The computation is performed on the time interval [0, 10.0] with $\Delta t = 0.04$. The Monte Carlo method for the fixed point computation uses 160 000 trajectories.

Figure 6 shows the results when $\sigma = \lambda = 1$ and the initial condition has 0 mean, hence $m_O(t) = 0$ (left part of the figure), or when the initial condition has mean 0.5 (right part of the figure where we observe that $m_Q(t)$ decreases to 0).

UNIVERSALITY OF MEAN-FIELD EQUATIONS 43

FIGURE 5. H-model: From left to right and top to bottom, $J =$ $-1, -2.5, -5$ and $\sigma = 1$. The computation is performed on the time interval [0, 5.0] with $\Delta t = 0.04$. The Monte Carlo method for the fixed point computation uses 100 000 trajectories.

Figure 7 shows the results when $\sigma = 2.0$ and the initial condition has 0 mean, hence $m_Q(t) = 0$ (left part of the figure), or when the initial condition has mean

FIGURE 6. S-model: $\sigma = 1$. Left: zero mean initial condition. $m_O(t) = 0$ at all times. Right: initial condition has mean 0.5. and $J = 1$. $m_Q(t)$ decreases to 0. The computation is performed on the time interval [0, 3.0] with $\Delta t = 0.01$. The Monte Carlo method for the fixed point computation uses 100 000 trajectories.

0.5 (right part of the figure and $J = 1$). We observe again that $m_O(t)$ decreases to 0. The bottom image shows the same case with a zero-mean initial condition. The estimation of the mean which we know to be equal to 0 is quite noisy.

7.3. Numerical verification of the universality of equations (19).

Thanks to [9] we know that the limit measure Q is independent of the actual distribution of the weights $J_{i \leftarrow j}$, subject to the conditions (15). As a consequence, the equations (19) that describe the limit dynamics are also independent of this distribution. We illustrate this with the following example. Consider the case of the H-model with Gaussian weights where the $J_{i\leftarrow j}$ are i.i.d. as Gaussians of the **n**-model with Gaussian weights where the $J_{i\leftarrow j}$ s are i.i.d. as Gaussians $\mathcal{N}(J/N, \sigma/\sqrt{N})$, and $\lambda = 1$. The results of the estimation of K_Q and \tilde{K}_Q are shown in Figure 4. According to the universality property the same limit Q is obtained when the $J_{i\leftarrow j}$ s are distributed, e.g., as

(95)
$$
J_{i \leftarrow j} \simeq \frac{J}{N} + \frac{\sigma}{\sqrt{N}} \left(\frac{\text{Ber}_{ij}(p)}{p} - 1 \right) \sqrt{\frac{p}{1 - p}},
$$

where the $Ber_{ij}(p)$ s are i.i.d. as Bernoulli random variables with parameter p, $0 \leq p \leq 1$. It is clear that $\mathbb{E}[J_{i \leftarrow j}] = J/N$ and $var(J_{i \leftarrow j}) = \sigma^2/N$. The limit

measures are therefore the same in the two cases *independently* of the value of p . We have simulated a fully connected network of N neurons defined by (6) where the $J_{i\leftarrow j}$ s are i.i.d. as (95) and estimated K_Q as

$$
K_Q(t,s) = \frac{1}{N} \sum_{i=1}^{N} f(X_t^i) f(X_s^i)
$$

We show in Figure 8 the covariance functions K_Q estimated from the network equations (6) and the mean-field dynamics (19). The agreement is qualitatively very good. In order to be slightly more precise in our comparison, we show in Figure 9 the variances $K_Q(t, t)$ (Left) and the means $m_Q(t)$ estimated from the network equations (6) (Left) and from the mean-field dynamics (19) (Right) together with the estimated standard deviations.

7.4. Miscellaneous results.

To finish this section of numerical experiments, we show results for two variations of the H-model. The first one is the model of Sompolinsky et al. [25, 7, 24, 8] where the function f is the odd sigmoid $f(x) = \tanh(x)$ shown in Figure 10.

Because this is an odd function, Corollary 28 applies and, given a zero-mean initial value, $m_Q(t) = 0$ for $0 \le t \le T$. This is shown in Figure 11.

We next show two results when the function f is the ReLu function, $f(x) =$ $\max(x, 0)$. It does not satisfy the boundedness hypothesis hence our results do not apply. The reason why we show some results is that this function is used in the field of AI as the main nonlinearity in deep recursive neural networks, see e.g. [18, 20]. Figure 12 shows two results. On the left the excitatory case when $J = 1$. The reader will appreciate the fact that $m_O(t)$ seems to be growing without limit (exploding), probably due to the fact that f is not upbounded, On the right we show the inhibitory case when $J = -1$. The situation seems very different, $m_Q(t)$ levels off, probably due to the fact that f is bounded below.

Acknowledgements The authors thank Romain Veltz for his help in developing the Julia code implementing the Λ algorithm. E.T. thanks the support of the ANR project ChaMaNe (ANR-19-CE40-0024, CHAllenges in MAthematical NEuroscience)

REFERENCES

- [1] G. Ben Arous and A. Guionnet. Large deviations for Langevin spin glass dynamics. Probab. Theory Related Fields, 102(4):455–509, 1995.
- [2] A. N. Borodin. Stochastic processes. Probability and its Applications. Birkhäuser/Springer, Cham, 2017.
- [3] T. Cabana and . D. Touboul. Large deviations for randomly connected neural networks: II. State-dependent interactions. Adv. in Appl. Probab., 50(3):983–1004, 2018.
- [4] T. Cabana and J. D. Touboul. Large deviations, dynamics and phase transitions in large stochastic and disordered neural networks. J. Stat. Phys., 153(2):211–269, 2013.

- [5] T. Cabana and J. D. Touboul. Large deviations for randomly connected neural networks: I. Spatially extended systems. Adv. in Appl. Probab., 50(3):944–982, 2018.
- [6] R. H. Cameron and W. T. Martin. Transformations of Wiener integrals under translations. Ann. of Math. (2), 45:386–396, 1944.
- [7] A. Crisanti and H. Sompolinsky. Dynamics of spin systems with randomly asymmetric bonds: Langevin dynamics and a spherical model. Phys. Rev. A (3), 36(10):4922–4939, 1987.
- [8] A. Crisanti and H. Sompolinsky. Path integral approach to random neural networks. Phys. Rev. E, 98:062120, Dec 2018.
- [9] A. Dembo, E. Lubetzky, and O. Zeitouni. Universality for Langevin-like spin glass dynamics. Ann. Appl. Probab., 31(6):2864–2880, 2021.
- [10] S. N. Ethier and T. G. Kurtz. Markov processes. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1986. Characterization and convergence.
- [11] O. Faugeras, J. Touboul, and B. Cessac. A constructive mean-field analysis of multi population neural networks with random synaptic weights and stochastic inputs. Front. Comput. Neurosci., 3, 2009.
- [12] I. V. Girsanov. On Transforming a Certain Class of Stochastic Processes by Absolutely Continuous Substitution of Measures. Theory of Probability $\mathcal C$ Its Applications, 5(3):285– 301, 1960.
- [13] A. Guionnet. *Dynamique de Langevin d'un verre de spins*. PhD thesis, Université Paris 11, 1995.
- [14] A. Guionnet. Averaged and quenched propagation of chaos for spin glass dynamics. Probab. Theory Related Fields, 109(2):183–215, 1997.
- [15] J. J. Hopfield. Neural networks and physical systems with emergent collective computational abilities. Proc. Nat. Acad. Sci. U.S.A., 79(8):2554–2558, 1982.
- [16] J. J. Hopfield. Neurons with graded response have collective computational properties like those of two-state neurons. Proceedings of the National Academy of Sciences, 81(10):3088– 3092, 1984.
- [17] J. J. Hopfield and D. W. Tank. Computing with neural circuits: A model. Science, 233(4764):625–633, 1986.
- [18] Y. LeCun, Y. Bengio, and G. Hinton. Deep learning. Nature, 521(7553):436–444, 2015.
- [19] O. Moynot and M. Samuelides. Large deviations and mean-field theory for asymmetric random recurrent neural networks. Probab. Theory Related Fields, 123(1):41–75, 2002.
- [20] E. Oostwal, M. Straat, and M. Biehl. Hidden unit specialization in layered neural networks: ReLU vs. sigmoidal activation. Phys. A, 564:Paper No. 125517, 14, 2021.
- [21] G. Peskir and A. Shiryaev. Optimal stopping and free-boundary problems. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2006.
- [22] M. Reed and B. Simon. Functional Analysis, (Methods of Modern Mathematical Physics), volume 1. Academic Press, 1981.
- [23] L. C. G. Rogers and D. Williams. Diffusions, Markov processes, and martingales. Vol. 2. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2000.
- [24] H. Sompolinsky, A. Crisanti, and H.-J. Sommers. Chaos in random neural networks. Phys. Rev. Lett., 61(3):259–262, 1988.
- [25] H. Sompolinsky and A. Zippelius. Relaxational dynamics of the Edwards-Anderson model and the mean-field theory of spin-glasses. Phys. Rev. B, 25:6860–6875, 1982.
- [26] F. G. Tricomi. Integral equations, volume Vol. V. Interscience Publishers, Inc., New York; Interscience Publishers Ltd., London, 1957.

[27] A. van Meegen, T. Kühn, and M. Helias. Large-deviation approach to random recurrent neuronal networks: parameter inference and fluctuation-induced transitions. Phys. Rev. Lett., 127(15):Paper No. 158302, 6, 2021.

O. Faugeras, CENTRE INRIA D'UNIVERSITÉ CÔTE D'AZUR, EPI CRONOS 2004, route des Lucioles, BP 93 06902 Sophia Antipolis Cedex FRANCE Olivier.Faugeras@inria.fr

E. Tanré, LABORATOIRE J.A.DIEUDONNÉ UMR CNRS-UNS N°7351 UNIVERSITÉ CÔTE D'AZUR PARC VALROSE 06108 NICE CEDEX 2 FRANCE Etienne.Tanre@inria.fr

FIGURE 7. S-model: $\sigma = 2$, $\lambda = 1$. Left: $J = 0$. $m_Q(t) = 0$ at all times. Right: initial condition has mean 0.5 and $J = 1$. $m_Q(t)$ decreases to 0. Bottom: $J = 1$, zero-mean initial condition. The computation is performed on the time interval [0, 3.0] with $\Delta t =$ 0.01. The Monte Carlo method for the fixed point computation uses 100 000 trajectories.

FIGURE 8. Comparison of the covariance functions $K_Q(t, s)$ estimated from the network equations (6) (Left) and from the meanfield dynamics (19) (Right, same as Figure 4, top left). The network size is 30,000. The Bernoulli parameter p is equal to 0.25.

FIGURE 9. Left: Comparison of the variance functions $K_Q(t, t)$ estimated from the network equations (6) and from the mean-field dynamics (19). Right: Comparison of the mean functions $m_Q(t)$. The Bernoulli parameter p is equal to 0.25.

FIGURE 10. The sigmoid function used in the work of Sompolinsky et al.. Note that it varies between -1 and 1 and is an odd function, as opposed to that of Figure 2.

FIGURE 11. H-model in the case of an odd sigmoïd function: $J =$ $\sigma = \lambda = 1$. Zero mean initial value. $m_Q(t) = 0$ at all times. The computation is performed on the time interval [0, 8.0] with $\Delta t =$ 0.04. The Monte Carlo method for the fixed point computation uses 100 000 trajectories.

UNIVERSALITY OF MEAN-FIELD EQUATIONS $51\,$

FIGURE 12. H-model in the case of the ReLu function. Left $J =$ $\sigma = \lambda = 1$. Zero mean initial value. $m_Q(t)$ appears to be growing without limit. Right: $-J = \sigma = \lambda = 1$. The computation is performed on the time interval [0, 10.0] with $\Delta t = 0.04$. The Monte Carlo method for the fixed point computation uses 100 000 trajectories.