



# Completion is an Instance of Abstract Canonical System Inference

Guillaume Burel, Claude Kirchner

## ► To cite this version:

Guillaume Burel, Claude Kirchner. Completion is an Instance of Abstract Canonical System Inference. 2006. inria-00000775v1

**HAL Id: inria-00000775**

**<https://inria.hal.science/inria-00000775v1>**

Submitted on 17 Nov 2005 (v1), last revised 12 Sep 2006 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Completion is an Instance of Abstract Canonical System Inference

Guillaume Burel and Claude Kirchner

LORIA (UMR 7503 CNRS-INPL-INRIA-Nancy2-UHP)  
Campus Scientifique, BP 239, 54506 Vandoeuvre-lès-Nancy Cedex, FRANCE

**Abstract.** Abstract canonical systems and inference (ACSI) were introduced in [1,2] to formalize the intuitive notions of good proof and good inference appearing typically in first-order logic or in Knuth-Bendix like completion procedures.

Since this abstract framework is intended to be generic, it is of fundamental interest to show its adequacy to represent the main systems of interest. This has been done for ground completion (where all equational axioms are ground) in [3] and was an open question for the general completion process.

By showing that the standard completion [4] is an instance of the ACSI framework we close the question. For this purpose, two proof representations, proof terms and proofs by replacement, are compared to build a proof ordering that provides an instantiation adapted to the abstract canonical system framework.

**Classification:** Logic in computer science, rewriting and deduction

## 1 Introduction

The notion of good proof is central when mechanizing deduction, in particular for defining useful tactics in proof assistant and theorem provers.

Motivated on one hand by this quest to *good proof* theory and on the other by the profound similarities between many proof search approaches, Dershowitz and Kirchner proposed in [1,5] a general framework based on ordering the set of proofs. In this context the best proofs are simply the minimal one.

Once one has defined what the best proofs are by the mean of a proof ordering, the next step is to obtain the best presentation of a theory, i.e. the set of axioms necessary for obtaining the best proofs for all the theory, but not containing anything useless.

To formalize this, the notion of *good inference* was introduced by Bonacina and Dershowitz [2]. Given a theory, its canonical presentation is defined as the set of the axioms needed to obtain the minimal proofs. It is general enough to produce all best proofs, leading to a notion of *saturation*, but it does not contain any redundant informations, hence the notion of *contraction*. Presentations, i.e. sets of axioms, are then transformed using appropriate deduction mechanisms to produce this canonical presentation.

This led to the Abstract Canonical Systems and Inference (ACSI) generic framework presented in [5,2].

The ACSI framework got its sources of inspiration from three related points. First, the early works on *Proof orderings* as introduced in [6] and [7] to prove the completeness of completion procedures *a la* Knuth-Bendix. Second, the developments about redundancy [8,9] to focus on the important axioms to perform further inferences. Last but not least, by the completion procedure [4], central in most theorem proving tools where an equality predicate is used. This procedure has been refined, mainly for two purposes: to have a more specific and thus more efficient algorithm when dealing with particular cases, or to increase the efficiency although remaining general. For the first case, a revue of specific completion procedures for specific algebraic structures can be found in [10]. For the second case, completion has been extended to equational completion [11,12,13], inductionless induction (initiated by Musser; see [14]), and ordered completion [15,8,7], to mention only a few.

But even if initially motivated by these three points, the ACSI framework has been developed as a full stand alone theory. This theory provide important abstract results based on basic hypothesis on proofs and a few postulates.

Therefore, a main question remains: is this framework indeed useful? Does this theory allows to *uniformly* understand and prove the main properties of a proof system, centered around the appropriate ordering on proofs?

At the price of a slight generalization of two postulates, it is shown in [16], that good proofs in natural deduction are indeed the cut free proofs as soon as proofs are compared using the ordering induced by beta reduction over the simply typed lambda-terms. For ground completion, the adequacy of the framework has been shown in [3], leaving the more general question of standard completion open.

This paper proves the adequacy of the framework to the standard completion procedure, generalizing in a non trivial way the result of [3] and showing the usefulness of abstract canonical systems.

This brings serious hopes that the ACSI framework is indeed well adapted and useful to uniformly understand and work with other algorithms, such as Buchberger's Gröbner basis algorithm for polynomial ideals [17,18].

The next section will summarize the framework of abstract canonical systems, as defined in [5,2], and briefly recall the standard completion. Section 3 deals with two representations of proofs in equational logic, namely as proof terms in the rewriting logic [19], and as proof by replacement [6]. We will show how to combine them to keep the tree structure of the first one, and the ordering associated with the second one, which is well adapted to prove the completeness of the standard completion. Finally, in Section 4, we will apply the abstract canonical systems framework to this proof representation to show the completeness of the standard completion. The proofs details are given in the Appendix.

## 2 Presentation

### 2.1 Abstract Canonical Systems

The results in this section are extracted from [5,2], which should be consulted for motivations, details and proofs.

Let  $\mathbb{A}$  be the set of all formulæ over some fixed vocabulary. Let  $\mathbb{P}$  be the set of all proofs. These sets are linked by two functions:  $[\cdot]^{Pm} : \mathbb{P} \rightarrow 2^{\mathbb{A}}$  gives the *premises* in a proof, and  $[\cdot]_{Cl} : \mathbb{P} \rightarrow \mathbb{A}$  gives its *conclusion*. Both are extended to sets of proofs in the usual fashion. The set of proofs built using assumptions in  $A \subseteq \mathbb{A}$  is noted by

$$Pf(A) \stackrel{!}{=} \{p \in \mathbb{P} : [p]^{Pm} \subseteq A\} .$$

The framework proposed here is predicated on two *well-founded* partial orderings over  $\mathbb{P}$ : a *proof ordering*  $>$  and a *subproof relation*  $\triangleright$ . They are related by a monotonicity requirement (postulate E). We assume for convenience that the proof ordering only compares proofs with the same conclusion ( $p > q \Rightarrow [p]_{Cl} = [q]_{Cl}$ ), rather than mention this condition each time we have cause to compare proofs.

We will use the term *presentation* to mean a set of formulæ, and *justification* to mean a set of proofs. We reserve the term *theory* for deductively closed presentations:

$$Th A \stackrel{!}{=} [Pf(A)]_{Cl} = \{[p]_{Cl} : p \in \mathbb{P}, [p]^{Pm} \subseteq A\} .$$

Theories are monotonic:

**Proposition 1 (Monotonicity).** *For all presentations A and B:*

$$A \subseteq B \Rightarrow Th A \subseteq Th B$$

Presentations  $A$  and  $B$  are *equivalent* ( $A \equiv B$ ) if their theories are identical:  $Th A = Th B$ . In addition to this, we assume the two following postulates:

**Postulate A (Reflexivity).** *For all presentations A:*

$$A \subseteq Th A$$

**Postulate B (Closure).** *For all presentations A:*

$$Th Th A \subseteq Th A$$

We call a proof *trivial* when it proves only its unique assumption and has no subproofs other than itself, that is, if  $[p]^{Pm} = \{[p]_{Cl}\}$  and  $p \triangleright q \Rightarrow p = q$ . We denote by  $\hat{a}$  such a trivial proof of  $a \in \mathbb{A}$  and by  $\hat{A}$  the set of trivial proofs of each  $a \in A$ .

We assume that proofs use their assumptions (postulate C), that subproofs don't use non-existent assumptions (postulate D), and that proof orderings are monotonic with respect to subproofs (postulate E):

**Postulate C (Trivial).** For all proofs  $p$  and formulæ  $a$ :

$$a \in [p]^{Pm} \Rightarrow p \triangleright \hat{a}$$

**Postulate D (Subproofs Premises Monotonicity).** For all proofs  $p$  and  $q$ :

$$p \triangleright q \Rightarrow [p]^{Pm} \supseteq [q]^{Pm}$$

**Postulate E (Replacement).** For all proofs  $p, q$  and  $r$ :

$$p \triangleright q > r \Rightarrow \exists v \in Pf([p]^{Pm} \cup [r]^{Pm}). p > v \triangleright r$$

We make no other assumptions regarding proofs or their structure and the proof ordering  $>$  is lifted to an ordering  $\succsim$  over presentations:

$$A \succsim B \text{ if } A \equiv B \text{ and } \forall p \in Pf(A) \exists q \in Pf(B). p \geq q .$$

We define what a *normal-form proof* is, i.e. the minimal proofs of  $Pf(Th A)$ :

$$Nf(A) \stackrel{!}{=} \mu Pf(Th A) \stackrel{!}{=} \{p \in Pf(Th A) : \neg \exists q \in Pf(Th A). p > q\} .$$

The *canonical presentation* contains those formulæ that appear as assumptions of normal-form proofs:

$$A^\# \stackrel{!}{=} [Nf(A)]^{Pm} .$$

So, we will say that  $A$  is *canonical* if  $A = A^\#$ .

A presentation  $A$  is *saturated* if it supports all possible normal form proofs:

$$Pf(A) \supseteq Nf(A) .$$

The set of all *redundant formulæ* of a given presentation  $A$  will be denoted as follows:

$$Red A \stackrel{!}{=} \{r \in A : A \succsim A \setminus \{r\}\} .$$

and a presentation  $A$  is *contracted* if

$$Red A = \emptyset .$$

The following main result can then be derived [1]:

**Theorem 1.** *A presentation is canonical iff it is saturated and contracted.*

We now consider inference and deduction mechanisms. A *deduction mechanism*  $\rightsquigarrow$  is a function from presentations to presentations and we call the relation  $A \rightsquigarrow B$  a *deduction step*. A sequence of presentations  $A_0 \rightsquigarrow A_1 \rightsquigarrow \dots$  is called a *derivation*. The *result* of the derivation is, as usual, its *persisting* formulæ:

$$A_\infty \stackrel{!}{=} \liminf_{j \rightarrow \infty} A_j = \bigcup_{j > 0} \bigcap_{i > j} A_i .$$

A deduction mechanism  $\rightsquigarrow$  is *sound* if  $A \rightsquigarrow B$  implies  $Th B \subseteq Th A$ . It is *adequate* if  $A \rightsquigarrow B$  implies  $Th A \subseteq Th B$ . It is *good* if proofs only get better:

$$\rightsquigarrow \subseteq \succsim .$$

A derivation  $A_0 \rightsquigarrow A_1 \rightsquigarrow \dots$  is *good* if  $A_i \succsim A_{i+1}$  for all  $i$ .

We now extend the notion of saturation and contraction to derivation:

- A derivation  $\{A_i\}_i$  is *saturating* if  $A_\infty$  is saturated.
- It is *contracting* if  $A_\infty$  is contracted.
- It is *canonical* if both saturating and contracting.

A canonical derivation can be used to build the canonical presentation of the initial presentation:

**Theorem 2.** *A good derivation is canonical if and only if*

$$A_\infty = A_0^\# .$$

## 2.2 The Standard Completion

The standard completion algorithm was first introduced by Knuth and Bendix in [4], hence the name it is often called. Its correctness was first shown by Huet in [20], using a fairness hypothesis. We use here a presentation of this algorithm as inference rules (see Fig. 1), as can be found in [6]. For basics on rewritings and completions, we refer to [21,22].

The Knuth-Bendix algorithm consists in 6 rules which apply to a couple  $E, R$  of a set of equational axioms and a set of rewriting rules. It takes a reduction ordering  $\gg$  over terms as argument. The rules are presented in Fig. 1.

Since [20], standard completion is associated with a fairness assumption (see [6, Lemma 2.8]): at the limit, all equations are oriented ( $E_\infty = \emptyset$ ) and all persistent critical pairs coming from  $R_\infty$  are treated by **Deduce** at least once.

Because we work with terms with variables, the reduction ordering  $\gg$  can not be total, so that **Orient** may fail. Therefore, the standard completion algorithm may either:

- terminate with success and yield a terminating, confluent set of rules;
- terminate with failure; or
- not terminate.

Here, the completeness of the standard completion will only be shown using the ACSI framework for the first case.

## 3 Proof Representations

Our goal is now to use the ACSI framework to directly show that standard completion inference rules are correct and complete. We have therefore first to find the right order on proofs. We have two main choices that we are now defining and relating.

<sup>1</sup>  $\blacktriangleright$  design the encompassment ordering,  $s \blacktriangleright t$  if a subterm of  $s$  in an instance of  $t$  but not vice versa.

<b>Deduce:</b> If $(s, t)$ is a critical pair of $R$	
	$E, R \rightsquigarrow E \cup \{s = t\}, R$
<b>Orient:</b> If $s \gg t$	
	$E \cup \{s = t\}, R \rightsquigarrow E, R \cup \{s \rightarrow t\}$
<b>Delete:</b>	
	$E \cup \{s = s\}, R \rightsquigarrow E, R$
<b>Simplify:</b> If $s \xrightarrow[R]{u}$	
	$E \cup \{s = t\}, R \rightsquigarrow E \cup \{u = t\}, R$
<b>Compose:</b> If $t \xrightarrow[R]{u}$	
	$E, R \cup \{s \rightarrow t\} \rightsquigarrow E, R \cup \{s \rightarrow u\}$
<b>Collapse</b> <sup>1</sup> : If $s \xrightarrow[v \rightarrow w \in R]{u}$ , and $s \blacktriangleright v$ ,	
	$E, R \cup \{s \rightarrow t\} \rightsquigarrow E \cup \{u = t\}, R$

Fig. 1. Standard Completion Inference Rules.

### 3.1 Proof Terms

Let us first consider the proof representation coming from the one used in rewriting logic (introduced by Meseguer [19]). Consider a signature  $\Sigma$ , and a set of equational axioms  $E$  and a set of rewrite rules  $R$  based on this signature. We consider the rules of the equational logic given in the Fig. 2. These inference rules define the *proof term* associated with a proof. The notation  $\pi : t \longrightarrow t'$  means that  $\pi$  is a proof term—that could also be seen as a trace—showing that the term  $t$  can be rewritten to the term  $t'$ .

Notice that the terms based on the signature  $\Sigma$  are plunged into the proof terms when they are formed with the rules **Reflexivity** and **Congruence**. Notice also that **Reflexivity** for  $t \longrightarrow t$  is not essential because it can be replaced by a tree of **Congruence** isomorph to  $t$ . The proof terms associated are furthermore the same in both case:  $t$ . Notice also that these proof terms are a restricted form of rho-terms [23].

Some proof terms defined here are “essentially the same”. For instance, the transitivity operator should be considered as associative, so that the proofs  $(\pi_1; \pi_2); \pi_3$  and  $\pi_1; (\pi_2; \pi_3)$  are equal. This can be done by quotienting the proof terms algebra by the congruence rules of figure 3.

The rules **Associativity**, **Identities** and **Inverse** give a group structure to the proof terms algebra. We can therefore deduce for instance that the proofs  $(\pi_1; \pi_2)^{-1}$  and  $\pi_2^{-1}; \pi_1^{-1}$  are equivalent. We

<b>Reflexivity:</b>	$\overline{t : t \longrightarrow t}$
<b>Congruence:</b>	$\frac{\pi_1 : t_1 \longrightarrow t'_1 \quad \dots \quad \pi_n : t_n \longrightarrow t'_n}{f(\pi_1, \dots, \pi_n) : f(t_1, \dots, t_n) \longrightarrow f(t'_1, \dots, t'_n)}$
<b>Replacement:</b>	For all rules or equational axiom $\ell = (g(x_1, \dots, x_n), d(x_1, \dots, x_n)) \in E \cup R$ ,
	$\frac{\pi_1 : t_1 \longrightarrow t'_1 \quad \dots \quad \pi_n : t_n \longrightarrow t'_n}{\ell(\pi_1, \dots, \pi_n) : g(t_1, \dots, t_n) \longrightarrow d(t'_1, \dots, t'_n)}$
<b>Transitivity:</b>	$\frac{\pi_1 : t_1 \longrightarrow t_2 \quad \pi_2 : t_2 \longrightarrow t_3}{\pi_1; \pi_2 : t_1 \longrightarrow t_3}$
<b>Symmetry:</b>	$\frac{\pi : t_1 \longrightarrow t_2}{\pi^{-1} : t_2 \longrightarrow t_1}$

**Fig. 2.** Inference Rules for Equational Logic

similarly have  $f(\pi_1, \dots, \pi_n)^{-1}$  equivalent to  $f(\pi_1^{-1}, \dots, \pi_n^{-1})$ , because

$$\begin{aligned}
f(\pi_1^{-1}, \dots, \pi_n^{-1}) &\equiv f(\pi_1^{-1}, \dots, \pi_n^{-1}); t' \\
&\equiv f(\pi_1^{-1}, \dots, \pi_n^{-1}); (f(\pi_1, \dots, \pi_n); f(\pi_1, \dots, \pi_n)^{-1}) \\
&\equiv (f(\pi_1^{-1}, \dots, \pi_n^{-1}); f(\pi_1, \dots, \pi_n)); f(\pi_1, \dots, \pi_n)^{-1} \\
&\equiv f(\pi_1^{-1}; \pi_1, \dots, \pi_n^{-1}; \pi_n); f(\pi_1, \dots, \pi_n)^{-1} \\
&\equiv f(t_1, \dots, t_n); f(\pi_1, \dots, \pi_n)^{-1} \\
&\equiv f(\pi_1, \dots, \pi_n)^{-1}.
\end{aligned}$$

### 3.2 Proofs by Replacement of Equal by Equal

This proof representation was introduced by [6] to prove the completeness of the Knuth-Bendix completion algorithm, using an ordering over such proofs that decreases for every completion step.

An *equational proof step* is an expression  $s \xleftarrow[e]{p} t$  where  $s$  and  $t$  are terms,  $e$  is an equational axiom  $u = v$ , and  $p$  is a position of  $s$  such that  $s|_p = \sigma(u)$  and  $t = s[\sigma(v)]_p$  for some substitution  $\sigma$ .

An *equational proof* of  $s_0 = t_n$  is any finite sequence of equational proof steps  $\left( s_i \xleftarrow[e_i]{p_i} t_i \right)_{i \in \{0, \dots, n\}}$  such that  $t_i = s_{i+1}$  for all  $i \in \{0, \dots, n-1\}$ . It is noted

$$s_0 \xleftarrow[e_0]{p_0} s_1 \xleftarrow[e_1]{p_1} s_2 \cdots s_n \xleftarrow[e_n]{p_n} t_n.$$



<b>Associativity:</b> For all proof terms $\pi_1, \pi_2, \pi_3$ ,
$\pi_1; (\pi_2; \pi_3) \equiv (\pi_1; \pi_2); \pi_3$
<b>Identities:</b> For all proof terms $\pi : t \longrightarrow t'$ ,
$\pi; t' \equiv t; \pi \equiv \pi$
<b>Preservation of Composition:</b> For all proof terms $\pi_1, \dots, \pi_n, \pi'_1, \dots, \pi'_n$ , for all function symbols $f$ ,
$f(\pi_1; \pi'_1, \dots, \pi_n; \pi'_n) \equiv f(\pi_1, \dots, \pi_n); f(\pi'_1, \dots, \pi'_n)$
<b>Parallel Move Lemma:</b> For all rewrite rules or equational axiom $\ell = (g(x_1, \dots, x_n), d(x_1, \dots, x_n)) \in E \cup R$ , for all proof terms $\pi_1 : t_1 \longrightarrow t'_1, \dots, \pi_n : t_n \longrightarrow t'_n$ ,
$\begin{aligned} \ell(\pi_1, \dots, \pi_n) &\equiv \ell(t_1, \dots, t_n); d(\pi_1, \dots, \pi_n) \\ &\equiv g(\pi_1, \dots, \pi_n); \ell(t'_1, \dots, t'_n) \end{aligned}$
<b>Inverse:</b> For all proof terms $\pi : t \longrightarrow t'$ ,
$\begin{aligned} \pi; \pi^{-1} &\equiv t \\ \pi^{-1}; \pi &\equiv t' \end{aligned}$

Fig. 3. Equivalence of Proof Terms

A *rewrite proof step* is an expression  $s \xrightarrow[\ell]{p} t$  or  $t \xleftarrow[\ell]{p} s$  where  $s$  and  $t$  are terms,  $\ell$  is a rewrite rule  $u \rightarrow v$ , and  $p$  is a position of  $s$  such that  $s|_p = \sigma(u)$  and  $t = s[\sigma(v)]_p$  for some substitution  $\sigma$ .

An *proof by replacement (of equal by equal)* of  $s_0 = t_n$  is any finite sequence of equational proof steps and rewrite proof step  $\left( s_i \xrightarrow[\ell_i]{p_i} t_i \right)_{i \in \{0, \dots, n\}}$  where  $\xrightarrow[\ell_i]{p_i} \in \{\longleftrightarrow, \longrightarrow, \longleftarrow\}$  for  $i \in \{0, \dots, n\}$  and such that  $t_i = s_{i+1}$  for all  $i \in \{0, \dots, n-1\}$ . It is noted

$$s_0 \xrightarrow[\ell_0]{p_0} s_1 \xrightarrow[\ell_1]{p_1} s_2 \cdots s_n \xrightarrow[\ell_n]{p_n} t_n$$

### 3.3 From Proof Terms to Proofs by Replacement

In order to have a one to one correspondence between proof representations, we use the equivalence of proof terms defined in Fig. 3. We can refine them to the proof term rewrite system  $\rightsquigarrow$  given in Fig. 4, in which  $\pi, \pi', \pi_1, \dots$  range over proof terms,  $t, t', t_1, \dots$  over  $\Sigma$ -terms,  $f, g, d$  over function symbols,  $\ell$  over rules and equational axioms labels and  $i$  over  $\{1, \dots, n\}$ .

The associativity is still considered in the congruence, so that all proof terms rewrite rules must be considered modulo the associativity of  $;$  which will be noted

<b>Delete Useless Identities:</b>	
	$\left. \begin{array}{l} \pi; t' \\ t; \pi \end{array} \right\} \rightsquigarrow \pi$
<b>Sequentialization:</b> If $\pi_k : t_k \longrightarrow t'_k$ and there exists $i \neq j \in \{1, \dots, n\}$ such that $\pi_i \neq t_i$ and $\pi_j \neq t_j$ ,	
	$f(\pi_1, \dots, \pi_n) \rightsquigarrow f(\pi_1, t_2, \dots, t_n); f(t'_1, \pi_2, \dots, t_n); \dots; f(t'_1, t'_2, \dots, \pi_n)$
<b>Composition Shallowing:</b> If $\pi_i : t_i \longrightarrow t'_i$ and $\pi'_i : t'_i \longrightarrow t''_i$ ,	
	$f(t_1, \dots, \pi_i; \pi'_i, \dots, t_n) \rightsquigarrow f(t_1, \dots, \pi_i, \dots, t_n); f(t_1, \dots, \pi'_i, \dots, t_n)$
<b>Parallel Moves:</b> If $\ell = (g(x_1, \dots, x_n), d(x_1, \dots, x_n))$ , $\pi_1 : t_1 \longrightarrow t'_1, \dots, \pi_n : t_n \longrightarrow t'_n$ , and if there exists $i \in \{1, \dots, n\}$ such that $\pi_i \neq t_i$ ,	
	$\ell(\pi_1, \dots, \pi_n) \rightsquigarrow \ell(t_1, \dots, t_n); d(\pi_1, \dots, \pi_n)$
<b>Delete Useless Inverses:</b>	
	$t^{-1} \rightsquigarrow t$
<b>Inverse Congruence:</b> If $\pi_i : t_i \longrightarrow t'_i$ ,	
	$f(t_1, \dots, \pi_i^{-1}, \dots, t_n) \rightsquigarrow f(t_1, \dots, \pi_i, \dots, t_n)^{-1}$
<b>Inverse Composition:</b>	
	$(\pi_1; \pi_2)^{-1} \rightsquigarrow \pi_2^{-1}; \pi_1^{-1}$

Fig. 4. Rewrite System for Proof Terms

$\sim$ . The class rewrite system that we consider will be therefore noted  $\rightsquigarrow / \sim$ . As it is linear, we can use the framework and results from [11].

We first prove that this rewrite system is included in the equivalence relation of Fig. 3.

**Proposition 2 (Correctness).** *For all proof terms  $\pi_1, \pi_2$ , if  $\pi_1 \rightsquigarrow \pi_2$  then  $\pi_1 \equiv \pi_2$ .*

The converse is false: for instance  $f(\ell_1, \ell_2) \equiv f(t_1, \ell_2); f(\ell_1, t'_2)$  but we do not have  $f(\ell_1, \ell_2) \xrightarrow[\rightsquigarrow]{*} f(t_1, \ell_2); f(\ell_1, t'_2)$ .

**Proposition 3 (Termination and Confluence).** *The proof term rewrite system  $\rightsquigarrow$  modulo  $\sim$  is terminating and confluent modulo  $\sim$ .*

The proof terms rewrite system  $\rightsquigarrow$  allow us to give a correspondence between proof terms and proofs by replacement of equal by equal: normal forms of proof terms correspond exactly to proofs by replacement. This fact is expressed in the following theorem, which is indeed a generalization of Lemma 3.6 in [19] for

equational logic. We also have operationalized the way to construct the chain of “one-step sequential rewrites”.

**Theorem 3 (Correspondence between Proof Representations).** *The normal form of a proof term  $\pi$  for the rewrite system  $\rightsquigarrow$ , noted  $nf(\pi)$ , has the following form:*

*there exists  $n \in \mathbb{N}$ , some contexts  $w_1[], \dots, w_n[]$ , some indices  $i_1, \dots, i_n \in \{-1, 1\}$ , some rule labels  $\ell_1, \dots, \ell_n$  and some terms  $t_1^1, \dots, t_{m_1}^1, \dots, t_1^n, \dots, t_{m_n}^n$  such that*

$$nf(\pi) = (w_1[\ell_1(t_1^1, \dots, t_{m_1}^1)])^{i_1}; \dots; (w_n[\ell_n(t_1^n, \dots, t_{m_n}^n)])^{i_n}$$

where  $\nu^1$  is a notation for  $\nu$ .

We will denote by  $nf(\pi)$  the normal form of a proof term  $\pi$ .

Such a proof term correspond with the following proof by replacement of equal by equal:

$$w_1[g_1(t_1^1, \dots, t_{m_1}^1)] \xrightarrow[\ell_1]{p_1} w_1[d_1(t_1^1, \dots, t_{m_1}^1)] \xrightarrow[\ell_2]{p_2} \dots \xrightarrow[\ell_n]{p_n} w_n[d_n(t_1^n, \dots, t_{m_n}^n)]$$

where for all  $j \in \{1, \dots, n\}$  we have:

- $\ell_j = (g_j, d_j)$ ,
- $p_j$  is the position of  $[]$  in  $w_j[]$ ,
- $\xrightarrow[\ell_j]{p_j} = \longrightarrow$  if  $i_j = 1$  and  $\ell_j \in R$ ,  
 $\longleftarrow$  if  $i_j = -1$  and  $\ell_j \in R$ ,  
 $\longleftrightarrow$  if  $\ell_j \in E$ .
- if  $j \neq n$ ,  $w_j[d_j(t_1^j, \dots, t_{m_j}^j)] = w_{j+1}[g_{j+1}(t_1^{j+1}, \dots, t_{m_{j+1}}^{j+1})]$ .

*Example 1.* Consider  $\pi = f(\ell_1(\ell_2), (\ell_3; r)^{-1})$  where  $\ell_1 : g(x) \longrightarrow d(x)$ ,  $\ell_2 : s = t$ ,  $\ell_3 : l \longrightarrow r$ , we have:

$$\begin{array}{ll} \pi \xrightarrow{\sim} f(\ell_1(s); d(\ell_2), (\ell_3; r)^{-1}) & \text{(Parallel Moves)} \\ \xrightarrow{\sim} f(\ell_1(s); d(\ell_2), r); f(d(t), (\ell_3; r)^{-1}) & \text{(Sequentialization)} \\ \xrightarrow{\sim} f(\ell_1(s); d(\ell_2), r); f(d(t), r^{-1}; \ell_3^{-1}) & \text{(Inverse Composition)} \\ \xrightarrow{\sim} f(\ell_1(s); d(\ell_2), r); f(d(t), r; \ell_3^{-1}) & \text{(Delete Useless Inverses)} \\ \xrightarrow{\sim} f(\ell_1(s); d(\ell_2), r); f(d(t), \ell_3^{-1}) & \text{(Delete Useless Identities)} \\ \xrightarrow{\sim} f(\ell_1(s), r); f(d(\ell_2), r); f(d(t), \ell_3^{-1}) & \text{(Composition Shallowing)} \\ \xrightarrow{\sim} f(\ell_1(s), r); f(d(\ell_2), r); f(d(t), \ell_3)^{-1} & \text{(Inverse Congruence)} \end{array}$$

This last term is the normal form proof term, and it is equivalent to the proof by replacement  $f(g(s), r) \xrightarrow[\ell_1]{1} f(d(s), r) \xleftarrow[\ell_2]{11} f(d(t), r) \xleftarrow[\ell_3]{2} f(d(t), l)$ .

Due to this theorem, normal forms of proof terms can be considered in the following indifferently as proof terms or as proofs by replacement.

### 3.4 Proofs Ordering

The representation of Bachmair by the mean of proof by replacement was defined to introduce an order on proofs [6]: given a reduction ordering  $\gg$ , to each single proof steps  $s \xrightarrow[\ell]{p} t$  is associated a *cost*. The cost of an equational proof step  $s \xrightarrow[u=v]{p} t$  is the triple  $(\{s, t\}, u, t)$ . The cost of a rewrite proof step  $s \xrightarrow[u \rightarrow v]{p} t$  is  $(\{s\}, u, t)$ . Proof steps are compared with each other according to their cost, using the lexicographic combination of the multiset  $\gg_{mult}$  extension of the reduction ordering over terms in the first component, the encompassment ordering  $\blacktriangleright$  on the second component, and the reduction ordering  $\gg$  on the last component. Proofs are compared as multisets of their proof steps. For two proofs by replacement  $p, q$ , we will write  $p >_{rep} q$  if  $p$  is greater than  $q$  for such an ordering.

Using theorem 3, we can translate Bachmair's proof ordering to proof terms:

**Definition 1 (Bachmair's Ordering on Proof Terms).**

For all proof terms  $\pi_1, \pi_2$ , we say that  $\pi_1 >_B \pi_2$  iff

$$nf(\pi_1) >_{rep} nf(\pi_2)$$

*Example 2.* Consider  $\pi_1 = f(\ell_1^{-1}; \ell_2)$  and  $\pi_2 = f(\ell_3)$  where  $\ell_1 = a \rightarrow b$ ,  $\ell_2 = a \rightarrow c$  and  $\ell_3 = b = c$ , and suppose  $a > b > c$ .

We have  $nf(\pi_1) = f(b) \xleftarrow[\ell_1]{1} f(a) \xrightarrow[\ell_2]{1} f(c)$  and  $nf(\pi_2) = f(b) \xrightarrow[\ell_3]{1} f(c)$ . The cost of  $nf(\pi_1)$  is  $\{(\{f(a)\}, a, f(b)), (\{f(a)\}, a, f(c))\}$ , the cost of  $nf(\pi_2)$  is  $\{(\{f(b), f(c)\}, b, f(c))\}$ , so  $nf(\pi_1) >_{rep} nf(\pi_2)$  and  $\pi_1 >_B \pi_2$ .

Nevertheless, we cannot hope to extend a RPO on  $\Sigma$ -terms to a RPO<sup>2</sup><sub>rpo</sub> on proof terms so that  $>_B$  and  $>_{rpo}$  coincide for the normal forms of proof terms:

*Counter-example 3.* Suppose we have  $\Sigma = \{f^1, a^0, b^0, c^0\}$  where the exponents of function symbols denote their arity, and a precedence  $f > a > b > c$ . We consider the RPO based on such a precedence.

Let  $\ell_f = f(a) \rightarrow c$  and  $\ell_b = b \rightarrow c$ .

We now want to extend the precedence to  $\ell_f$  and  $\ell_b$  in order to extend the RPO to proof terms. If we have  $\ell_f < \ell_b$ ,  $f(a) \xrightarrow[\ell_f]{\epsilon} c >_{rep} b \xrightarrow[\ell_b]{\epsilon} c$  but  $\ell_f <_{rpo} \ell_b$ .

If we suppose  $f > \ell_f > \ell_b$  we have  $f(a) \xrightarrow[\ell_f]{\epsilon} c >_{rep} f(b) \xrightarrow[\ell_b]{1} f(c)$  but  $\ell_f <_{rpo} f(\ell_b)$ .

If we suppose  $\ell_f > \ell_b$  and  $\ell_f > f$ , then  $f(f(b)) \xrightarrow[\ell_b]{11} f(f(c)) >_{rep} f(a) \xrightarrow[\ell_f]{\epsilon} c$  but  $f(f(\ell_b)) <_{rpo} \ell_f$ .

Such an extension is therefore impossible, there is no extension of  $>_{rpo}$  on proof terms so that for all proof terms  $\pi_1, \pi_2$ , we have  $nf(\pi_1) >_{rpo} nf(\pi_2)$  if and only if  $nf(\pi_1) >_B nf(\pi_2)$ .

<sup>2</sup> or better an ordering compatible with associativity, such as the AC-RPO [24]

In the following, proofs will be represented by proof terms, the proof ordering  $>$  between them will be the ordering  $>_B$  restricted to proofs with the same conclusion, and the subproof relation  $\triangleright$  will be the subterm relation.

## 4 Standard Completion is an Instance of Abstract Canonical System

### 4.1 Adequacy to the Postulates

Adequacy to postulates A, B, C and D comes from the tree structure of the proof terms representation.

Postulate E is not trivially verified, because of the definition of the ordering as translation of an ordering over proof by replacement. Nevertheless, we can show

**Theorem 4 (Postulate E for Equational Proofs).** *For all contexts  $w[\ ]$ , for all proof terms  $q, r$ ,*

$$q > r \text{ implies } w[q] > w[r] .$$

### 4.2 Standard Completion is Sound and Adequate

This is shown in [6, Lemma 2.1]: if  $E, R \rightsquigarrow E', R'$ , then  $\xrightarrow[E \cup R]{*}$  and  $\xrightarrow[E' \cup R']{*}$  are the same. To prove this, one has simply to verify it for each inference rule of standard completion.

### 4.3 Standard Completion is Good

This is shown in [6, Lemma 2.5, 2.6]: if  $E, R \rightsquigarrow E', R'$ , then proofs in  $E, R$  can be transformed to proofs in  $E', R'$  using following rules:

$$\begin{array}{lcl}
s \xrightarrow[E]{\leftarrow} t & \rightsquigarrow & s \xrightarrow[R']{\rightarrow} t \\
s \xrightarrow[E]{\leftarrow} t & \rightsquigarrow & s \xrightarrow[R']{\rightarrow} u \xrightarrow[E']{\leftarrow} t \\
s \xrightarrow[E]{\leftarrow} s & \rightsquigarrow & s \\
s \xleftarrow[R]{\leftarrow} u \xrightarrow[R]{\rightarrow} t & \rightsquigarrow & s \xleftarrow[E']{\leftarrow} t \\
s \xleftarrow[R]{\leftarrow} u \xrightarrow[R]{\rightarrow} t & \rightsquigarrow & s \xrightarrow[R']{*} v \xleftarrow[R']{*} t \\
s \xrightarrow[R]{\rightarrow} t & \rightsquigarrow & s \xrightarrow[R']{\rightarrow} v \xleftarrow[R']{\leftarrow} t \\
s \xrightarrow[R]{\rightarrow} t & \rightsquigarrow & s \xrightarrow[R']{\rightarrow} v \xleftarrow[E']{\leftarrow} t
\end{array}$$

We have  $\xrightarrow[\rightsquigarrow]{\rightarrow} \subseteq >$ , so these proofs become indeed better.

#### 4.4 Standard Completion is Canonical

We can now show the following theorem:

**Theorem 5 (Completeness of Standard Completion).** *Standard completion results—at the limit, when it terminates without failure—in the canonical, Church-Rosser basis.*

*Proof.* We can show  $R_\infty = E_0^\sharp$ , and because standard completion is good we can use Theorem 2.

*Remark 1.* When standard completion does not terminate, we can show that  $E_0^\sharp = R_\infty^\sharp \subseteq R_\infty$ . Consequently, the resulting set  $R_\infty$  is then *saturated*, but it is not necessarily *reduced*.

This shows that the standard completion is an instance of the framework of the abstract canonical systems, when we choose the convenient proof representation.

### 5 Conclusion

We presented a proof of the adequacy of standard completion to the theory of abstract canonical systems. This conducted us to think about the relations between equational proof representations. The first one, proof terms as presented in [19], is convenient to consider proofs as terms, with a subterm relation and substitutions. The other one, initiated in [6], is fully adapted to the study of the completeness of the standard completion procedure. We presented a way to pass from one representation to another by the mean of the proof term rewrite rules presented in Fig. 4. Thanks to this, we extended the ordering introduced with the proof by replacement to the proof terms and thus combine the advantages of both representations. This therefore answer to the question whether the abstract canonical systems, centered in a quite general way around the notion of proof ordering, are indeed the right framework to uniformly prove the completeness of completion.

We plan now to understand how the proof we have presented here can be extended to other completion procedures. Bachmair introduced another proof ordering to prove the completeness of the completion modulo [6], so that the generalization seems rather natural. We plan also to look at other kinds of deduction mechanisms, such as Buchberger’s algorithm or resolution.

Furthermore, proof terms as presented by [19] are one of the basis for the  $\rho$ -calculus [<http://rho.loria.fr>]. The link between the completion procedure and the sequent systems mentioned above can probably be found here and be related to Dowek’s work proving that confluent rewrite rules can be linked with **Cut**-free proofs of some sequent systems [25].

*Acknowledgments* This paper benefited greatly from suggestions, discussions and the enthusiasm of Nachum Dershowitz.

## References

1. Dershowitz, N., Kirchner, C.: Abstract saturation-based inference. In Kolaitis, P., ed.: *Proceedings of LICS 2003*, Ottawa, Ontario, iee (2003)
2. Bonacina, M.P., Dershowitz, N.: Abstract Canonical Inference. *ACM Transactions on Computational Logic* (2005)
3. Dershowitz, N.: Canonicity. In Dahn, I., Vigneron, L., eds.: *Fourth International Workshop on First-Order Theorem Proving (FTP)*. Volume 86., *Electronic Notes in Theoretical Computer Science* (2003)
4. Knuth, D.E., Bendix, P.B.: Simple word problems in universal algebras. In Leech, J., ed.: *Computational Problems in Abstract Algebra*. Pergamon Press, Oxford (1970) 263–297
5. Dershowitz, N., Kirchner, C.: Abstract Canonical Presentations. *Theoretical Computer Science* (To appear)
6. Bachmair, L.: Proof methods for equational theories. PhD thesis, University of Illinois, Urbana-Champaign, (Ill., USA) (1987) Revised version, August 1988.
7. Bachmair, L., Dershowitz, N.: Equational inference, canonical proofs, and proof orderings. *Journal of Association for Computing Machinery* **41** (1994) 236–276
8. Hsiang, J., Rusinowitch, M.: Proving refutational completeness of theorem proving strategies: The transfinite semantic tree method. *Journal of the ACM* **38** (1991) 559–587
9. Bachmair, L., Ganzinger, H.: Resolution theorem proving. In Robinson, A., Voronkov, A., eds.: *Handbook of Automated Reasoning*. Volume I. Elsevier Science (2001) 19–99
10. Le Chenadec, P.: Canonical Forms in Finitely Presented Algebras. John Wiley & Sons (1986)
11. Huet, G.: Confluent reductions: Abstract properties and applications to term rewriting systems. *Journal of the ACM* **27** (1980) 797–821 Preliminary version in 18th Symposium on Foundations of Computer Science, IEEE, 1977.
12. Peterson, G., Stickel, M.E.: Complete sets of reductions for some equational theories. *Journal of the ACM* **28** (1981) 233–264
13. Jouannaud, J.P., Kirchner, H.: Completion of a set of rules modulo a set of equations. *SIAM Journal of Computing* **15** (1986) 1155–1194
14. Kapur, D., Musser, D.R.: Proof by consistency. *Artificial Intelligence* **13** (1987) 125–157
15. Lankford, D.S.: Canonical inference. Technical report, Louisiana Tech. University (1975)
16. Burel, G.: Systèmes Canoniques Abstraits : Application à la Dédution Naturelle et à la Complétion. Master’s thesis, Université Denis Diderot – Paris 7 (2005)
17. Buchberger, B.: A critical-pair/completion algorithm for finitely generated ideals in rings. In Börger, E., Hasenjaeger, G., Rödding, D., eds.: *Proceedings of Logic and Machines: Decision problems and Complexity*. Volume 171 of *Lecture Notes in Computer Science*., Springer-Verlag (1983) 137–161
18. Winkler, F.: Knuth-Bendix procedure and Buchberger algorithm - A synthesis. In: *Proceedings of the ACM-SIGSAM 1989 International Symposium on Symbolic and Algebraic Computation*, Portland (Oregon, USA), ACM Press (1989) 55–67
19. Meseguer, J.: Conditional rewriting logic as a unified model of concurrency. *Theoretical Computer Science* **96** (1992) 73–155
20. Huet, G.: A complete proof of correctness of the Knuth–Bendix completion algorithm. *Journal of Computer and System Sciences* **23** (1981) 11–21 Also as: Rapport 25, INRIA, 1980.

21. Baader, F., Nipkow, T.: Term *Rewriting* and *all That*. Cambridge University Press (1998)
22. Kirchner, C., Kirchner, H.: Rewriting, solving, proving. A preliminary version of a book available at [www.loria.fr/~ckirchne/rsp.ps.gz](http://www.loria.fr/~ckirchne/rsp.ps.gz) (1999)
23. Cirstea, H., Kirchner, C.: The rewriting calculus — Part I *and* II. Logic Journal of the Interest Group in Pure and Applied Logics **9** (2001) 427–498
24. Rubio, A., Nieuwenhuis, R.: A total AC-compatible ordering based on RPO. Theoretical Computer Science **142** (1995) 209–227
25. Dowek, G.: Confluence as a cut elimination property. In Nieuwenhuis, R., ed.: RTA. Volume 2706 of Lecture Notes in Computer Science., Springer (2003) 2–13



## A Proofs for Section 3 and 4

### A.1 From Proof Terms to Proof by Replacement

To prove the termination of  $\rightsquigarrow / \sim$ , we need a reduction ordering compatible with the associativity. We consider only associativity here, although most of the existing works use associativity and commutativity. Therefore, we need the following lemma.

**Lemma 1.** *If  $A \subseteq B$  then  $>$  is  $B$ -compatible implies  $>$  is  $A$ -compatible.*

*Proof.* Just notice that  $s' \xleftarrow[A]{*} s > t \xleftarrow[A]{*} t'$  implies  $s' \xleftarrow[B]{*} s > t \xleftarrow[B]{*} t'$ .

We can therefore use the  $AC$ -RPO ordering: a total  $AC$ -compatible simplification ordering on ground terms is defined in [24], as an extension of the RPO. To compare terms, they are interpreted using flattening and interpretation rules. As we consider here that the associative commutative symbols have the lowest precedence, we do not need the interpretation rules, and we will only present the flattening rules: terms are reduced using a set of rules

$$f(x_1, \dots, x_n, f(y_1, \dots, y_r), z_1, \dots, z_m) \rightarrow f(x_1, \dots, x_n, y_1, \dots, y_r, z_1, \dots, z_m) \quad (1)$$

for all  $AC$ -symbols  $f$  with  $n + m \geq 1$  and  $r \geq 2$ . Such a rewrite system is terminating as shown in [24].

For all terms  $t$ , let  $snf(t)$  denote the *set of normal forms* of  $t$  using rules (1).

Given a precedence  $>$  on function symbols, let  $>_{rpo}$  denote the recursive path ordering with precedence  $>$  where  $AC$  function symbols have multiset status and other symbols have lexicographic status.

If  $f(s_1, \dots, s_n)$  is the normal form of a term  $s$  rewriting by (1) only at topmost position, then  $tf(s) \stackrel{!}{=} (s_1, \dots, s_n)$ .

**Definition 2 ( $AC$ -RPO).** *For all terms  $s, t$ ,  $s >_{AC-rpo} t$  if:*

- $\forall t' \in snf(t) \exists s' \in snf(s), s' >_{AC-rpo} t'$  **or**
- $\forall t' \in snf(t) \exists s' \in snf(s), s' \geq_{rpo} t'$  **and**  $tf(s) = f(s_1, \dots, s_m)$  **and**  $tf(t) = (t_1, \dots, t_n)$  **and**
  - if the head of  $s$  is  $AC$  then  $\{s_1, \dots, s_m\} >_{AC-rpo\_mult} \{t_1, \dots, t_n\}$  **or**
  - if the head of  $s$  is not  $AC$  then  $(s_1, \dots, s_m) >_{AC-rpo\_lex} (t_1, \dots, t_n)$ .

**Proposition 4 ([24]).** *The  $AC$ -RPO is an  $AC$ -compatible simplification ordering which is total for non  $AC$ -equivalent ground terms.*

We define a precedence  $>$  such that for all function symbols  $f$  and for all rule labels  $\ell$  we have  $\ell > f > \cdot^{-1} > \cdot$ . The  $AC$ -RPO built with this precedence will be noted  $\succ$ .

To show termination, we also need the following lemma:

**Lemma 2.** *For all proof terms  $\pi : t \rightarrow t'$ , we have  $\pi \succeq t$  and  $\pi \succeq t'$ .*

*Proof.* By induction on the structure of the proof term  $\pi$ .

For **Reflexivity**,  $\pi = t = t'$ .

For **Congruence**,  $\pi = f(\pi_1, \dots, \pi_n)$ ,  $t = f(t_1, \dots, t_n)$  and  $t' = f(t'_1, \dots, t'_n)$ . By induction hypothesis, for all  $i \in \{1, \dots, n\}$ , we have  $\pi_i \succeq t_i, t'_i$ . Furthermore,  $\pi$  is not reducible on the top position using rules (1), so that  $snf(\pi) = \{f(\pi'_1, \dots, \pi'_n) : \forall i, \pi'_i \in snf(\pi_i)\}$ , whereas  $t$  and  $t'$  are not reducible. Consequently, by definition of an  $AC$ -RPO,  $\pi \succeq t, t'$ .

For **Replacement**,  $\pi = \ell(\pi_1, \dots, \pi_n)$ ,  $t = g(t_1, \dots, t_n)$  and  $t' = d(t'_1, \dots, t'_n)$  where  $\ell = (g, d) \in E \cup R$ . With the same arguments than for **Congruence**, we can conclude that  $\pi \succeq t, t'$  (recall that  $\ell > g, d$ ).

For **Transitivity**,  $\pi = \pi_1; \pi_2$  where  $\pi_1 : t \rightarrow t''$  and  $\pi_2 : t'' \rightarrow t'$ . By induction hypothesis,  $\pi_1 \succeq t$  and  $\pi_2 \succeq t'$ . As  $\succ$  is a simplification ordering,  $\pi \succ \pi_1, \pi_2 \succeq t, t'$ .

For **Symmetry**,  $\pi = \pi'^{-1}$  where  $\pi' : t' \rightarrow t$ . By induction hypothesis and because  $\succ$  is a simplification ordering,  $\pi \succ \pi' \succeq t', t$ .

**Proposition 5 (Termination).** *The rewrite system  $\rightsquigarrow$  of Fig. 4 modulo  $\sim$  is terminating for ground proof terms.*

*Proof.* We can show that  $\rightsquigarrow \subseteq \succ$ , thus proving the termination of  $\rightsquigarrow / \sim$ :

For **Delete Useless Identities**, it comes from the fact that  $\succ$  is a simplification ordering.

For **Sequentialization**, rules (1) are not applicable on the left side whereas they lead on the right side to  $(f(\pi_1, t_2, \dots, t_n), f(t'_1, \pi_2, \dots, t_n), \dots, f(t'_1, t'_2, \dots, \pi_n))$ . We have  $f > \cdot$ , thus by definition of a RPO, we must then prove that for all  $i \in \{1, \dots, n\}$  we have  $f(\pi_1, \dots, \pi_n) \succ_{RPO} f(t'_1, \dots, t'_{i-1}, \pi_i, t_{i+1}, \dots, t_n)$ , i.e.  $(\pi_1, \dots, \pi_n) \succ_{RPO}^{lex} (t'_1, \dots, t'_{i-1}, \pi_i, t_{i+1}, \dots, t_n)$ . By hypothesis there exists at least a  $j \in \{1, \dots, n\} \setminus \{i\}$  such that  $\pi_j \neq t_j$ , so we can conclude with the preceding lemma.

For **Composition Shallowing**, both sides are not reducible using rules (1). We have  $f >$ ; thus we have to show:  $f(t_1, \dots, \pi_i; \pi'_i, \dots, t_n) \succ_{RPO} f(t_1, \dots, \pi_i, \dots, t_n)$  and  $f(t_1, \dots, \pi_i; \pi'_i, \dots, t_n) \succ_{RPO} f(t_1, \dots, \pi'_i, \dots, t_n)$ . Both comparisons hold by definition of a RPO.

For **Parallel Moves**, both sides are not reducible using rules (1). We have  $\ell >$ ; thus we have to prove that  $\ell(\pi_1, \dots, \pi_n) \succ_{RPO} \ell(t_1, \dots, t_n)$  and  $\ell(\pi_1, \dots, \pi_n) \succ_{RPO} d(\pi_1, \dots, \pi_n)$ . The first comparison holds because of the lemma and because there exists a  $i \in \{1, \dots, n\}$  such that  $\pi_i \neq t_i$ ; the second one holds because  $\ell > d$ .

For **Delete Useless Inverses**, this comes from the fact that  $\succ$  is a simplification ordering.

For **Inverse Congruence**, both sides are not reducible using rules (1), therefore this is a consequence of  $f > \cdot^{-1}$ .

For **Inverse Composition**, both sides are not reducible using rules (1), therefore this is a consequence of  $\cdot^{-1} >$ .

We can also prove confluence:

**Proposition 6 (Confluence).** *The rewrite system  $\rightsquigarrow$  is confluent modulo  $\sim$  on ground proof terms.*

*Proof.* The class rewrite system is linear and terminating, so we just have to check that the critical pairs are confluent [11].

For  $\xleftarrow[R]{\circ} \xrightarrow[R]{\circ}$ , it is easy to check for most of the critical pairs that they are confluent. We only detail the most problematic one. For two possible applications of **Sequentialization**, we have for instance  $f(g(\nu_1, \dots, \nu_m), \pi_1, \dots, \pi_n)$  that can be rewritten to  $f(g(\nu_1, \dots, \nu_m), t_1, \dots, t_n); f(g(s_1, \dots, s_m), \pi_1, \dots, t_n); \dots; f(g(s_1, \dots, s_m), t'_1, \dots, \pi_n)$  and to  $f(g(\nu_1, \dots, s_m); \dots; g(s'_1, \dots, \nu_m), \pi_1, \dots, \pi_n)$ . Both of them reduce to  $f(g(\nu_1, \dots, s_m); \dots; g(s'_1, \dots, \nu_m), t_1, \dots, t_n); f(g(s_1, \dots, s_m), \pi_1, \dots, t_n); \dots; f(g(s_1, \dots, s_m), t'_1, \dots, \pi_n)$ .

For  $\xleftarrow[R]{\circ} \xrightarrow[A]{\circ}$ , the only rules that can interfere with  $\sim$  are **Delete Useless Identities**, **Composition Shallowing** and **Inverse Composition**. We can check that all critical pairs are confluent.

**Theorem 6 (Correspondence between Proof Representations).** *The normal form of a proof term  $\pi$  for the rewrite system  $\rightsquigarrow$ , noted  $nf(\pi)nf(\pi)$ —textbf, has the following form:*

*there exists  $n \in \mathbb{N}$ , some contexts  $w_1[], \dots, w_n[]$ , some indices  $i_1, \dots, i_n \in \{-1, 1\}$ , some rule labels  $\ell_1, \dots, \ell_n$  and some terms  $t_1^1, \dots, t_{m_1}^1, \dots, t_1^n, \dots, t_{m_n}^n$  such that*

$$nf(\pi) = (w_1[\ell_1(t_1^1, \dots, t_{m_1}^1)])^{i_1}; \dots; (w_n[\ell_n(t_1^n, \dots, t_{m_n}^n)])^{i_n}$$

where  $\nu^1$  is a notation for  $\nu$ .

We will denote by  $nf(\pi)$  the normal form of a proof term  $\pi$ .

Such a proof term correspond with the following proof by replacement of equal by equal:

$$w_1[g_1(t_1^1, \dots, t_{m_1}^1)] \xleftrightarrow[\ell_1]{p_1} w_1[d_1(t_1^1, \dots, t_{m_1}^1)] \xleftrightarrow[\ell_2]{p_2} \dots w_n[g_n(t_1^n, \dots, t_{m_n}^n)] \xleftrightarrow[\ell_n]{p_n} w_n[d_n(t_1^n, \dots, t_{m_n}^n)]$$

where for all  $j \in \{1, \dots, n\}$  we have:

- $\ell_j = (g_j, d_j)$ ,
- $p_j$  is the position of  $[]$  in  $w_j[]$ ,  
 $\longrightarrow$  if  $i_j = 1$  and  $\ell_j \in R$ ,
- $\xleftrightarrow[\ell_j]{p_j} = \xleftarrow{\quad}$  if  $i_j = -1$  and  $\ell_j \in R$ ,  
 $\longleftrightarrow$  if  $\ell_j \in E$ .
- if  $j \neq n$ ,  $w_j[d_j(t_1^j, \dots, t_{m_j}^j)] = w_{j+1}[g_{j+1}(t_1^{j+1}, \dots, t_{m_{j+1}}^{j+1})]$ .

*Proof.* We first have to check that proof terms in that form are indeed irreducible by  $\rightsquigarrow$ , what is left to the reader.

Then, suppose that we have an irreducible proof term. Because **Sequentialization** can not be applied, there is at most one ; under all function symbols. Because **Composition Shallowing** can not be applied, there are no ; under all function symbols. Because **Inverse Congruence** and **Inverse Composition** can not be applied,  $\cdot^{-1}$  is applied between ; and function symbols. Irreducible proof term are therefore application of ; over eventually  $\cdot^{-1}$  over base terms composed of function symbols and rule labels.

Because **Delete Useless Identities** and **Delete Useless Inverse** can not be applied, there is a least one non-trivial proof (i.e a proof with a label in it) in each of these base terms. Because **Sequentialization** can not be applied, there is at most one non-trivial proof in each of them. Because **Parallel Moves** can not be applied, the subterms of the labels are  $\Sigma$ -terms. Consequently, each base term contains one and only one rule label, applied to  $\Sigma$ -terms.

## A.2 Adequacy to the Postulates

**Postulate A** The proof of  $(u, v) \in E \cup R$  labeled by  $\ell$  is  $\ell(x_1, \dots, x_n)$  where  $x_1, \dots, x_n$  are the free variables of  $(u, v)$ .

**Postulate B** We can replace the assumption  $\ell(\pi_1, \dots, \pi_n)$  of something proved by its proof where the free variables are replaced by the proofs  $\pi_1, \dots, \pi_n$ .

**Postulate C and D** These postulates hold because of the tree structure of proofs.

**Postulate E** This one does not trivially hold.

We first show the following lemma:

**Lemma 3.** For all function symbols  $f$  of arity  $n + 1$ , for all proof terms  $\pi_1, \dots, \pi_n$ ,  $q$  and  $r$ ,

$$q > r \text{ implies } f(\pi_1, \dots, q, \dots, \pi_n) > f(\pi_1, \dots, r, \dots, \pi_n)$$

*Proof.* Suppose  $q > r$ , thus by definition  $nf(q) >_{rep} nf(r)$ . To compare  $f(\pi_1, \dots, q, \dots, \pi_n)$  and  $f(\pi_1, \dots, r, \dots, \pi_n)$ , we have to transform them to proof by replacement. As  $\xrightarrow{\sim/\sim}$  is Church-Rosser, the way it is applied does not matter.

We have

$$\begin{aligned} f(\pi_1, \dots, q, \dots, \pi_n) &\xrightarrow{\sim}^* f(\pi_1, t_2, \dots, t_n); \dots; f(t'_1, \dots, q, \dots, t_n); \dots; f(t'_1, \dots, \pi_n) \\ &\xrightarrow{\sim}^* f(\pi_1, t_2, \dots, t_n); \dots; \underline{f(t'_1, \dots, nf(q), \dots, t_n)}; \dots; f(t'_1, \dots, \pi_n) \end{aligned}$$

Then, if  $nf(q)$  contains ; the underlined term will be split by **Composition Shallowing**. If it contains  $^{-1}$  the rule **Inverse Congruence** will be applied. Some terms outside the underline corresponding to identity will be removed by **Delete Useless Identities**, and the normal form will look like

$$f(\pi_1, t_2, \dots, t_n); \dots; \underline{f(t'_1, \dots, q_1, \dots, t_n)^{i_1}; \dots; f(t'_1, \dots, q_m, \dots, t_n)^{i_m}}; \dots; f(t'_1, \dots, \pi_n)$$

with  $nf(q) = q_1^{i_1}; \dots; q_m^{i_m}$ .

The same will apply with  $r$ , and therefore, to compare the initial proofs, we just have to compare the costs of the underlined terms.

The cost of  $nf(q)$  will look like  $\{(\{s_1\}, u_1, h_1), \dots, (\{s_m\}, u_m, h_m)\}$ . Then the cost of  $f(t'_1, \dots, q_1, \dots, t_n)^{i_1}; \dots; f(t'_1, \dots, q_m, \dots, t_n)^{i_m}$  will be

$$\{(\{f(t'_1, \dots, s_1, \dots, t_n)\}, u_1, f(t'_1, \dots, h_1, \dots, t_n)), \dots, (\{f(t'_1, \dots, s_m, \dots, t_n)\}, u_m, f(t'_1, \dots, h_m, \dots, t_n))\}$$

For  $nf(r)$  they will be respectively  $\{(\{g_1\}, v_1, d_1), \dots, (\{g_p\}, v_p, d_p)\}$  and

$$\{(\{f(t'_1, \dots, g_1, \dots, t_n)\}, v_1, f(t'_1, \dots, d_1, \dots, t_n)), \dots, (\{f(t'_1, \dots, g_p, \dots, t_n)\}, v_p, f(t'_1, \dots, d_p, \dots, t_n))\}$$

$\gg$ , which is used to compare the first and the third components of each part of the cost, is a reduction ordering, so that  $nf(q) >_{rep} nf(r)$  implies for instance  $f(t'_1, \dots, q_1, \dots, t_n)^{i_1}; \dots; f(t'_1, \dots, q_m, \dots, t_n)^{i_m} >_{rep} f(t'_1, \dots, r_1, \dots, t_n)^{i_1}; \dots; f(t'_1, \dots, r_p, \dots, t_n)^{i_p}$ .

The same is true for labels:

**Lemma 4.** For all rule labels  $\ell$ , for all proof terms  $\pi_1, \dots, \pi_n$ ,  $q$  and  $r$ ,

$$q > r \text{ implies } \ell(\pi_1, \dots, q, \dots, \pi_n) > \ell(\pi_1, \dots, r, \dots, \pi_n)$$

*Proof.*  $\ell(\pi_1, \dots, q, \dots, \pi_n)$  and  $\ell(\pi_1, \dots, r, \dots, \pi_n)$  can be reduced by **Parallel Moves** to  $\ell(t_1, \dots, t_n); d(\pi_1, \dots, q, \dots, \pi_n)$  and  $\ell(t_1, \dots, t_n); d(\pi_1, \dots, r, \dots, \pi_n)$ . We can therefore conclude using the preceding lemma.

This allow us to show

**Theorem 7 (Postulate E for Equational Proofs).** For all proof terms  $p, r$ , for all position  $i$  of  $p$ ,

$$p|_i > r \text{ implies } p > p[r]_i$$

*Proof.* This is proved by induction on  $i$ . For  $i = \epsilon$  this is trivial. For  $i \neq \epsilon$ , by induction hypothesis, the result holds for the subproofs of  $p$ . Consider the head of  $p$ :

- for **Symmetry**, it is trivial;
- for **Transitivity**, it comes from the fact that equational proofs are compared as the multiset of their equational proof steps;
- for **Congruence**, it comes from lemma 3;
- for **Replacement**, it comes from lemma 4.

### A.3 Standard Completion is Canonical

Remember that by fairness assumption,  $E_\infty = \emptyset$ .

**Lemma 5.** *For all standard completion derivations  $(E_i, R_i)_i$ ,*

$$E_0^\sharp \subseteq R_\infty$$

*Proof.* By contradiction, suppose there is  $(a, b) \in E_0^\sharp \setminus R_\infty$ , labeled  $\ell$ . Because completion is adequate, there exists  $p \in \mu Pf(R_\infty)$  proving  $a = b$ . Because  $a = b \in E_0^\sharp$ ,  $\ell(x_1, \dots, x_n) \in Nf(E_0) = Nf(R_\infty)$  where  $(x_i)_i$  are the free variables of  $\ell$ , so that

$$p > \ell(x_1, \dots, x_n)$$

- If there are no peak in  $nf(p)$ , then  $nf(p)$  is a valley proof, and it is easy to show that it is smaller than  $\ell(x_1, \dots, x_n)$ , which is a contradiction with the preceding comparison.
- If there is a parallel peak, for instance  $s[c, e] \xleftarrow[\ell_1]{i} s[d, e] \xrightarrow[\ell_2]{j} s[d, f]$ , then the proof by replacement where this peak is replaced by  $s[c, e] \xrightarrow[\ell_2]{j} s[c, f] \xleftarrow[\ell_1]{i} s[d, f]$  is smaller, thus leading to a contradiction with the minimality of  $p$  in  $Pf(R_\infty)$ .
- If there is a critical peak, then by fairness assumption there is some step  $k$  where this critical peak is treated by **Deduce**. The proof of the conclusion of the critical peak at the step  $k + 1$  is therefore smaller. Because standard completion is good, it can only go smaller, so that at the limit we can find by replacement of the critical peak by this proof a smaller proof of  $a = b$ , thus leading to a contradiction with the minimality of  $p$  in  $Pf(R_\infty)$ .

**Lemma 6.** *For all standard completion derivations  $(E_i, R_i)_i$  which terminate without failure,*

$$R_\infty \subseteq E_0^\sharp$$

*Proof.* By contradiction, suppose there is  $(a, b) \in R_\infty \setminus E_0^\sharp$ , labeled by  $\ell$ . Then there exists a proof  $p \in \mu Pf(E_0^\sharp)$  such that  $\ell(x_1, \dots, x_n) > p$  where  $x_1, \dots, x_n$  are the free variables of  $\ell$ .

Rules comes from orientation of equational axioms through **Orient**, so that  $a \gg b$ . The cost of  $\ell(x_1, \dots, x_n)$  is then  $\{(\{a\}, a, b)\}$ . Consider the leftmost step of  $nf(p)$ . It is of the form  $a \xrightarrow[(c,d)]{i} a[d]_i$  where  $c = a|_i$ . If it is  $a \xleftarrow[d \rightarrow c]{i} a[d]_i$  then the cost of this proof step would be  $\{(\{a[d]_i\}, d, a)\}$ , which is then greater than  $\{(\{a\}, a, b)\}$ , thus leading to a contradiction with the fact that  $\ell(x_1, \dots, x_n) > p$ . If  $a \xrightarrow[c=d]{i} a[d]_i$  then the cost of this proof step would be  $\{(\{a, a[d]_i\}, c, a[d]_i)\}$ , which is then greater than  $\{(\{a\}, a, b)\}$ , thus leading to a contradiction with the fact that  $\ell(x_1, \dots, x_n) > p$ . If it is  $a \xrightarrow[c \rightarrow d]{i} a[d]_i$  then there is a critical pair  $(b, a[d]_i)$  in  $R_\infty$  (we just proved that  $E_0^\sharp \subseteq R_\infty$ ). The fairness assumption will therefore apply, and therefore **Deduce** will produce the equational axiom  $b = a[d]_i$ , which will be oriented, and  $a \longrightarrow b \in R_\infty$  will be simplified through **Compose** or **Collapse**. Because  $a \longrightarrow b$  is persisting, it must be generated once again, thus contradicting the termination of the completion.

**Theorem 8 (Completeness of Standard Completion).** *Standard completion results — at the limit, when it terminates without failure — in the canonical, Church-Rosser basis.*

*Proof.* There is nothing more to prove, because we have  $R_\infty = E_0^\sharp$ , and standard completion is good so we can use Theorem 2.