



The complexity of class polynomial computation via floating point approximations

Andreas Enge

► To cite this version:

Andreas Enge. The complexity of class polynomial computation via floating point approximations. 2007. inria-00001040v2

HAL Id: inria-00001040

<https://inria.hal.science/inria-00001040v2>

Submitted on 24 Apr 2007 (v2), last revised 25 Jul 2008 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The complexity of class polynomial computation via floating point approximations

Andreas Enge

INRIA Futurs & Laboratoire d'Informatique (CNRS/UMR 7161)

École polytechnique, 91128 Palaiseau Cedex, France

enge@lix.polytechnique.fr

April 23, 2006

Abstract

We analyse the complexity of computing class polynomials, that are an important ingredient for CM constructions of elliptic curves, via complex floating point approximations of their roots. The heart of the algorithm is the evaluation of modular functions in several arguments. The fastest one of the presented approaches uses a technique devised by Dupont to evaluate modular functions by Newton iterations on an expression involving the arithmetic-geometric mean. It runs in time

$$O(|D| \log^5 |D| \log \log |D|) = O(|D|^{1+\varepsilon}) = O(h^{2+\varepsilon})$$

for any $\varepsilon > 0$, where D is the CM discriminant and h is the degree of the class polynomial. Another fast algorithm uses multipoint evaluation techniques known from symbolic computation; its asymptotic complexity is worse by a factor of $\log |D|$. Up to logarithmic factors, this running time matches the size of the constructed polynomials. The estimate also relies on a new result concerning the complexity of enumerating the class group of an imaginary quadratic order and on a rigorously proven upper bound for the height of class polynomials.

1 Motivation and results

The theory of complex multiplication yields an efficient approach to the construction of elliptic curves over a finite field having a given endomorphism ring of discriminant D , as long as $|D|$ is not too large. Exploiting the link between the endomorphism ring of an elliptic curve and its number of points, it is possible to efficiently obtain curves with specific properties. Applications include primality proving [AM93], the construction of classical, discrete logarithm based elliptic curve cryptosystems and of identity based cryptosystems [MNT01, DEM05, BLS03, BW05].

The classical approach to effective complex multiplication is to compute minimal polynomials of special, algebraic values of modular functions on the upper complex halfplane (more details are provided in Section 2). The complexity of this algorithm remains a shady issue; one of the reasons a serious analysis has not been undertaken so far are the purported numerical difficulties with the computations with complex floating point numbers, an issue that actually does not arise in practice (cf. the short discussion at the end of Section 7). Moreover, figures found in the existing literature need not necessarily reflect reality. For instance, in [ALV04] it is claimed in a reference to [LL90] that the floating point approach has a complexity of $O(|D|^2 \log^2 |D|)$, while even the most naïve algorithm presented in Section 6.1 in fact has a complexity of only $O(|D|^{3/2+\varepsilon})$. The authors of [CNST98] even pretend that the complexity is exponential in the class number h and thus in $|D|$.

In [CH02] the authors present an algebraic generalisation of the complex multiplication algorithms to p -adic fields, that yields the same minimal polynomials. Besides ruling out any possible numerical instabilities, they obtain a complexity of $O(|D|^{1+\varepsilon})$, which is better than a straightforward implementation of the complex approach, and asymptotically optimal (up to logarithmic factors) due to the size of the constructed objects.

The present article thus has two goals. First, it provides an accurate account of the complexity of different algorithms for class polynomial computation via floating point approximations. Second, it shows that asymptotically optimal algorithms exist also in the complex setting, with a complexity that is linear (up to logarithmic factors) in the output size. These new algorithms are presented in Sections 6.3 and 6.4, and the faster one achieves the following:

2000 Mathematics subject classification: primary 11Y16, secondary 11G15

Theorem 1

Let f be a fixed modular function that is a class invariant for a family of discriminants D of class numbers $h = h(D)$. Then the algorithm of Section 6.4, which computes a floating point approximation to the class polynomial for f , runs in time

$$O(hn \log^2 n \log \log n)$$

when executed with complex floating point numbers of $n = n(D)$ bits precision.

The floating point precision n required to carry out the computations is clearly bounded from below by the height of the class polynomial. The following theorem provides a rigorously proven upper bound on the heights, that is close to experimental findings. A bound of the same shape is given in [ALV04, §3.3] as a heuristic, and the gist of the proof appears already in [LL90, §5.10] and [Sch91, Section 2].

Theorem 2

The logarithmic height of the class polynomial for j for the discriminant D of class number h is bounded above by

$$\begin{aligned} & c_5 h + c_1 N \left(\log^2 N + 4\gamma \log N + \frac{\log N + \gamma + 1}{N} + c_6 \right) \\ & \leq c_1 N \log^2 N + c_2 N \log N + c_3 N + c_1 \log N + c_4 \end{aligned}$$

where $N = \sqrt{\frac{|D|}{3}}$, $\gamma = 0.577 \dots$ is Euler's constant, $c_1 = \sqrt{3}\pi = 5.441 \dots$, $c_2 = 17.512 \dots$, $c_3 = 15.424 \dots$, $c_4 = 11.056 \dots$, $c_5 = 2.474 \dots$ and $c_6 = 2.309 \dots$.

The asymptotic upper bound of

$$O\left(\sqrt{|D|} \log^2 |D|\right)$$

holds for any other class invariant as well.

In practice, one observes that rounding errors do not disturb the result. It suffices to take n as an approximation of the height plus a few guard digits to be able to round the floating point approximation of the class polynomial to the correct polynomial with integral coefficients. A rigorous error analysis, however, appears to be out of reach. So Theorems 1 and 2 can be brought together only in the form of a heuristic, assuming that the computations with floating point numbers of n bits yield an approximation of the class polynomial that is correct on essentially n bits.

Corollary 3 (heuristic)

Taking $n = O\left(\sqrt{|D|} \log^2 |D|\right)$ in Theorem 1 and using the bound on the class number $h = O\left(\sqrt{|D|} \log |D|\right)$ proved at the end of Section 4, the algorithm of Section 6.4 computes the class polynomial for f in time

$$O(|D| \log^5 |D| \log \log |D|) = O(|D|^{1+\varepsilon}) = O(h^{2+\varepsilon})$$

for any $\varepsilon > 0$.

Notice that up to logarithmic factors, this complexity corresponds to the output of the algorithm, namely the size of the class polynomials. Notice also that the correctness of the output can be checked by a probabilistic, Monte-Carlo type algorithm; namely one may check that the reduction of the class polynomial modulo sufficiently many suitable primes p yields elliptic curves over \mathbb{F}_p with complex multiplication by \mathcal{O}_D . Indeed, the main application of class polynomials is to compute elliptic curves over finite fields with given complex multiplication, and in this situation it can be verified independently that the curves are correct.

As an ingredient for the proof of Theorem 1 we obtain in Section 5 the following result for computing class groups of imaginary quadratic orders:

Theorem 4

The class group of the imaginary-quadratic order of discriminant D can be enumerated by a probabilistic algorithm in time

$$O\left(\sqrt{|D|} \log^2 |D| (\log \log |D|)^2 \log \log \log |D|\right).$$

Again the algorithm is essentially optimal in its output size.

2 Complex multiplication and class polynomials

2.1 The basic approach

For proofs of the following facts on complex multiplication of elliptic curves see, for instance, [Cox89].

Let us first consider the situation over the complex numbers. Let $D < 0$ be an imaginary quadratic discriminant, and $\mathcal{O}_D = \left[1, \frac{D+\sqrt{D}}{2}\right]_{\mathbb{Z}}$ the (not necessarily maximal) order of discriminant D in $K = \mathbb{Q}(\sqrt{D})$. The ideal class number of \mathcal{O}_D is denoted by $h = h_D$. By Siegel's theorem [Sie36], $\frac{\log h}{\log |D|} \rightarrow \frac{1}{2}$ ($|D| \rightarrow \infty$), so that $|D| \in O(h^{2+\varepsilon})$ and $h \in O(|D|^{1/2+\varepsilon})$ for any $\varepsilon > 0$. There are h isomorphism classes of elliptic curves over \mathbb{C} having *complex multiplication* by \mathcal{O}_D , that is, curves with \mathcal{O}_D as endomorphism ring. Namely, let $j : \mathbb{H} = \{z \in \mathbb{C} : \Im(z) > 0\} \rightarrow \mathbb{C}$ denote the absolute modular invariant, and let $\tau_i = \frac{-B_i + \sqrt{D}}{2A_i}$ run through the roots of the reduced quadratic forms $[A_i, B_i, C_i] = A_i X^2 + B_i X + C_i$ of discriminant $D = B_i^2 - 4A_i C_i$, representing the ideal classes of \mathcal{O}_D ; then the j -invariants of the elliptic curves are given by the $j(\tau_i)$. Moreover, these $j(\tau_i)$ are algebraic integers; in fact, they generate the so-called *ring class field* K_D for \mathcal{O}_D , the Galois extension of K whose Galois group is isomorphic to the class group of \mathcal{O}_D (the isomorphism being given by the Artin map). The minimal polynomial of the $j(\tau_i)$ over K , $H_D(X) = \prod_{i=1}^h (X - j(\tau_i))$, has in fact coefficients in \mathbb{Z} and is called a *class polynomial*. In the special case that D is a fundamental discriminant, the ring class field K_D is also called the *Hilbert class field* of K .

Let now $\mathbb{F}_q = \mathbb{F}_{p^m}$ be a finite field of characteristic p . Suppose that p splits in $K = \mathbb{Q}(\sqrt{D})$ and that $p \nmid D$; then p is unramified in K_D . If furthermore q may be written as $4q = U^2 + DV^2$ with $U, V \in \mathbb{Z}$, the prime ideals above p in K_D are of inertia degree m . The reductions of the complex elliptic curves with complex multiplication by \mathcal{O}_D modulo any of these prime ideals thus live in \mathbb{F}_q . By Deuring's reduction and lifting theorems [Deu41, Einleitung, par. 5], these h curves are precisely the elliptic curves over \mathbb{F}_q with complex multiplication by \mathcal{O}_D . They may be obtained as follows: Compute the class polynomial $H_D \in \mathbb{Z}[X]$ and reduce it modulo p . It splits completely over \mathbb{F}_q , and each of its roots is the j -invariant of an elliptic curve over \mathbb{F}_q with the desired endomorphism ring.

2.2 Class invariants

Unfortunately, H_D has very large coefficients (see the discussion in Section 4), so that its computation requires a high precision to be accurate. In practice, one may often gain a constant factor for the required number of digits by using instead of j modular functions f that are invariant under $\Gamma^0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{Sl}_2(\mathbb{Z}) : N|b \right\}$ for some positive integer N ; that is, $f\left(\frac{az+b}{cz+d}\right) = f(z)$ for any such matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Under suitable conditions on the discriminant D and suitable normalisations of the τ_i , derived from Shimura's reciprocity law, the *singular values* $f(\tau_i)$ are still elements of the class field K_D [Sch76, Gee99, GS98, Sch02]; we then call f a *class invariant* and the minimal polynomial $H_D[f](X) = \prod_{i=1}^h (X - f(\tau_i))$ again a *class polynomial*. The first such class invariants are given by Weber's functions \mathfrak{f} , \mathfrak{f}_1 , \mathfrak{f}_2 , γ_2 and γ_3 [Web08].

Two parameterised families of class invariants are exhibited in [ES04], where N is the product of two primes and $H_D[f] \in \mathbb{Z}[X]$, and [EM07], where N is prime and $H_D[f]$ has coefficients in the maximal order of $\mathbb{Q}(\sqrt{D})$. Again, these class polynomials split completely after reduction in \mathbb{F}_q . The corresponding elliptic curves may be recovered from some polynomial relationship between f and j : If the modular curve $X_0(N)$ has genus 0, then j is sometimes given by a rational formula in f ; otherwise it has been suggested in [ES04] to look for a root of the modular polynomial $\Phi(f, j)$ after reducing modulo p and specialising in the value found for f in \mathbb{F}_q . (Both cases require that all coefficients be rational to make sense modulo p , which holds for all exhibited class invariants.)

3 Complexity of arithmetics

This section discusses the well-known complexity of the basic multiprecision and polynomial arithmetic underlying the computations. Throughout the remainder of this article we use llog as an abbreviation for $\log \log$, and lllog for $\log \log \log$.

3.1 Multiprecision floating point arithmetic

Let $M(n)$ be the bit complexity of multiplying two n -bit integers; Schönhage–Strassen multiplication implies $M(n) \in O(n \log n \log \log n)$. Then the four basic arithmetic operations and the square roots of real floating point numbers of precision n have a complexity of $O(M(n))$, the inversion and square roots being realised by Newton iterations as explained in Lemmata 2.2 and 2.3 of [Bre76]. The same article shows that \exp , \sin and the constant π can be computed in $O(\log n M(n))$.

Thus, the four basic operations on complex floating point numbers of precision n can be executed in time $O(M(n))$. Letting $c = \sqrt{\frac{a+\sqrt{a^2+b^2}}{2}}$ for $a > 0$, one obtains $\sqrt{a+bi} = c + \frac{b}{2c}i$ in time $O(M(n))$. Finally, the complex exponential may be reduced to the real exponential and the real sine and cosine functions and is thus computed with complexity $O(\log n M(n))$.

3.2 Polynomial arithmetic

Concerning operations with polynomials, we assume that a floating point precision of n bits has been fixed and is the same for all input and output polynomials. Multiplying two polynomials of degree d over \mathbb{C} by the FFT takes $O(d \log d)$ multiplications in \mathbb{C} , whence it has a complexity of $O(d \log d M(n))$ once the necessary roots of unity have been computed.

The following algorithm ([GG99, Algorithm 10.3]) obtains a monic polynomial of degree h from its roots by organising the computations in a binary tree.

Algorithm 5 (Poly_from_roots)

Input: $h \in \mathbb{N}$ and $x_1, \dots, x_h \in \mathbb{C}$

Output: binary tree $T_{k,i} = (X - x_{(i-1)2^k+1}) \cdots (X - x_{\min(i2^k, h)})$
for $k = 0, \dots, t = \lceil \log_2 h \rceil$, $i = 1, \dots, 2^{t-k}$;
in particular, the root $T_{t,1} = (X - x_1) \cdots (X - x_h)$

1. **for** $i = 1$ **to** h
 $T_{0,i} \leftarrow X - x_i$
2. **for** $i = h+1$ **to** 2^t
 $T_{0,i} \leftarrow 1$
3. **for** $k = 1$ **to** t
for $i = 1$ **to** 2^{t-k}
 $T_{k,i} \leftarrow T_{k-1,2i-1} \cdot T_{k-1,2i}$
4. **return** T

If the needed roots of unity are precomputed, the algorithm has a complexity of $O(h \log^2 h M(n))$. Notice that the roots of unity of order $2^{\lceil \log_2 h \rceil}$ suffice to carry out all the FFTs in Algorithm 5. By Section 3.1, computing a primitive root of unity takes $O(\log n M(n))$, all others can be obtained by successive multiplications in $O(h M(n))$. Thus, the total complexity of Algorithm 5 becomes

$$O((h \log^2 h + \log n)M(n)).$$

One of the asymptotically optimal algorithms for class polynomials relies on a related technique of symbolic computation (see [GG99, §10.1]). Let $f(X)$ be a polynomial of degree d over \mathbb{C} that is to be evaluated in h different arguments x_1, \dots, x_h . The key observation for doing so fast is that $f(x_i)$ is nothing but $f(X) \bmod X - x_i$. So the first step of the algorithm is the construction of the binary tree $T_{k,i}$ as in Algorithm 5, containing products of more and more of the $X - x_i$. Then a matching tree $R_{k,i} = f \bmod T_{k,i}$ is computed in the converse order, from its root to its leaves, such that the leaves contain the desired values of f .

Algorithm 6 (Multi_eval)

Input: $h \in \mathbb{N}$, $f \in \mathbb{C}[X]$ and $x_1, \dots, x_h \in \mathbb{C}$

Output: $f(x_1), \dots, f(x_h)$

1. $T \leftarrow \text{Poly_from_roots}(h, x_1, \dots, x_h)$
2. $t \leftarrow \lceil \log_2 h \rceil$, $R_{t,1} \leftarrow f \bmod T_{t,1}$

3. **for** $k = t - 1$ **downto** 0
 for $i = 1$ **to** 2^{t-k}
 $R_{k,i} \leftarrow R_{k+1, \lfloor \frac{i+1}{2} \rfloor} \bmod T_{k,i}$
 4. **return** $R_{0,1}, \dots, R_{0,h}$

A division with remainder of a polynomial of degree d by a polynomial of degree d' has the same complexity as multiplying polynomials of degree bounded by d ([GG99, §9.1]). The algorithm first computes an inverse modulo some power of X by Newton iterations and obtains the quotient with one multiplication; the remainder then requires another multiplication. So after the first reduction of f modulo the product of all the $X - x_i$, the complexity of Algorithm 6 on n -bit numbers is the same as that of Algorithm 5. Including the first reduction, it is given by

$$O((d \log d + h \log^2 h + \log n)M(n)).$$

4 The height of class polynomials

The running time of the algorithms working with floating point approximations depends crucially on the precision n . To be able to round the approximated class polynomial to a polynomial over the integers (\mathbb{Z} or the maximal order of an imaginary-quadratic number field), n has to be at least the bit size of the largest coefficient, or otherwise said, the logarithmic height of the polynomial. In this section, we prove Theorem 2 by developing an explicit upper bound without hidden constants for the height of the class polynomial for j . The result is of independent interest; for instance, it allows to bound the precision also for the p -adic algorithms, which yields a proof of correctness for their output.

As shown in [EM02], the height of class polynomials for a different f changes asymptotically by a constant factor, depending on the degrees in f and j of the modular polynomial connecting the two functions. So this section proves an asymptotic bound of $O(\sqrt{|D|} \log^2 |D|)$ for the height of any class polynomial. To obtain a more explicit bound for invariants other than j , the techniques of this section may be used with the necessary adaptations.

Let us first give the intuitive basis for the bound. Since the values of j are usually large, more often than not the largest coefficient of the class polynomial is its constant one. Then approximating $j(\tau) = q^{-1} + 744 + \sum_{\nu=1}^{\infty} c_{\nu} q^{\nu}$ by the first term, q^{-1} , one obtains a heuristic estimate for the height of

$$\pi \sqrt{|D|} \sum_{[A,B,C]} \frac{1}{A},$$

where the sum is taken over all reduced primitive quadratic forms of discriminant D . This heuristic estimate, that can also be found in [ALV04], is very well confirmed by experimental findings, see [EM02].

Turning this idea into an explicit bound is mainly a matter of computations that are sketched in the following.

The upper bound $\frac{e^{4\pi\sqrt{\nu}}}{\sqrt{2}\nu^{3/4}}$ on c_{ν} of [BP05] yields

$$|j(\tau) - q^{-1}| \leq 744 + \sum_{\nu=1}^{\infty} \frac{e^{4\pi\sqrt{\nu}}}{\sqrt{2}\nu^{3/4}} e^{-\pi\sqrt{3}} = k_1 = 1370.566 \dots$$

for τ in the fundamental domain \mathcal{F} , so that

$$|j(\tau)| \leq |q^{-1}| + k_1 \leq k_2 |q^{-1}|$$

with $k_2 = \frac{k_1}{e^{\pi\sqrt{3}}} = 5.939 \dots$

Let the A_i be numbered in increasing order. Then the logarithm of the absolute value of the coefficient in front of X^j is bounded above

$$\log \left(\binom{h}{j} \prod_{i=1}^j (k_2 |q_i^{-1}|) \right) \leq \log \left(2^h k_2^h \prod_{i=1}^h |q_i^{-1}| \right) \leq \log(2k_2)h + \pi \sqrt{|D|} \sum_{i=1}^h \frac{1}{A_i},$$

which is in fact independent of j .

The next step consists of estimating $\sum_{i=1}^h \frac{1}{A_i}$. [Sch91, Lemma 2.2] proves that $\sum_{i=1}^h \frac{1}{A_i} = O(\log^2 |D|)$. We shall derive a bound that makes the involved constants explicit. Consider the

number of possible B for a given A . This number is certainly bounded above by the number of $B \in]-A, A]$ that satisfy $B = D \pmod{2}$ and $\frac{B^2 - D}{4} = 0 \pmod{A}$. Assuming the worst case that the quadratic equation has a root modulo each prime divisor of A and considering the cases of odd and even A separately, one obtains an upper bound on the number of B of $2 \cdot 2^{\omega(A)} \leq 2\tau(A)$, where $\omega(A)$ denotes the number of prime factors of A and $\tau(A)$ its number of divisors. Hence,

$$\sum_{i=1}^h \frac{1}{A_i} \leq 2 \sum_{A=1}^{\sqrt{\frac{|D|}{3}}} \frac{\tau(A)}{A},$$

and this sum may be bounded by standard techniques from analytic number theory. We use the estimates

$$\log N + \gamma \leq \sum_{n=1}^N \frac{1}{n} \leq \log N + \gamma + \frac{1}{2N} \quad (1)$$

with Euler's constant $\gamma = 0.577 \dots$ and

$$\sum_{n=1}^N \frac{\log n}{n} \geq \frac{\log 2}{2} + \int_3^{N+1} \frac{\log n}{n} \geq \frac{1}{2} \log^2 N - k_3 \quad (2)$$

with $k_3 = \frac{1}{2}(\log^2 3 - \log 2) = 0.256 \dots$. We have

$$\begin{aligned} 2 \sum_{A=1}^N \frac{\tau(A)}{A} &= 2 \sum_{A=1}^N \sum_{1 \leq m, n: mn=A} \frac{1}{mn} = 2 \sum_{1 \leq m, n: mn \leq N} \frac{1}{mn} = 2 \sum_{m=1}^N \frac{1}{m} \sum_{n=1}^{\lfloor N/m \rfloor} \frac{1}{n} \\ &\leq 2 \sum_{m=1}^N \frac{1}{m} \left(\log \frac{N}{m} + \gamma + \frac{1}{2 \lfloor \frac{N}{m} \rfloor} \right) \\ &\leq \log^2 N + 4\gamma \log N + 2\gamma^2 + \frac{\log N + \gamma}{N} + k_3 + \sum_{m=1}^N \frac{1}{m \lfloor \frac{N}{m} \rfloor} \text{ by (1) and (2)} \end{aligned}$$

The last sum can be bounded by

$$\sum_{m=1}^N \frac{1}{m \lfloor \frac{N}{m} \rfloor} \leq \sum_{m=1}^{\lfloor \frac{N}{2} \rfloor} \frac{1}{m (\frac{N}{m} - 1)} + \sum_{m=\lfloor \frac{N}{2} \rfloor + 1}^N \frac{1}{m} \leq 2 \sum_{m=\lfloor \frac{N}{2} \rfloor}^N \frac{1}{m} \leq 2 \log 2 + \frac{1}{N} \text{ by (1)}$$

Combining the inequalities, the height of the class polynomial is bounded from above by

$$k_5 h + \pi \sqrt{|D|} \left(\log^2 N + 4\gamma \log N + \frac{\log N + \gamma + 1}{N} + k_4 \right) \quad (3)$$

with $N = \sqrt{\frac{|D|}{3}}$, $k_4 = k_3 + 2 \log 2 + 2\gamma^2 = 2.309 \dots$ and $k_5 = \log(2k_2) = 2.474 \dots$

Similarly to the argumentation above one shows that

$$h \leq 2 \sum_{A=1}^n \tau(A) \leq 2N(\log N + \gamma) + 1, \quad (4)$$

which also implies $h = O(\sqrt{|D|} \log |D|)$.

Combining (3) and (4) yields the bound of Theorem 2.

5 Class group enumeration

Computing the class group of \mathcal{O}_D by enumerating all reduced primitive quadratic forms of discriminant D is a step of the algorithm that can be neglected in practice. In the new algorithms of Sections 6.3 and 6.4, however, it risks to come close to the complexity of the crucial parts. One has to determine all coprime $[A_i, B_i, C_i]$ such that $|B_i| \leq A_i \leq C_i$, $B_i > 0$ if one of the inequalities is not strict, and $D = B_i^2 - 4A_i C_i$. These conditions imply that $A_i \leq \sqrt{\frac{|D|}{3}}$. The following trivial

algorithm is implemented in the code described in more detail in Section 7: Loop over A and B such that $0 \leq B \leq A \leq \sqrt{\frac{|D|}{3}}$ and compute the corresponding C , if it exists. While it can be seen in the figures of Table 1 that this approach takes virtually no time, it carries out $O(|D|)$ arithmetic operations with integers of bit size in $O(\log |D|)$ and thus has a complexity of

$$O(|D| M(\log |D|)) = O(|D| \log |D| \log |D| \log |D|),$$

which is smaller than the bound of Theorem 1 only by logarithmic factors.

An asymptotically faster (and again up to logarithmic factors optimal) algorithm is inspired by the theoretical considerations of Section 4.

Looping over A in the interval $1 \leq A \leq \sqrt{\frac{|D|}{3}}$, one solves the congruence $B = D \bmod 2$ and $\frac{B^2 - D}{4} = 0 \bmod A$, and keeps the form if $C = \frac{B^2 - D}{4A} \geq A$. As shown in Section 4, the total number of B to consider is on $O(\sqrt{|D|} \log |D|)$.

To find the square root of D modulo A , the prime factorisation of A is required, and the most convenient approach is to enumerate the A in factored form. To this purpose, one needs the primes up to $\sqrt{\frac{|D|}{3}}$, which can be obtained in time $O\left(\frac{\sqrt{|D|} \log |D|}{\log |D|}\right)$ by [AB04]. All possible A are then computed with an amortised cost of one multiplication per value in total time $O(\sqrt{|D|} M(\log |D|))$.

For any given A and prime $p|A$, a square root of D modulo p is computed in $O(\log p M(\log p))$ by Cipolla's algorithm [Cip03], lifting to a root modulo p^e for $p^e || A$ requires an additional $O(e M(\log p))$. For one A , this takes

$$O\left(\sum_{p^e || A} \log(p^e) M(\log p)\right) = O(\log A M(\log A)) = O(\log |D| M(\log |D|)),$$

or $O(\sqrt{|D|} \log |D| M(\log |D|))$ for all A .

The roots modulo prime powers have to be recombined by the Chinese remainder theorem to form a candidate B . Organising the computations in a tree with the leaves indexed by the $p^e || A$, one may use the analogues for numbers of the fast algorithms for polynomials of Section 3.2. The tree has $\omega(A) = O(\log A)$ leaves and thus a height of $O(\log A)$, and the root contains a number of $O(\log A)$ digits, so that one computation of Chinese remainders takes

$$O(\log A M(\log A)).$$

This is to be multiplied by the total number of B , which yields an overall complexity of

$$O(\sqrt{|D|} \log |D| \log |D| M(\log |D|)) = O(\sqrt{|D|} \log^2 |D| \log^2 |D| \log |D|).$$

So Chinese remaindering is the dominant step of the class group enumeration, and Theorem 4 is proved.

When working with a class invariant other than j that is not invariant under $\text{Sl}_2(\mathbb{Z})$, but only under $\Gamma^0(N)$ for some $N > 0$, these representatives of the class group need to be normalised to obtain a coherent set of algebraic conjugates. Such a normalisation is, for instance, given by an N -system as defined in [Sch02]. It has the property that

$$\gcd(A_i, N) = 1 \text{ and } B_i \equiv B_1 \pmod{2N} \text{ for } 1 \leq i \leq h$$

and may be obtained by applying suitable unimodular transformations to the original $[A_i, B_i, C_i]$. The coefficients of these transformations are defined modulo N , so that for fixed N , trying all possibilities requires a constant number of arithmetic operations per form and increases the size of the A_i , B_i and C_i by a constant factor. Thus, the total complexity of this step becomes $O(h M(\log |D|))$, which is negligible compared to the enumeration step. (In practice, one would use a more intelligent approach to lower the complexity with respect to N , cf. the constructive proof of Proposition 3 in [Sch02].)

6 Complexity of class polynomial computation

We are now able to provide the generic complexity for class polynomial computation via floating point approximations. Different approaches to evaluating the class invariants lead to algorithms with different overall complexities; these are examined below.

Let n be the precision in bits used for the computations, let h be the class number of \mathcal{O}_D and denote by $E(h, n)$ the time needed to evaluate the class invariant with precision n in h values. Then the algorithm takes

- $O(h^{3/2+\varepsilon})$ for enumerating the class group according to Section 5;
- $E(h, n)$ for evaluating the class invariant and
- $O((h \log^2 h + \log n)M(n))$ for reconstructing the class polynomial from its roots according to Algorithm 5.

The required precision n is actually larger than h (see Section 4), so that the class group computation indeed becomes negligible since $E(h, n)$ is at least of the order $hn \geq h^2$: This is the time needed to write down the conjugates. The discussion of the following sections shows that the evaluations of the modular function also dominate the construction of the polynomial from its roots, so from now on we examine in more detail the quantity $E(h, n)$.

We propose four different algorithms for evaluating modular functions. The first two of them are well-known, the third one is a novel application of the techniques of symbolic computation presented in Section 3.2 and almost allows to obtain the complexity stated in Theorem 1. The fourth one finally proves Theorem 1 by gaining an additional logarithmic factor.

The class polynomial $H_D[f] = \prod_{i=1}^h (X - f(\tau_i))$ is obtained by evaluating the function f in the h different arguments $\tau_i = \frac{-B_i + \sqrt{D}}{2A_i}$ for an N -system $[A_i, B_i, C_i]$ (see Section 5), where N depends on f and is assumed to be fixed. Since $f(z)$ is modular for $\Gamma^0(N)$, it is invariant under the translation $z \mapsto z + N$ and admits a Fourier transform, that is, a Laurent series expansion in the variable $q^{1/N} = e^{2\pi iz/N}$. Thus, the $f(\tau_i)$ may be obtained by first computing the corresponding $q_i^{1/N}$ and then evaluating the q -expansion in these arguments. By Section 6.3, the q_i can be computed in time $O(h \log n M(n)) = O(hn \log^2 n \log n)$, which will be dominated by the actual function evaluations.

6.1 The naïve approach

The straightforward technique for evaluating $f = \sum_{\nu=\nu_0}^{\nu_1} c_\nu (q^{1/N})^\nu$ consists of a Horner scheme for the polynomial part $\sum_{\nu=\nu_0}^{\nu_1-\nu_0} c_{\nu+\nu_0} (q^{1/N})^\nu$ and a multiplication by $(q^{1/N})^{\nu_0}$. (Notice that usually $\nu_0 < 0$.) Its complexity is $O((\nu_1 - \nu_0 + \log \nu_0)M(n))$, the c_ν having been precomputed to precision n . Actually, ν_0 is a constant depending only on f ; ν_1 , however, depends not only on the c_ν , but also on the desired precision n and on $|q^{1/N}|$ and thus on the function argument.

Consider first the classical case of f being j , which is invariant under $\text{Sl}_2(\mathbb{Z})$. We may then assume that the arguments are transformed by a matrix in $\text{Sl}_2(\mathbb{Z})$ into the standard fundamental domain

$$\mathcal{F} = \left\{ z \in \mathbb{C} : |z| \geq 1, -\frac{1}{2} \leq \Re(z) < \frac{1}{2} \right\}$$

prior to evaluating j . Then $|q| \leq \left| e^{2\pi i \frac{1+\sqrt{-3}}{2}} \right| = e^{-\pi\sqrt{3}}$ is bounded from above by a constant. On the other hand, it is shown in [BP05] that $0 \leq c_\nu \leq \frac{e^{4\pi\sqrt{\nu}}}{\sqrt{2}\nu^{3/4}}$ for $\nu \geq 1$. Thus, it is possible to fix $\nu_1 \in O(n)$ to obtain a precision of $O(n)$ digits.

The total complexity of evaluating j at h values then becomes

$$O(hn M(n)),$$

or

$$O\left(|D|^{3/2} \log^6 |D| \log \log |D|\right)$$

with $n = O(\sqrt{|D|} \log^2 |D|)$ and $h = O(\sqrt{|D|} \log |D|)$ according to Section 4.

Concerning alternative class invariants, it holds unfortunately that the fundamental domain for $\Gamma^0(N)$ with $N > 1$ contains at least one rational number (called *cusp*). In such a cusp, $|q| = 1$, and the q -expansion usually diverges; in a neighbourhood of the cusp, it may converge arbitrarily slowly. In this case, it is possible to use a different expansion in the neighbourhood by transporting the cusp to infinity via a matrix in $\text{Sl}_2(\mathbb{Z})$. We will not pursue this discussion, since the approach of Section 6.2 provides a faster and simpler solution for all class invariants actually suggested in the literature.

6.2 Using the sparsity of η

Virtually all class invariants suggested in the literature are in some way derived from Dedekind's η -function. This is the case for the Weber functions $\mathfrak{f}(z) = e^{-\pi i/24} \frac{\eta(\frac{z+1}{2})}{\eta(z)}$, $\mathfrak{f}_1(z) = \frac{\eta(\frac{z}{2})}{\eta(z)}$ and $\mathfrak{f}_2(z) = \sqrt{2} \frac{\eta(2z)}{\eta(z)}$ already examined in [Web08], the generalised Weber functions $\mathfrak{w}_N(z) = \frac{\eta(\frac{z}{N})}{\eta(z)}$ suggested in [EM07], the double η quotients $\mathfrak{w}_{p_1, p_2}(z) = \frac{\eta(\frac{z}{p_1})\eta(\frac{z}{p_2})}{\eta(z)\eta(\frac{z}{p_1 p_2})}$ proposed in [ES04], and even for j . In fact, j is most conveniently computed as $j = \left(\frac{\mathfrak{f}_1^{24} + 16}{\mathfrak{f}_1^8}\right)^3$.

The definition of η in [Ded76] is closely related to the partition generating function:

$$\eta = q^{1/24} \prod_{\nu \geq 1} (1 - q^\nu);$$

evaluating the product to precision n requires $O(n)$ arithmetic operations. An expression better suited for computation is given by Euler's pentagonal number theorem [Eul83]:

$$\eta = q^{1/24} \left(1 + \sum_{\nu=1}^{\infty} (-1)^\nu \left(q^{\nu(3\nu-1)/2} + q^{\nu(3\nu+1)/2} \right) \right).$$

Since the occurring exponents are values of quadratic polynomials, the series is very sparse: To reach an exponent of order $O(n)$, only $O(\sqrt{n})$ terms need to be computed, and this process can be implemented with $O(\sqrt{n})$ multiplications. Recall that any polynomial of fixed degree can be evaluated in an arithmetic progression with a constant number of arithmetic operations per additional value, once the first few values are known; the employed algorithm relies on iterated differences. In the special case of η , the following recursion yields two additional terms of the series at the expense of four multiplications by recursively computing q^ν , $q^{2\nu-1}$, $q^{\nu(3\nu-1)/2}$ and $q^{\nu(3\nu+1)/2}$ as follows:

$$\begin{aligned} q^{\nu+1} &= q^\nu \cdot q \\ q^{2(\nu+1)-1} &= q^{2\nu-1} \cdot q^2 \\ q^{\nu(3(\nu+1)-1)/2} &= q^{\nu(3\nu+1)/2} \cdot q^{2(\nu+1)-1} \\ q^{\nu(3(\nu+1)+1)/2} &= q^{\nu(3\nu+1)-1)/2} \cdot q^{\nu+1} \end{aligned}$$

Besides the sparse and regular series expression, the η function has a second crucial property that makes it well suited for computation: It is a modular form of weight $1/2$. As such, unlike j , it is not invariant under transformations in $\text{Sl}_2(\mathbb{Z})$. However, its transformation behaviour is explicitly known (cf. [Deu58, §4]) and easily computable. Thus, to obtain $\eta(z)$ for an arbitrary value of z , one should first transform z into the fundamental domain \mathcal{F} , so that the series can be truncated at an exponent of order $O(n)$. Then the evaluation of $O(\sqrt{n})$ terms of the η series for $O(h)$ distinct values (the constant being at most 4 for the \mathfrak{w}_{p_1, p_2} mentioned in the beginning of this section) with a floating point precision of $O(n)$ digits can be carried out in time

$$O(h\sqrt{n} M(n)),$$

or

$$O\left(|D|^{5/4} \log^5 |D| \log \log |D|\right)$$

as $h = O(\sqrt{|D|} \log |D|)$ and $n = O(\sqrt{|D|} \log^2 |D|)$ suffices according to Section 4.

It remains, however, to verify that transforming the arguments into the fundamental domain is dominated by the cost of the series evaluation. The arguments being roots of an N -system $[A_i, B_i, C_i]$ with $A_i, |B_i| \in O(N^2 \sqrt{|D|}) = O(\sqrt{|D|})$, they may be transformed into \mathcal{F} by symbolically reducing the quadratic forms in time $O(h M(\log |D|))$ (see [Coh93, Prop. 5.4.3]), which is negligible. The same holds for arguments such as $\frac{z+1}{2}$ or $\frac{z}{N}$, corresponding to quadratic forms whose discriminants have absolute values in $O(|D|)$.

6.3 Multipoint evaluation

The algorithms of Sections 6.1 and 6.2 compute the values of modular functions one at a time; but for the sake of class polynomial computation, we need the values in many points. For polynomials, Algorithm 6 provides a fast way of doing exactly this. But from a numerical point of view, a

class invariant can be seen as a polynomial via its truncated q -expansion. Either one considers the function directly as done for j in Section 6.1, or one proceeds via η as in Section 6.2. In both cases, a polynomial of degree $O(n)$ has to be evaluated in $O(h)$ points, which by Algorithm 6 can be done in time

$$\begin{aligned} O((n \log n + h \log^2 h)M(n)) &= O(n^2 \log^2 n \log n + nh \log^2 h \log n \log n) \\ &= O(|D| \log^6 |D| \log \log |D|). \end{aligned}$$

6.4 Newton iterations on the arithmetic-geometric mean

A new approach for evaluating modular functions in single arguments is described in [Dup07]. It is based on the arithmetic-geometric mean and Newton iterations on a function involving it. The basic algorithm underlying Theorem 4 of [Dup07] computes the modular function k' , whose square λ satisfies

$$\frac{256(1 - \lambda + \lambda^2)^3}{(\lambda(1 - \lambda))^2} = j.$$

For an argument with imaginary part bounded by a constant and the precision n tending to infinity, it has a complexity of $O(\log nM(n))$.

During class polynomial computations, the precision and the imaginary part of the arguments are tightly coupled, so that this algorithm is not sufficient to derive the desired complexity result. The modification of Theorem 5 of [Dup07] obtains the same complexity of $O(n \log^2 n \log \log n)$ uniformly in the argument. If the imaginary part of the argument is of the order of the required precision, then the algorithm of Section 6.2 or even the naïve algorithm of Section 6.1 already yield the desired result with a constant number of arithmetic operations. Otherwise, the argument is repeatedly divided by 2 until its imaginary part is smaller than a constant, which can be compensated by iterations of the arithmetic-geometric mean. Then the basic algorithm converges sufficiently fast.

For other modular functions f , one may have the evaluation of k' followed by Newton iterations on the modular polynomial relating k' and f . For fixed f , this phase does not increase the complexity. The approach does not work, however, for η , which is a modular form of weight 1/2 instead of a modular function (of weight 0). The algorithm of Section 7 of [Dup07] computes first θ_{00}^2 , a certain modular form of weight 1, as the inverse of the arithmetic-geometric mean of 1 and k' , and then η as the twelfth root of $\lambda(1 - \lambda)\theta_{00}^2/16$ by a suitably initialised Newton process. Again, one obtains a complexity of $O(\log nM(n))$ for an evaluation at precision n , uniformly in the argument.

The complexity for evaluating at h arguments then becomes

$$O(h \log nM(n)) = O(|D| \log^5 |D| \log \log |D|)$$

with the estimates of Section 4 for h and n . This proves Theorem 1.

7 Implementation

The algorithms of this article have been implemented using `gmp` [Gra] with an assembly patch for 64 bit AMD processors [Gau05], `mpfr` [HLPZ] and `mpc` [EZ] for the multiprecision arithmetic and a library of the author's for the polynomial operations. Table 1 provides running times for class numbers between 2500 and 100000, obtained on an AMD Opteron 250 with 2.4 GHz. All timings are given in seconds and rounded to two significant digits. (The computations for class number 100000 have been carried out on a 2.2 GHz machine disposing of more memory, and the running times have been scaled accordingly.) For each class number, the discriminant with smallest absolute value has been chosen. Only the algorithms of Sections 6.2 to 6.4 are taken into account; the naïve approach of Section 6.1 is clearly inferior to the one exploiting the sparsity of η . The chosen class invariant is the double η -quotient $\mathfrak{w}_{3,13}$, and its values are obtained by precomputing a table for the values of η at the h reduced quadratic forms.

The first lines of the table provide some general information. The precision (1) of the floating point computations is obtained by increasing the estimate of Section 4 by 1% to account for potential rounding errors. As the chosen class invariant is not j , a correction factor depending on the class invariant needs to be used, see Section 4. This factor is correct only asymptotically, which explains why the actual height (2) is a bit smaller than the precision estimate. For j , the two are closer to each other. $M(n)$ is measured by computing the first 100000 successive powers of $\pi + i\gamma$ with Euler's constant γ .

		5000	10000	20000	40000	100000
	$ D $	6961631	23512271	98016239	357116231	2093236031
(1)	precision n (bits)	9540	20317	45179	96701	264727
(2)	height (in base 2)	8431	18114	40764	87842	242410
(3)	$M(n)$	7.3	23	75	230	1080
(4)	class group	0.1	0.1	0.4	1.3	6.8
(5)	conjugates from η	3.4	21	140	890	10000
(6)	poly. from roots	13	93	730	5200	120000
sparse series						
(7)	η	12	98	900	7700	140000
(8)	of which q_i	3.0	22	170	1300	20000
(9)	total time	28	210	1800	14000	270000
multipoint eval.						
(10)	η	93	640	5700	42000	aborted
(11)	total time	110	750	6500	48000	
(10) / (7)		7.8	6.5	6.3	5.5	—
AGM						
(12)	η	32	200	1400	9900	130000
(13)	total time	48	320	2300	16000	260000
(12) / (7)		2.7	2.0	1.6	1.3	0.93

Table 1: Running times

The second block of lines provides timings for the steps that are independent of the algorithm used for evaluating the modular function. As can be seen, the class group computation (4) is completely negligible. Line (5) corresponds to the effort of deriving all values of the class invariant from the tabulated η values (reduction of quadratic forms, multiplication by 24-th roots of unity and computation of the η quotients). The computation of the polynomial from its roots (6) corresponds precisely to Algorithm 5, and provides a measure for the complexity of the operations with polynomials. For constructing the largest polynomial of degree 100000 from its roots, the polynomial FFT has been disabled during the last steps and replaced by Toom–Cook multiplication, since the FFT consumed too much memory; this explains the jump from $h = 40000$.

The third block contains the timings for evaluating η in the reduced quadratic forms (7) using the sparse series representation as described in Section 6.2, and the total running time for computing the class polynomials using this technique (9). Line (8) details the time spent in (7) (and also in (10)) for computing the q_i ; it essentially measures the complex exponential.

The fourth block represents the corresponding results for the multipoint evaluation approach of Section 6.3, and the last block corresponds to the asymptotically optimal evaluation of Section 6.4, for which an implementation by Dupont has been used. As explained in Section 6.4, the algorithm requires to switch to the sparse series evaluation when the imaginary part of the argument becomes too large. In the implementation, the AGM code is disabled for an imaginary part larger than 5.

Comparing first the evaluation of η as a sparse series or by multipoint evaluation, one notices that the asymptotically faster algorithm is about 5 to 8 times slower on the examples and that it appears to catch up with growing class numbers. However, this happens so slowly that one cannot expect it to beat the algorithm in $O(|D|^{1.25+\varepsilon})$ in the foreseeable future for any tractable instances.

The approach using Newton iterations on the AGM is faster than multipoint evaluation, but still hardly beats the asymptotically slower algorithm in $O(|D|^{1.25+\varepsilon})$: the biggest computed example of class number 100000 lies just beyond the cross-over point!

One notices that the growth rates of the running times of all algorithms, instead of behaving like $|D|$ or $|D|^{1.25}$, come closer to $|D|^{1.4}$ or $|D|^{1.6}$. A small part of this can be explained by the rather peculiar choice of discriminants. Taking the first one with a given class number, the precision n is rather large compared to h and $\sqrt{|D|}$, since there are many forms with small A so that the sum $\sum_{i=1}^h \frac{1}{A_i}$ becomes comparatively large.

However, the major reason for the faster than predicted growth of computing time is that for the floating point arithmetic the range for asymptotically fast algorithms is not yet reached. The threshold for switching to the FFT in gmp on the test machine is set to about 500000 bits; so the examples lie still in the Karatsuba or Toom–Cook range, which accounts for a growth of $M(n)$ of $n^{\log 3 / \log 2}$ resp. $n^{\log 5 / \log 3}$ instead of n .

A possible improvement of the multipoint evaluation approach consists of conveniently grouping

the arguments. For instance, in the example of class number 5000, the function η has to be evaluated in 2501 arguments (corresponding to the two ambiguous forms and 2499 pairs of opposite non-ambiguous forms). Assuming the worst case of $|q| \approx e^{-\pi\sqrt{3}}$, that is almost reached for the largest values of A , one can approximate η by a polynomial of degree 1190. As multipoint evaluation should be most efficient when the number of arguments is about half the degree, it makes sense to perform four evaluations in 625 resp. 626 arguments each. But sorting them by their absolute values, the smaller ones do not actually require such a high degree approximation of η . In the example, an approximation of degree 260, 551, 805 resp. 1190 is sufficient for the four chunks of q . Then the time used for multipoint evaluation drops from 93 s to 64 s. Experimenting with different partitions of the arguments (three resp. five parts of the same size, parts of different sizes adapted to the degree of the approximations, etc.) yields similar results, far from competing with the sparse series evaluation.

Another point to take into account are the space requirements of the algorithms. When each root of the class polynomial is computed separately, only $O(hn)$ bits need to be stored, which is linear in the output size. Multipoint evaluation as described in Section 3.2, however, requires that the tree constructed in Step 1 of Algorithm 6 be maintained in memory, so that the occupied space grows by a logarithmic factor to reach $O(hn \log h)$. This logarithmic factor could be saved by evaluating in chunks of $O\left(\frac{h}{\log h}\right)$ arguments, as explained in Lemma 2.1 of [GS92].

Anyway, the example class polynomial of degree 100000 uses over 5 GB as an uncompressed text file and is computed in about 3 days. This shows that the limiting factor is the memory requirement rather than the running time, as can be expected from algorithms that have a close to linear complexity with respect to their output size.

As a final remark, one notices that the algorithms behave numerically well, even though rounding errors do occur during floating point computations. For the algorithm of Section 6.2, this can be explained by the sparsity of the η series and the fact that all coefficients are $+1$ and -1 . Indeed, if the last few digits of a term are erroneous, these errors propagate to subsequent terms. However, the absolute magnitude of such errors decreases rapidly, so that the wrong digits in later terms actually do not intervene in the additions. (Otherwise said, the computations may as well be carried out with fixed point numbers, and indeed a simulation of fixed point arithmetic using floating point numbers of decreasing precision yields accurate results.)

8 Comparison to other approaches

8.1 Chinese remaindering

In [ALV04], the authors suggest an approach for directly computing class polynomials for j modulo a prime p . The basic idea is to derive the polynomial modulo many small primes by enumerating all elliptic curves modulo these small primes and only retaining those having complex multiplication by \mathcal{O}_D . Then a Chinese remainder technique allows to obtain the class polynomial modulo p .

Unfortunately, it is not sufficient to gather only $O(\log p)$ bits of information per coefficient of the class polynomial modulo small primes, although this is the information contained in the final output. In fact, so many small primes are needed that the class polynomial could be reconstructed over \mathbb{Z} instead of only modulo p . The complexity derived in Section 3.2 of [ALV04] is

$$O\left(|D|^{3/2} \log^{10} |D| + |D| \log^2 |D| \log p + \sqrt{|D|} \log^2 p\right),$$

and already the term depending only on $|D|$ is worse than what is obtained with the practically employed algorithm of Section 6.2, not to speak of the asymptotically fast versions of Sections 6.3 and 6.4.

Thus, it is asymptotically faster to compute the class polynomial over \mathbb{Z} using floating point approximations and to reduce it modulo p afterwards.

8.2 p -adic algorithms

Couveignes and Henocq in [CH02] suggest a p -adic approach for computing class polynomials for j . The basic idea is to look for a small prime p and (by exhaustive search) an elliptic curve modulo p with complex multiplication by \mathcal{O}_D . This curve is then lifted to the p -adic numbers with a high enough precision so that the class polynomial may be reconstructed. Alternatively, they show how to work with supersingular curves. The complexity of their approach is $O(|D|^{1+\varepsilon})$, where the exact

power of the logarithmic factor has not been worked out. By nature, the algorithm is not affected by rounding errors, and using the explicit bound of Theorem 2 its output is certified to be correct.

So the complex and the p -adic approach are both essentially linear in the size of the class polynomials. All variants have been implemented (for more details on the complex implementation, see [EM02]; for the ordinary p -adic algorithm, see [BS04]; for the supersingular one, [LR04]), and all seem to work reasonably well in practice. The floating point algorithms are easy to implement with arbitrary class invariants using the results of [Sch02]. The p -adic approach with ordinary curves has been made to work with class invariants other than j , see [BS04] and [Brö06, Chapter 6]. The considerable overhead involved makes it unclear whether it is competitive with the complex approach.

Acknowledgements

I am grateful to Régis Dupont for letting me use his code to test the fast evaluation of η via the arithmetic-geometric mean. I thank Karim Belabas for pointing out to me the basic idea of Section 4, Pierrick Gaudry and François Morain for numerous discussions and Dan Bernstein, Reinier Bröker and Kristin Lauter for helpful comments.

References

- [AB04] A. O. L. Atkin and D. J. Bernstein. Prime sieves using binary quadratic forms. *Mathematics of Computation*, 73(246):1023–1030, 2004.
- [ALV04] Amod Agashe, Kristin Lauter, and Ramarathnam Venkatesan. Constructing elliptic curves with a known number of points over a prime field. In Alf van der Poorten and Andreas Stein, editors, *High Primes and Misdemeanours: Lectures in Honour of the 60th Birthday of Hugh Cowie Williams*, volume 41 of *Fields Institute Communications*. American Mathematical Society, 2004.
- [AM93] A. O. L. Atkin and F. Morain. Elliptic curves and primality proving. *Mathematics of Computation*, 61(203):29–68, July 1993.
- [BLS03] Paulo S. L. M. Barreto, Ben Lynn, and Michael Scott. Constructing elliptic curves with prescribed embedding degrees. In Stelvio Cimato, Clemente Galdi, and Giuseppe Persiano, editors, *Security in Communication Networks — Third International Conference, SCN 2002, Amalfi, Italy, September 2002*, volume 2576 of *Lecture Notes in Computer Science*, pages 257–267, Berlin, 2003. Springer-Verlag.
- [BP05] N. Brisebarre and G. Philibert. Effective lower and upper bounds for the Fourier coefficients of powers of the modular invariant j . *Journal of the Ramanujan Mathematical Society*, 20:255–282, 2005.
- [Bre76] Richard P. Brent. Fast multiple-precision evaluation of elementary functions. *Journal of the ACM*, 23(2):242–251, 1976.
- [Brö06] Reinier Bröker. *Constructing elliptic curves of prescribed order*. PhD thesis, Universiteit Leiden, 2006.
- [BS04] Reinier Bröker and Peter Stevenhagen. Elliptic curves with a given number of points. In Duncan Buell, editor, *Algorithmic Number Theory — ANTS-VI*, volume 3076 of *Lecture Notes in Computer Science*, pages 117–131, Berlin, 2004. Springer-Verlag.
- [BW05] Friederike Brezing and Annegret Weng. Elliptic curves suitable for pairing based cryptography. *Designs, Codes and Cryptography*, 37(1):133–141, 2005.
- [CH02] Jean-Marc Couveignes and Thierry Henocq. Action of modular correspondences around CM points. In Claus Fieker and David R. Kohel, editors, *Algorithmic Number Theory — ANTS-V*, volume 2369 of *Lecture Notes in Computer Science*, pages 234–243, Berlin, 2002. Springer-Verlag.
- [Cip03] M. Cipolla. Un metodo per la risoluzione della congruenza di secondo grado. *Napoli Rend.*, 9:153–163, 1903.

- [CNST98] Jinhui Chao, Osamu Nakamura, Kohji Sobataka, and Shigeo Tsujii. Construction of secure elliptic cryptosystems using CM tests and liftings. In Kazuo Ohta and Dingyi Pei, editors, *Advances in Cryptology — ASIACRYPT '98*, volume 1514 of *Lecture Notes in Computer Science*, pages 95–109, Berlin, 1998. Springer-Verlag.
- [Coh93] Henri Cohen. *A Course in Computational Algebraic Number Theory*. Graduate Texts in Mathematics. Springer-Verlag, New York, 1993.
- [Cox89] David A. Cox. *Primes of the Form $x^2 + ny^2$ — Fermat, Class Field Theory, and Complex Multiplication*. John Wiley & Sons, New York, 1989.
- [Ded76] R. Dedekind. Erläuterungen zu den vorstehenden Fragmenten. In R. Dedekind and H. Weber, editors, *Bernhard Riemann's gesammelte mathematische Werke und wissenschaftlicher Nachlaß*, pages 438–447. Teubner, Leipzig, 1876.
- [DEM05] Régis Dupont, Andreas Enge, and François Morain. Building curves with arbitrary small MOV degree over finite prime fields. *Journal of Cryptology*, 18(2):79–89, 2005.
- [Deu41] Max Deuring. Die Typen der Multiplikatorenringe elliptischer Funktionenkörper. *Abhandlungen aus dem mathematischen Seminar der hamburgischen Universität*, 14:197–272, 1941.
- [Deu58] Max Deuring. Die Klassenkörper der komplexen Multiplikation. In *Enzyklop. d. math. Wissenschaften*, volume I 2 Heft 10. Teubner, Stuttgart, 2 edition, 1958.
- [Dup07] Régis Dupont. Fast evaluation of modular functions using Newton iterations and the AGM. To appear in *Mathematics of Computation*; available at http://www.lix.polytechnique.fr/Labo/Regis.Dupont/preprints/Dupont_FastEvalMod.ps.gz, 2007.
- [EM02] Andreas Enge and François Morain. Comparing invariants for class fields of imaginary quadratic fields. In Claus Fieker and David R. Kohel, editors, *Algorithmic Number Theory — ANTS-V*, volume 2369 of *Lecture Notes in Computer Science*, pages 252–266, Berlin, 2002. Springer-Verlag.
- [EM07] Andreas Enge and François Morain. Further investigations of the generalised Weber functions. In preparation, 2007.
- [ES04] Andreas Enge and Reinhard Schertz. Constructing elliptic curves over finite fields using double eta-quotients. *Journal de Théorie des Nombres de Bordeaux*, 16:555–568, 2004.
- [Eul83] Leonhard Euler. Evolutio producti infiniti $(1-x)(1-xx)(1-x^3)(1-x^4)(1-x^5)(1-x^6)$ etc. in seriem simplicem. *Acta academiae scientiarum Petropolitanae*, 1780:I:125–169, 1783. *Opera Omnia* I.3:472–479.
- [EZ] Andreas Enge and Paul Zimmermann. *mpc* — a library for multiprecision complex arithmetic with exact rounding. Version 0.4.5, available at <http://www.lix.polytechnique.fr/Labo/Andreas.Engel/Software.html>.
- [Gau05] Pierrick Gaudry. Assembly support for GMP on AMD64, April 2005. Available at http://www.lix.polytechnique.fr/Labo/Pierrick.Gaudry/mpn_AMD64/.
- [Gee99] Alice Gee. Class invariants by Shimura's reciprocity law. *Journal de Théorie des Nombres de Bordeaux*, 11(1):45–72, 1999.
- [GG99] Joachim von zur Gathen and Jürgen Gerhard. *Modern Computer Algebra*. Cambridge University Press, 1999.
- [Gra] Torbjörn Granlund et. al. *gmp* — GNU multiprecision library. Version 4.2.1, available at <http://www.swox.com/gmp>.
- [GS92] Joachim von zur Gathen and Victor Shoup. Computing Frobenius maps and factoring polynomials. *Computational Complexity*, 2:187–224, 1992.
- [GS98] Alice Gee and Peter Stevenhagen. Generating class fields using Shimura reciprocity. In J. P. Buhler, editor, *Algorithmic Number Theory — ANTS-III*, volume 1423 of *Lecture Notes in Computer Science*, pages 441–453, Berlin, 1998. Springer-Verlag.

- [HLPZ] Guillaume Hanrot, Vincent Lefèvre, Patrick Pélicier, and Paul Zimmermann et. al. `mpfr` — a library for multiple-precision floating-point computations with exact rounding. Version 2.2.0, available at <http://www.mpfr.org>.
- [LL90] A. K. Lenstra and H. W. Lenstra Jr. Algorithms in number theory. In Jan van Leeuwen, editor, *Algorithms and Complexity*, volume A of *Handbook of Theoretical Computer Science*, chapter 12, pages 673–715. Elsevier, Amsterdam, 1990.
- [LR04] R. Lercier and E. Riboulet-Deyris. Elliptic curves with complex multiplication. Posting to the Number Theory List, available at <http://listserv.nodak.edu/cgi-bin/wa.exe?A2=ind0401&L=nmbirthry&P=R305>, 2004.
- [MNT01] Atsuko Miyaji, Masaki Nakabayashi, and Shunzou Takano. New explicit conditions of elliptic curve traces for FR-reduction. *IEICE Trans. Fundamentals*, E84-A(5):1234–1243, May 2001.
- [Sch76] Reinhard Schertz. Die singulären Werte der Weberschen Funktionen $f, f_1, f_2, \gamma_2, \gamma_3$. *Journal für die reine und angewandte Mathematik*, 286/287:46–74, 1976.
- [Sch91] René Schoof. The exponents of the groups of points on the reductions of an elliptic curve. In G. van der Geer, F. Oort, and J. Steenbrink, editors, *Arithmetic Algebraic Geometry*, pages 325–335, Boston, 1991. Birkhäuser.
- [Sch02] Reinhard Schertz. Weber’s class invariants revisited. *Journal de Théorie des Nombres de Bordeaux*, 14(1):325–343, 2002.
- [Sie36] Carl Ludwig Siegel. Über die Classenzahl quadratischer Zahlkörper. *Acta Arithmetica*, 1:83–86, 1936.
- [Web08] Heinrich Weber. *Lehrbuch der Algebra*, volume 3. Chelsea Publishing Company, New York, 3rd edition, 1908.