



HAL
open science

The union of Unit Balls has Quadratic Complexity, even if They all Contain the Origin

Hervé Brönnimann, Olivier Devillers

► **To cite this version:**

Hervé Brönnimann, Olivier Devillers. The union of Unit Balls has Quadratic Complexity, even if They all Contain the Origin. RR-3758, INRIA. 1999. inria-00072904

HAL Id: inria-00072904

<https://inria.hal.science/inria-00072904v1>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*The union of unit balls has quadratic complexity,
even if they all contain the origin*

Hervé Brönnimann — Olivier Devillers

N° 3758

Septembre 1999

THÈME 2



*Rapport
de recherche*

The union of unit balls has quadratic complexity, even if they all contain the origin

Hervé Brönnimann , Olivier Devillers

Thème 2 — Génie logiciel
et calcul symbolique
Projet Prisme

Rapport de recherche n° 3758 — Septembre 1999 — 8 pages

Abstract: We provide a lower bound construction showing that the union of unit balls in \mathbb{R}^3 has quadratic complexity, even if they all contain the origin. This settles a conjecture of Sharir.

Key-words: Computational geometry, Union of balls, Geometric example

Une union de boules unités d'intersection non vide a une complexité quadratique

Résumé : Nous proposons une construction d'un ensemble de boules de \mathbb{R}^3 dont l'union a un nombre quadratique de faces et d'arêtes. Ces boules ont la particularité d'avoir le même rayon et d'avoir une intersection non vide. Cette note démontre une conjecture énoncé par M. Sharir.

Mots-clés : Géométrie algorithmique, Union de boules, Exemple géométrique

1 Introduction

The union of a set of n balls in \mathbb{R}^3 has quadratic complexity $\Theta(n^2)$, even if they all have the same radius. All the already known constructions have balls scattered around, however, and Sharir posed the problem whether a quadratic complexity could be achieved if all the balls (of same radius) contained the origin.

In this note, we show a construction of n unit balls, all containing the origin, whose union has complexity $\Theta(n^2)$. As a trivial observation, we observe that the centers are arbitrarily close to the origin in our construction. In fact, if the centers are forced to be at least pairwise ε apart, for some constant $\varepsilon > 0$, then no more than $O(\frac{1}{\varepsilon^3})$ can meet in a single point, and hence the union has complexity at most $O(\frac{1}{\varepsilon^3}n) = O_\varepsilon(n)$. It is an interesting open question what a condition should be so that the union have subquadratic complexity and yet the balls have arbitrarily close centers.

By contrast, the *intersection* of n balls can have quadratic complexity if their radii are not constrained, but the complexity is linear if all the radii are the same [2]. Similarly, the convex hull of n balls can have also quadratic complexity [1], but that complexity is linear if they all have the same radius.

2 Construction

Let m and k be any integers. We define two families of unit balls: the first consists of k unit balls whose centers lie on a small vertical segment; the second consists of m unit balls whose centers lie on a small circle under the segment. (See Figure 3.) We show below that their union has quadratic $O(km)$ complexity.

The balls $B_1 \dots B_k$. We denote by $B(p, r)$ the ball centered at p and of radius r . Let $n = k + m$ and P_i denote the point of coordinates $(0, 0, (i - 1)/n^4)$, and $B_i = B(P_i, 1)$, for $i = 1, \dots, k$. It is clear that the boundary of $\cup_{1 \leq i \leq k} B_i$ consists of two hemispheres belonging to B_1 and B_k linked by a narrow cylinder of height less than $k/n^4 \leq 1/n^3$. This cylinder contains all the circles $\partial B_i \cap \partial B_{i+1}$ for $i = 1, \dots, k - 1$. (See Figure 1.)




Figure 1: The union $\cup_{1 \leq i \leq k} B_i$.

The balls $B_{k+1} \dots B_{k+m}$. Let R be the point of coordinates $(x, 0, z)$ with

$$x = \frac{2n^2 - 4}{n^4}, \quad z = -\frac{2n^2 - 4}{n^3}.$$

(Any values satisfying the constraints $P_k H < 1$ in (1) and $\ell < \frac{2}{n}$ in (2) below would do.) We define θ as the rotation around the z -axis of angle $2\pi/m$, and for each $j = 1, \dots, m$, $R_{k+j} = \theta^{j-1}(R)$ and $B_{k+j} = B(R_{k+j}, 1)$.

3 Analysis

By our choice of x and z , we prove below that the boundaries of B_{k+1} and of the union $\cup_{i=1}^k B_i$ depicted in Figure 1 meet along a curve γ which satisfies the two claims below. The situation is depicted on Figure 2.

Claim 1 *The curve γ intersects all the balls B_i for $i = 0, \dots, k, .$*

Claim 2 *The portion of γ which does not belong to B_1 (equivalently, which belongs to the union $\cup_{i=2}^k B_i$) is contained in an angular sector of angle at most $2\pi/m$.*

From claim 2, we conclude that the portion of γ which does not belong to B_1 is contained in the boundary of the union of the $n = k + m$ balls, From claim 1, we conclude that the portion of γ which does not belong to B_1 has complexity $\Omega(k)$. From claim 2, that it is contained in a small angular sector, hence appears completely on the boundary of the union of the $n = k + m$ balls, and it is replicated m times, once for each of the balls B_j , $j = 1, \dots, m$. It




Figure 2: The union $\bigcup_{1 \leq i \leq k} B_i \cup B_{k+1}$. The curve γ consists of a portion that belongs to B_1 and of another portion which is contained in a dihedral sector of angle less than π/m .

follows that the union of all the balls B_i for $i = 1, \dots, k + m$ has quadratic complexity $\Omega(km)$. Moreover, all the balls contain the origin. The union of the n balls is depicted on Figure 3.

The proofs involve only elementary geometry and trigonometry. The situation is depicted in Figure 4 and 5. Figure 4 depicts a section in the xz -plane of the spheres ∂B_i and ∂B_{k+1} and the point M , the highest point of intersection of the bounding spheres. The point M is also depicted on Figure 2.

Proof of Claim 1. It suffices to prove that M is higher than P_k , since then γ extends higher than P_k as well and passes through M by symmetry. The lowest point of γ belongs to B_1 and is clearly below the origin. The two facts together prove that γ must intersect all the balls between B_1 and B_k .




Figure 3: The union $\cup_{1 \leq i \leq k+m} B_i$. On the right, a blow-up of the centers.

Let H be the point in the xz -plane on the median bisector of R and P_k , with same z -ordinate as P_k . (See Figure 4.) In order to prove that M is higher than P_k , it suffices to prove that H belongs to B_k , since then M is farther along the bisector. The two triangles QP_kH and KRP_k have equal angles, hence they are similar. It follows that

$$P_k H = P_k R \frac{P_k Q}{R K} = \frac{P_k R^2}{2 R K} = \frac{x^2 + (z - z_k)^2}{2x}, \quad (1)$$

where $z_k = \frac{k-1}{n^4}$. For x and z as given in the construction, we have

$$P_k H = 1/16 \frac{-40 n^4 - 15 n^2 + 68 + 16 n^6 - 16 n^3 + 28 n}{n^4 (n^2 - 2)}$$

which is smaller than 1 for $n \geq 2$.

Proof of Claim 2. It is easy to see that the intersection of γ and a ball B_i ($2 \leq i \leq k$) consists of at most two arcs of circle, any of which is monotone in angular coordinates around the z -axis, and that any such arc is entirely above the plane $z = 0$. Hence the intersections of γ with the xy -plane belong to B_1 and B_{k+1} . It suffices to show that these intersections are at a distance ℓ at most $\frac{2}{n} \leq \sin \frac{\pi}{m}$ from the x -axis. (See Figure 5.)

In the xy -plane section, B_1 is a unit circle, and B_{k+1} is a circle of radius $r = \sqrt{1 - z^2}$ and center R' of coordinates $(x, 0)$. (Recall that the center of




Figure 4: Figure for Claim 1.

B_{k+1} has coordinates $(x, 0, z)$.) Hence ℓ is the height of a triangle with base x and sides 1 and $r < 1$. It is elementary to compute that

$$\ell = \sqrt{1 - \left(\frac{z^2 + x^2}{2x}\right)^2}. \quad (2)$$

For our choice of x and z , this yields

$$\ell = \sqrt{\frac{2n^6 + 3n^4 - 4n^2 - 4}{n^8}}$$

which is smaller than $2/n$ for $n \geq 2$.

Acknowledgments. Thanks to Micha Sharir for pointing out the problem to us. It was also pointed out that Alon Efrat might have a construction which leads to a quadratic lower bound as well. We have derived our construction independently.

References

- [1] Jean-Daniel Boissonnat, André Cérézo, Olivier Devillers, Jacqueline Duquesne, and Mariette Yvinec. An algorithm for constructing the con-




Figure 5: Figure for Claim 2.

vex hull of a set of spheres in dimension d . *Comput. Geom. Theory Appl.*, 6:123–130, 1996.

- [2] A. Heppes. Beweis einer Vermutung von A. Vázsonyi. *Acta Math. Acad. Sci. Hungar.*, 7:463–466, 1956.



Unité de recherche INRIA Sophia Antipolis
2004, route des Lucioles - B.P. 93 - 06902 Sophia Antipolis Cedex (France)

Unité de recherche INRIA Lorraine : Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - B.P. 101 - 54602 Villers lès Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38330 Montbonnot St Martin (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - B.P. 105 - 78153 Le Chesnay Cedex (France)

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, B.P. 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399