



**HAL**  
open science

## Solution of a Minimax Problem with Additive Final Cost

Laura S. Aragone, Silvia C. Di Marco, Roberto L.V. González

► **To cite this version:**

Laura S. Aragone, Silvia C. Di Marco, Roberto L.V. González. Solution of a Minimax Problem with Additive Final Cost. [Research Report] RR-3394, INRIA. 1998. inria-00073295

**HAL Id: inria-00073295**

**<https://inria.hal.science/inria-00073295>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

*Solution of a Minimax Problem  
with Additive Final Cost*

Laura S. Aragone - Silvia C. Di Marco  
Roberto L.V. González

N° 3394

Avril 1998

THÈME 4

*R*apport  
de recherche

Les rapports de recherche de l'INRIA  
sont disponibles en format postscript sous  
ftp.inria.fr (192.93.2.54)

si vous n'avez pas d'accès ftp  
la forme papier peut être commandée par mail :  
e-mail : dif.gesdif@inria.fr  
(n'oubliez pas de mentionner votre adresse postale).

par courrier :  
Centre de Diffusion  
INRIA  
BP 105 - 78153 Le Chesnay Cedex (FRANCE)

INRIA research reports  
are available in postscript format  
ftp.inria.fr (192.93.2.54)

if you haven't access by ftp  
we recommend ordering them by e-mail :  
e-mail : dif.gesdif@inria.fr  
(don't forget to mention your postal address).

by mail :  
Centre de Diffusion  
INRIA  
BP 105 - 78153 Le Chesnay Cedex (FRANCE)

# Solution of a minimax problem with additive final cost \*

Laura S. Aragone<sup>†</sup>, Silvia C. Di Marco<sup>‡</sup> & Roberto L.V. González<sup>‡</sup>

## Abstract

In this paper we study a minimax optimal control problem with finite horizon and additive final cost. We introduce a generalized problem where the original one can be embedded. For this generalized problem, we establish a dynamical programming principle and we present the associated Hamilton-Jacobi-Bellman (HJB) equation. This equation is defined in terms of a discontinuous Hamiltonian and we prove that the optimal cost of the generalized problem is the unique viscosity solution of this HJB equation.

**Key words:** *minimax problem, dynamical programming principle, Hamilton-Jacobi-Bellman equation, viscosity solution*

---

\*This paper is included in the activities developed in the frame of the Cooperation Project INRIA-Instituto Beppo Levi, Coordinators of the project: E. Rofman - R. González.

<sup>†</sup>CONICET-Universidad Nacional de Quilmes, Argentina.

<sup>‡</sup>CONICET - Inst. Beppo Levi, FCEIA, Universidad Nacional de Rosario, Argentina.

## Solution d'un problème d'optimisation minimax avec coût final additif

**Résumé:** On analyse ici un problème de contrôle optimal de type minimax avec coût final additif. On introduit un problème auxiliaire généralisant le problème original et permettant d'utiliser les techniques de la programmation dynamique. Pour le problème généralisé, on établit un principe de la programmation dynamique et on présente aussi l'équation de Hamilton-Jacobi-Bellman (HJB) associée. On démontre que le coût optimal du problème généralisé est l'unique solution de viscosité de cette équation de HJB.

**Motclefs:** *problèmes minimax, principe de la programmation dynamique, équation de Hamilton-Jacobi-Bellman, solution de viscosité*

**AMS Classification:** 49S35, 49L05, 49L20, 49L25

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The optimization problem</b>	<b>2</b>
2.1	Presentation of the problem . . . . .	2
2.2	General assumptions . . . . .	2
<b>3</b>	<b>Auxiliary problem</b>	<b>3</b>
3.1	Auxiliary variable and problem reformulation . . . . .	3
3.2	Properties of the optimal cost $v$ . . . . .	3
3.3	Dynamical programming equation . . . . .	4
<b>4</b>	<b>Hamilton-Jacobi-Bellman equation</b>	<b>4</b>
4.1	The optimal cost as a viscosity solution . . . . .	5
4.2	Uniqueness of the viscosity solution . . . . .	8

# 1 Introduction

The optimization of dynamic systems where the criterion is the maximum value of a function is a frequent problem in technology, economics and industry. This problem appears for example, when we attempt to minimize the maximum deviation of controlled trajectories with respect to a given "model" trajectory. Minimax problems differs from those usually considered in the optimal control literature where a cumulative cost is minimized. Since in some cases, minimax problems describe more appropriately decision problems arisen in controlled systems whose performance is evaluated with a unique scalar parameter, the minimax optimization has received considerable attention in recent publications (see, for example, [3, 4, 5, 6, 7, 8, 9, 10, 19]). Furthermore, the relation of minimax problems with the design of robust controllers can be seen in [11].

In addition, from the academic point of view, the minimax optimal control problem is of interest in the area of game theory because minimax problems can be seen as a game – see [14] – where a player applies ordinary controls and the other one – using complete and privileged information – chooses a stopping time. Problems of this type lead to the treatment of nonlinear partial differential inequalities akin to those appearing in the *obstacle problem* (with obstacle given in explicit or implicit form, see [7]). To find solutions of these systems we must consider generalized solutions – even discontinuous solutions – since commonly there does not exist classical solutions of such systems (see [1, 2]). The treatment of the infinite horizon problem also presents great analytical difficulties, because the optimal cost is neither necessarily lower semicontinuous nor upper semicontinuous. Moreover, the optimal cost cannot be approximated with a sequence of finite horizon problems. Studies concerning these issues can be seen in [15, 18]. Besides, it is also important to develop numerical methods to compute these solutions in approximate way because in general it is not possible to find exact analytical solutions. Numerical methods to obtain approximated open loop optimal controls are analyzed in [20] and for closed loop, see [4, 14].

Here, we analyze a minimax optimal control problem where the functional to be optimized, not only depends on the maximum of a function along the complete trajectory of the system but also it takes into account (in an additive fashion) another function of the final state of the system.

For a problem with this structure, a dynamical programming principle cannot be formulated merely in terms of the initial time and the initial state. To obtain a dynamical programming principle, we introduce an auxiliary parameter which "remembers" the past maximum values (the use of a similar procedure can be seen in [17, 18]). Using this parameter, we present an auxiliary problem which gives the solution of the original problem using a particular value of the auxiliary parameter. For this second optimal control problem, we establish the associated dynamical programming principle and a Hamilton-Jacobi-Bellman (HJB) equation. Finally, we prove that the optimal cost of the auxiliary problem is the unique solution of the associated HJB equation.

The paper is organized as follows. In §2, we present the optimization problem. In §3, we describe the auxiliary problem and its relation with the original one. We also establish there the dynamical programming equation. In §4, we give the HJB equation associated to the problem and we prove the uniqueness of solution in the viscosity sense.

## 2 The optimization problem

### 2.1 Presentation of the problem

We consider a minimax optimal control problem with finite horizon and final cost. More specifically, the problem consists in minimizing the functional  $J$

$$\begin{aligned} J : [0, T] \times \mathbb{R}^m \times A &\mapsto \mathbb{R} \\ (t, x, \alpha(\cdot)) &\mapsto J(t, x, \alpha(\cdot)) = \operatorname{ess\,sup}_{s \in [t, T]} f(s, y(s), \alpha(s)) + \Psi(y(T)), \end{aligned}$$

where  $y(\cdot)$  represents the state of a dynamic system which evolves accordingly to the following differential equation,

$$\begin{cases} y'(s) = g(s, y(s), \alpha(s)) & \forall t \leq s < T \\ y(t) = x, & x \in \mathbb{R}^m. \end{cases}$$

$A = L^\infty([0, T], A)$  is the set of admissible control policies and  $A$  is the control set.

The optimal cost function is given by

$$u(t, x) = \inf_{\alpha(\cdot) \in A} J(t, x, \alpha(\cdot)). \quad (1)$$

This problem is an extension of another one analyzed by Barron-Ishii in [7] and by Di Marco-González in [12, 13]. We are interested in analyzing the new problem from the continuous viewpoint.

After introducing an auxiliary problem, which generalized the problem presented above, we establish the dynamical programming principle corresponding to the new problem, and we present a Hamilton-Jacobi-Bellman (HJB) equation associated to the optimal cost. Finally, we prove that the optimal cost is the unique solution of this HJB equation.

### 2.2 General assumptions

Let  $BUC([0, T] \times \mathbb{R}^m \times A)$  be the set of bounded and uniformly continuous functions in  $[0, T] \times \mathbb{R}^m \times A$ . We assume the following hypotheses hold:

$$(H_1) \quad g : [0, T] \times \mathbb{R}^m \times A \mapsto \mathbb{R}^m, \quad g \in BUC([0, T] \times \mathbb{R}^m \times A),$$

$$\|g(t, x, a)\| \leq M_g,$$

$$\|g(t, x, a) - g(\hat{t}, \hat{x}, a)\| \leq L_g(|t - \hat{t}| + \|x - \hat{x}\|), \quad \forall t, \hat{t} \in [0, T], \forall x, \hat{x} \in \mathbb{R}^m, \forall a \in A.$$

$$(H_2) \quad f : [0, T] \times \mathbb{R}^m \times A \mapsto \mathbb{R}, \quad f \in BUC([0, T] \times \mathbb{R}^m \times A),$$

$$m_f \leq f(t, x, a) \leq M_f,$$

$$|f(t, x, a) - f(\hat{t}, \hat{x}, a)| \leq L_f(|t - \hat{t}| + \|x - \hat{x}\|), \quad \forall t, \hat{t} \in [0, T], \forall x, \hat{x} \in \mathbb{R}^m, \forall a \in A.$$

$$(H_3) \quad \Psi : \mathbb{R}^m \mapsto \mathbb{R}, \quad \Psi \in BUC(\mathbb{R}^m) \text{ and}$$

$$|\Psi(x) - \Psi(\hat{x})| \leq L_\Psi \|x - \hat{x}\|, \quad \forall x, \hat{x} \in \mathbb{R}^m.$$

$$(H_4) \quad A \text{ is compact.}$$



### 3 Auxiliary problem

#### 3.1 Auxiliary variable and problem reformulation

In the problem presented above it is not possible to establish a dynamical programming principle only in terms of the variables  $(t, x)$ . In order to develop the analytical and numerical treatment of the problem we extend the state of the system introducing an auxiliary variable  $y_{m+1}(\cdot)$ . The initial value of this variable is  $y_{m+1}(t) = \rho$  and the evolution of  $y_{m+1}$  is given by the following expression:

$$y_{m+1}(\tau) = \max \left\{ y_{m+1}(\sigma), \operatorname{ess\,sup}_{s \in [\sigma, \tau]} f(s, y(s), \alpha(s)) \right\}, \quad \sigma \in [t, T]. \quad (2)$$

**Note 3.1** *The interpretation of the additional variable  $y_{m+1}$  is the following one:  $y_{m+1}$  remembers the maximum values of the function  $f$  from the initial time  $t$  to the current time  $\tau$ . To make it clear, let us consider  $m_f(t, x) = \min_{a \in A} f(t, x, a)$  and  $\rho \leq m_f(x)$ . Then, from (2), it follows that*

$$y_{m+1}(T) = \operatorname{ess\,sup}_{s \in [t, T]} f(s, y(s), \alpha(s)).$$

This variable  $y_{m+1}$  is a solution of the differential inclusion given by

$$\frac{dy_{m+1}}{ds}(s) \in \overline{G}(f(s, y(s), \alpha(s)) - y_{m+1}(s)), \quad (3)$$

where

$$\overline{G}(w) = \begin{cases} 0 & \text{if } w < 0, \\ [0, \infty] & \text{if } w = 0, \\ \infty & \text{if } w > 0. \end{cases}$$

In this way, the original problem can be studied as an ordinary optimal control problem, i.e. a problem with pure final cost because  $J(t, x, \alpha(\cdot)) = y_{m+1}(T) + \Psi(y(T))$ . In this case, the optimal cost  $v$  is:

$$v(t, x, \rho) = \inf \{ y_{m+1}(T) + \Psi(y(T)) : \alpha(\cdot) \in \mathcal{A} \}, \quad (4)$$

where  $t \in [0, T]$ ,  $x \in \mathbb{R}^m$  and  $\rho \in \mathbb{R}$ , being  $\rho$  the initial value for equation (3), i.e.  $y_{m+1}(t) = \rho$ .

**Remark 3.1** *From (2), it is clear that  $\{y_{m+1}(s) : s \in (t, T]\} \subseteq [\nu, M_f]$ , where  $\nu = \max\{\rho, m_f\}$ . Moreover, if  $\rho \geq M_f$ , then  $y_{m+1}(s) = \rho, \forall s \in [t, T]$ .*

#### 3.2 Properties of the optimal cost $v$

The following properties brings some relations between the optimal costs  $u$  and  $v$  of the original and auxiliary problems. They are almost evident and can be proved without difficulties using the definitions of the original and auxiliary problems described above. Here we omit the complete proofs for the sake of brevity and we only sketch the lines of argument.

- If  $\rho \leq \min_{a \in A} f(t, x, a)$ , then

$$v(t, x, \rho) = u(t, x). \quad (5)$$

By replacing in (4) and comparing with (1), it follows that  $v(t, x, \rho) = u(t, x)$ ,

- If  $\rho \geq \max_{a \in A} f(t, x, a)$ , then  $v(t, x, \rho) = \hat{u}(t, x) + \rho$ , where  $\hat{u}$  is the optimal cost of the control problem (1) when  $f \equiv 0$ , i.e. where the functional to be minimized is  $\Psi(y(T))$ .

In this case  $y_{m+1}(s) = \rho, \forall s \in [t, T]$ . Then, by replacing in (4), we have that

$$v(t, x, \rho) = \rho + \inf_{\alpha(\cdot) \in A} \Psi(y(T)). \quad (6)$$

- If  $\rho_1 > \rho_2$ , then

$$v(t, x, \rho_1) \geq v(t, x, \rho_2). \quad (7)$$

Let  $\rho_1 > \rho_2$  and  $y_{m+1}^{\rho_i}(\cdot)$  the solution of the differential inclusion (3) when the initial condition is  $\rho_i$ . Then,

$$\begin{aligned} y_{m+1}^{\rho_1}(T) &= \max\{\rho_1, \text{ess sup}_{s \in [t, T]} f(s, y(s), \alpha(s))\} \\ &\geq \max\{\rho_2, \text{ess sup}_{s \in [t, T]} f(s, y(s), \alpha(s))\} = y_{m+1}^{\rho_2}(T). \end{aligned} \quad (8)$$

By replacing in (4) it results  $v(t, x, \rho_1) \geq v(t, x, \rho_2)$ .

- $v$  is bounded  $\forall \rho \in [-K, K]$ .

This property follows from the definition of  $v$  and the boundedness of  $f$  and  $\Psi$ .

- $v$  is Lipschitz continuous with respect to the three variables  $t, x, \rho$ .

This property follows from the hypotheses verified by  $f, g$  and  $\Psi$ .

- $v$  may be not semiconcave.

In effect, if  $\Psi \equiv 0$ .  $u$  can be not semiconcave (see [12]), then  $v$  has the same property (see examples in [14]).

### 3.3 Dynamical programming equation

It is clear that in this new problem with the state augmented by the variable  $\rho$ , the dynamical programming equation is given by

$$v(t, x, \rho) = \inf_{\alpha(\cdot) \in L^\infty(t, s], A)} v(s, y(s), y_{m+1}(s)) \quad (9)$$

We also have the final condition

$$v(T, x, \rho) = \max\left\{\rho, \min_{a \in A} f(T, x, a)\right\} + \Psi(x). \quad (10)$$

## 4 Hamilton-Jacobi-Bellman equation

Let us define

$$H_A(t, x, \rho, v) = \inf \left\{ v(t, x, f(t, x, a)) - v(t, x, \rho) : \right. \\ \left. f(t, x, a) > \rho, v(t, x, f(t, x, a)) \leq v(t, x, \rho) \right\},$$

$$H(t, x, \rho, \nabla v) = \inf \left\{ \frac{\partial v}{\partial t}(t, x, \rho) + \frac{\partial v}{\partial x}(t, x, \rho)g(t, x, a) : f(t, x, a) < \rho \right\},$$

$$H_*(t, x, \rho, \nabla v) = \inf \left\{ \frac{\partial v}{\partial t}(t, x, \rho) + \frac{\partial v}{\partial x}(t, x, \rho)g(t, x, a) : f(t, x, a) \leq \rho \right\}.$$

Taking into account the dynamical programming equation we obtain the following differential formulation. Strictly, it takes the following form

$$\begin{cases} \min \{H_A(t, x, \rho, v), H(t, x, \rho, \nabla v)\} \geq 0, \\ \min \{H_A(t, x, \rho, v), H_*(t, x, \rho, \nabla v)\} \leq 0, \end{cases} \quad (11)$$

with final condition

$$v(T, x, \rho) = \max \left\{ \rho, \min_{a \in A} f(T, x, a) \right\} + \Psi(x). \quad (12)$$

The solution in the viscosity sense of this system (11) is defined as follows.

- The function  $w$  is a *subsolution* in the viscosity sense if,
  - $w$  is upper-semicontinuous on  $(0, T) \times \mathbb{R}^m \times \mathbb{R}$ ,
  - $w$  is upper-bounded on  $(0, T) \times \mathbb{R}^m \times [-K, K]$ ,  $\forall K \in \mathbb{R}^+$ ,
  - $w$  is a non-decreasing function of the variable  $\rho$ ,
  - given  $(t, x, \rho) \in (0, T) \times \mathbb{R}^m \times \mathbb{R}$  and  $\Phi \in C^1((0, T) \times \mathbb{R}^m \times \mathbb{R})$  such that  $\Phi - w$  has a minimum in  $(t, x, \rho)$  in a neighborhood  $\mathcal{N}(t, x, \rho)$ , it results

$$H(t, x, \rho, \nabla \Phi) \geq 0. \quad (13)$$

- The function  $z$  is a *supersolution* in the viscosity sense if,
  - $z$  is lower-semicontinuous on  $(0, T) \times \mathbb{R}^m \times \mathbb{R}$ ,
  - $z$  is lower-bounded on  $(0, T) \times \mathbb{R}^m \times [-K, K]$ ,  $\forall K \in \mathbb{R}^+$ ,
  - given  $(t, x, \rho) \in (0, T) \times \mathbb{R}^m \times \mathbb{R}$  and  $\Phi \in C^1((0, T) \times \mathbb{R}^m \times \mathbb{R})$  such that  $\Phi - z$  has a maximum in  $(t, x, \rho)$  in a neighborhood  $\mathcal{N}(t, x, \rho)$ , it results

$$\min \{H_A(t, x, \rho, z), H_*(t, x, \rho, \nabla \Phi)\} \leq 0. \quad (14)$$

By virtue of the especial definition of viscosity solution given above, we can use the following equivalent form of the HJB equation:

$$\begin{cases} H(t, x, \rho, \nabla v) \geq 0, \\ \min \{H_A(t, x, \rho, v), H_*(t, x, \rho, \nabla v)\} \leq 0. \end{cases} \quad (15)$$

## 4.1 The optimal cost as a viscosity solution

**Theorem 4.1** *The optimal cost  $v$  is a solution in the viscosity sense of the system (11).*

**Proof.** The final condition is trivially verified from (9) and it was proved that  $v$  is continuous, bounded and non-decreasing with respect to  $\rho$ . So it remains to prove the last property of the subsolution's definition and the last one of the supersolution's definition.

First, we will prove that  $v$  is a *subsolution* of (11).

Let  $(t, x, \rho) \in (0, T) \times \mathbb{R}^m \times \mathbb{R}$  and  $\Phi \in C^1((0, T) \times \mathbb{R}^m \times \mathbb{R})$  such that  $\Phi - v$  has a minimum in  $(t, x, \rho)$  in a neighborhood  $\mathcal{N}(t, x, \rho)$ .

Let us prove that  $H(t, x, \rho, \nabla \Phi) \geq 0$ . Let  $a \in A$  such that  $f(t, x, a) < \rho$  and  $\{\alpha_n\}$  be a control sequence such that  $\alpha_n(s) = a$ , for all  $s \in [t, t + n^{-1}]$ .

From equation (9), we have that

$$v(t, x, \rho) \leq v(t + n^{-1}, y_n(t + n^{-1}), \max\{\rho, \operatorname{esssup}_{s \in [t, t + n^{-1}]} f(s, y_n(s), \alpha_n(s))\}) \quad (16)$$

Since  $\lim_{n \rightarrow \infty} \operatorname{esssup}_{s \in [t, t+n^{-1}]} f(s, y_n(s), \alpha_n(s)) = f(t, x, a)$ , for all  $n \geq n_0$ , we get

$$\max \left\{ \rho, \operatorname{esssup}_{s \in [t, t+n^{-1}]} f(s, y_n(s), \alpha_n(s)) \right\} = \rho$$

and then, from (16), we obtain

$$v(t, x, \rho) \leq v(t + n^{-1}, y_n(t + n^{-1}), \rho). \quad (17)$$

By the minimality of  $\Phi - v$  at  $(t, x, \rho)$ , it follows that

$$\Phi(t + n^{-1}, y_n(t + n^{-1}), \rho) - \Phi(t, x, \rho) \geq v(t + n^{-1}, y_n(t + n^{-1}), \rho) - v(t, x, \rho) \geq 0. \quad (18)$$

Since

$$\lim_{n \rightarrow \infty} \frac{y_n(t + n^{-1}) - x}{n^{-1}} = \lim_{n \rightarrow \infty} n \int_t^{t+n^{-1}} g(s, y(s), \alpha_n(s)) ds = g(t, x, a),$$

dividing by  $n^{-1}$  in (18) and taking limit when  $n \rightarrow \infty$ , we have that

$$\frac{\partial \Phi}{\partial t}(t, x, \rho) + \frac{\partial \Phi}{\partial x}(t, x, \rho) g(t, x, a) \geq 0.$$

Then, since  $a$  is arbitrary, it results that

$$H(t, x, \rho, \nabla \Phi) \geq 0. \quad (19)$$

So, (13) and (19) imply that  $v$  is subsolution.

Let us see that  $v$  is a *supersolution* of (11).

Let  $(t, x, \rho) \in (0, T) \times \mathbb{R}^m \times \mathbb{R}$  and  $\Phi \in C^1((0, T) \times \mathbb{R}^m \times \mathbb{R})$  such that  $\Phi - v$  has a maximum in  $(t, x, \rho)$  in a neighborhood  $\mathcal{N}(t, x, \rho)$ .

We assume that  $\exists \eta > 0$  such that

$$\min \{H_A(t, x, \rho, v), H_*(t, x, \rho, \nabla \Phi)\} \geq \eta > 0. \quad (20)$$

Let  $\{\alpha_n\}$  be a minimizing control sequence such that

$$v(t, x, \rho) + \frac{1}{n^2} \geq v(t + n^{-1}, y_n(t + n^{-1}), \max\{\rho, \operatorname{esssup}_{s \in [t, t+n^{-1}]} f(s, y_n(s), \alpha_n(s))\}). \quad (21)$$

By the compactness of  $A$ , there exists  $a \in A$  and a subsequence of  $\{\alpha_n\}$  such that

$$\lim_{n \rightarrow \infty} \operatorname{esssup}_{s \in [t, t+n^{-1}]} f(s, y_n(s), \alpha_n(s)) = f(t, x, a).$$

The case  $f(t, x, a) > \rho$  can be ruled out.

In effect, the inequality (20) implies that  $H_A(t, x, \rho, v) \geq \eta$ . But taking limit in (21) and by virtue of (20) and (7), it follows that  $v(t, x, \rho) \geq v(t, x, \rho) + \eta$ , which is absurd.

So, we have that  $f(t, x, a) \leq \rho$ . We will analyze the effect of this condition on the relation  $H_*(t, x, \rho, \nabla \Phi) \geq \eta > 0$  (valid by virtue of (20)).

As  $f(t, x, a) \leq \rho$ , then for all  $\varepsilon > 0$ ,  $f(t, x, a) < \rho + \varepsilon$ . In consequence, there exists  $n_\varepsilon$  such that  $\forall n \geq n_\varepsilon$ ,

$$\operatorname{ess\,sup}_{s \in [t, t+n^{-1}]} f(s, y_n(s), \alpha_n(s)) < \rho + \varepsilon. \quad (22)$$

Eventually, by redefining the controls  $\alpha_n(\cdot)$  in zero measure sets, it is possible to affirm that  $\forall n \geq n_\varepsilon, \forall s \in [t, t+n^{-1}]$ ,

$$f(s, y_n(s), \alpha_n(s)) < \rho + \varepsilon. \quad (23)$$

Since  $f$  is Lipschitz-continuous, we have that  $\forall n \geq n_\varepsilon, \forall s \in [t, t+n^{-1}]$ ,

$$f(t, x, \alpha_n(s)) < \rho + \varepsilon + L_f(1 + M_g)n^{-1} < \rho + 2\varepsilon. \quad (24)$$

We consider the set of controls  $Z_\varepsilon$  given by

$$Z_\varepsilon = \{a \in A : f(t, x, a) \leq \rho + 2\varepsilon\}.$$

Then, for any  $n \geq n_\varepsilon$  and  $s \in [t, t+n^{-1}]$ , it results  $\alpha_n(s) \in Z_\varepsilon$ .

As  $(t, x, \rho)$  is a maximum point for the function  $\Phi - v$ , from (21), we have

$$\Phi(t+n^{-1}, y_n(t+n^{-1}), \rho) - \Phi(t, x, \rho) \leq v(t+n^{-1}, y_n(t+n^{-1}), \rho) - v(t, x, \rho) \leq n^{-2}. \quad (25)$$

It is possible to prove that there exists  $\bar{g} \in \overline{Co(g(t, x, Z_\varepsilon))}$ , where  $Co(E)$  is the convex hull of the set  $E$ , such that (eventually using a suitable subsequence)

$$\lim_{n \rightarrow \infty} n \int_t^{t+n^{-1}} g(s, y_n(s), \alpha_n(s)) ds = \bar{g}$$

and so, by virtue of (25), we get

$$\lim_{n \rightarrow \infty} \frac{\Phi(t+n^{-1}, y_n(t+n^{-1}), \rho) - \Phi(t, x, \rho)}{n^{-1}} = \frac{\partial \Phi}{\partial t}(t, x, \rho) + \frac{\partial \Phi}{\partial x}(t, x, \rho) \bar{g} \leq 0. \quad (26)$$

On the other hand, as  $Z_\varepsilon$  is closed, we have that  $\overline{Co(g(t, x, Z_\varepsilon))} = Co(g(t, x, Z_\varepsilon))$ .

As  $\bar{g} \in Co(g(t, x, Z_\varepsilon)) \forall \varepsilon > 0$ , we get that  $\bar{g} \in Co(g(t, x, Z_0))$ . Then, there exists a probability measure  $\mu(\cdot)$  with support in  $Z_0$  such that

$$\bar{g} = \int_{Z_0} g(t, x, a) d\mu(a). \quad (27)$$

From (20), if  $a \in Z_0$ , then  $\frac{\partial \Phi}{\partial t}(t, x, \rho) + \frac{\partial \Phi}{\partial x}(t, x, \rho)g(t, x, a) \geq \eta > 0$ .

Using this inequality and the integral expression (27) in (26) we have that,

$$0 \geq \frac{\partial \Phi}{\partial t}(t, x, \rho) + \frac{\partial \Phi}{\partial x}(t, x, \rho) \bar{g} = \int_{Z_0} \left( \frac{\partial \Phi}{\partial t}(t, x, \rho) + \frac{\partial \Phi}{\partial x}(t, x, \rho)g(t, x, a) \right) d\mu(a) \geq \eta. \quad (28)$$

This inequality contradicts the initial assumption (20) and so, we obtain by *reductio ad absurdum* that  $v$  is a supersolution. □

## 4.2 Uniqueness of the viscosity solution

**Theorem 4.2** *There is a unique solution of the system (11).*

**Proof.** Let  $w$  be a subsolution and  $z$  a supersolution of (11).

We will prove by *reductio ad absurdum* that  $w \leq z$ .

Let us suppose that  $\exists K > 0$  such that

$$r := \sup \{w(t, x, \rho) - z(t, x, \rho) : (0, T) \times \mathbb{R}^m \times [-K, K]\} > 0, \quad (29)$$

then given  $0 < \delta < \frac{r}{2}$  there exists  $(t_\delta, x_\delta, \rho_\delta) \in (0, T) \times \mathbb{R}^m \times [-K, K]$  such that

$$w(t_\delta, x_\delta, \rho_\delta) - z(t_\delta, x_\delta, \rho_\delta) > \sup_{(0, T) \times \mathbb{R}^m \times [-K, K]} (w(t, x, \rho) - z(t, x, \rho)) - \delta > \frac{r}{2}.$$

Let  $M > 0$  such that

$$\begin{cases} w(t, x, \rho) \leq M, & \text{in } [0, T] \times \mathbb{R}^m \times [-2K, 2K], \\ z(t, x, \rho) \geq -M, & \text{in } [0, T] \times \mathbb{R}^m \times [-2K, 2K]. \end{cases} \quad (30)$$

As  $w$  is upper semicontinuous,  $\exists \omega_w : \mathbb{R}^+ \mapsto \mathbb{R}^+$  such that

$$\sup \{w(t', x', \rho') : |t' - t| + \|x' - x\| + |\rho' - \rho| \leq \eta\} < w(t, x, \rho) + \omega_w(\eta). \quad (31)$$

For each  $\epsilon$  we consider the hump function

$$\Phi(t, x, s, y) = -\frac{|t-s|^2}{\epsilon^2} - \frac{\|x-y\|^2}{\epsilon^2} - \sigma\left(\frac{T}{t} - 1\right) - \frac{r}{8}\left(1 - \frac{t}{T}\right) + (\delta + \omega_w(\sqrt{\epsilon}))\xi(\|x - x_\delta\|). \quad (32)$$

The elements of  $\Phi$  have the following properties:

- $\xi(\cdot) \in C^1(\mathbb{R})$  verifies

$$\begin{cases} \xi(\tau) = 0 & \tau = [0, 1], \\ \xi(\tau) = -1 & \tau \geq 3, \\ |\xi'(\tau)| \leq 1. \end{cases}$$

- $\sigma < \min \left\{ M, \frac{r t_\delta}{8(T - t_\delta)} \right\}$ .

- The derivatives of  $\Phi$  are

$$\begin{cases} \frac{\partial \Phi}{\partial t}(t, x, s, y) = -2\frac{(t-s)}{\epsilon^2} + \sigma\frac{T}{t^2} + \frac{r}{8T}, \\ \frac{\partial \Phi}{\partial x}(t, x, s, y) = -2\frac{(x-y)}{\epsilon^2} + (\delta + \omega_w(\sqrt{\epsilon}))\xi'(\|x - x_\delta\|)\frac{(x - x_\delta)}{\|x - x_\delta\|}, \\ \frac{\partial \Phi}{\partial s}(t, x, s, y) = 2\frac{(t-s)}{\epsilon^2}, \\ \frac{\partial \Phi}{\partial y}(t, x, s, y) = 2\frac{(x-y)}{\epsilon^2}. \end{cases} \quad (33)$$

We define the function

$$\phi^{\sqrt{\epsilon}}(t, x, \rho, s, y) = z(t, x, \rho) - w(s, y, \rho + \sqrt{\epsilon}) - \Phi(t, x, s, y). \quad (34)$$

As  $w$  is nondecreasing, from (29) and (31) we have

$$w(t, x, \rho + \sqrt{\epsilon}) - z(t, x, \rho) \leq r + \omega_w(\sqrt{\epsilon}),$$

so the function  $\phi^{\sqrt{\epsilon}}$  has a minimum in  $(0, T) \times \mathbb{R}^m \times [-K, K] \times [0, T] \times \mathbb{R}^m$ .

Let  $(t_0, x_0, \rho_0, s_0, y_0)$  be one of these minimizing points. From (30) and (32), we get

$$\frac{|t_0 - s_0|^2}{\epsilon^2} + \frac{\|x_0 - y_0\|^2}{\epsilon^2} + \sigma \left( \frac{T - t_0}{t_0} \right) + \frac{r}{8} \left( 1 - \frac{t_0}{T} \right) \leq 2M + \sigma \left( \frac{T - t_\delta}{t_\delta} \right) + \frac{r}{8} \left( 1 - \frac{t_\delta}{T} \right).$$

If we define  $M_1 = 2M + \sigma \frac{T}{t_\delta} + \frac{r}{8}$ , the last inequality and the definition of  $\Phi$  imply that

$$\begin{cases} \frac{\sigma T}{M_1} < t_0, \\ |t_0 - s_0| < \sqrt{M_1} \epsilon, \\ \|x_0 - y_0\| < \sqrt{M_1} \epsilon, \\ \|x_0 - x_\delta\| \leq 3. \end{cases} \quad (35)$$

In particular, as  $(t_0, x_0, \rho_0, s_0, y_0)$  is a minimizing point

$$\phi^{\sqrt{\epsilon}}(t_0, x_0, \rho_0, s_0, y_0) \leq \phi^{\sqrt{\epsilon}}(t_0, x_0, \rho_0, t_0, x_0).$$

and so, we have

$$\frac{|t_0 - s_0|^2}{\epsilon^2} + \frac{\|x_0 - y_0\|^2}{\epsilon^2} \leq w(s_0, y_0, \rho_0 + \sqrt{\epsilon}) - w(t_0, x_0, \rho_0 + \sqrt{\epsilon}) \leq \omega_w(\sqrt{\epsilon}). \quad (36)$$

From (36), we get

$$\begin{cases} |t_0 - s_0| < \epsilon \sqrt{\omega_w(\sqrt{\epsilon})}, \\ \|x_0 - y_0\| < \epsilon \sqrt{\omega_w(\sqrt{\epsilon})}. \end{cases} \quad (37)$$

Let us define the function  $\phi_1 \in C^1((0, T) \times \mathbb{R}^m \times \mathbb{R})$  as follows:

$$\phi_1(t, x, \rho) = w(s_0, y_0, \rho + \sqrt{\epsilon}) + \Phi(t, x, s_0, y_0).$$

From (34),  $\phi_1 - z$  has a maximum in  $(t_0, x_0, \rho_0)$ , then, as  $z$  is supersolution, it follows that

$$\min(H_A(t_0, x_0, \rho_0, z), H_*(t_0, x_0, \rho_0, \nabla \phi_1)) \leq 0. \quad (38)$$

We can suppose w.l.g. that  $H_A(t_0, x_0, \rho_0, z) = +\infty$ . (see the justification of this argument in the Note 4.1 given below).

Therefore, it must be  $\min(H_A(t_0, x_0, \rho_0, z), H_*(t_0, x_0, \rho_0, \nabla \phi_1)) = H_*(t_0, x_0, \rho_0, \nabla \phi_1)$ , so

$$H_*(t_0, x_0, \rho_0, \nabla \phi_1) \leq 0. \quad (39)$$

Then, it must exist  $\bar{a}$  such that  $f(t_0, x_0, \bar{a}) \leq \rho_0$  and

$$\frac{\partial \phi_1}{\partial t}(t_0, x_0, \rho_0) + \frac{\partial \phi_1}{\partial x}(t_0, x_0, \rho_0) g(t_0, x_0, \bar{a}) \leq 0. \quad (40)$$

On the other hand, by defining  $\phi_2 \in C^1((0, T) \times \mathbb{R}^m \times \mathbb{R})$

$$\phi_2(s, y, \rho) = z(t_0, x_0, \rho - \sqrt{\epsilon}) - \Phi(t_0, x_0, s, y),$$

from (34) we have that  $\phi_2 - w$  has a minimum in  $(s_0, y_0, \rho_0 + \sqrt{\epsilon})$ .

As  $w$  is a subsolution we have

$$H(s_0, y_0, \rho_0 + \sqrt{\epsilon}, \nabla \phi_2) \geq 0. \quad (41)$$

Since  $|s_0 - t_0| \leq \epsilon \sqrt{\omega_w(\sqrt{\epsilon})}$  and  $\|x_0 - y_0\| \leq \epsilon \sqrt{\omega_w(\sqrt{\epsilon})}$  we can suppose that  $\epsilon$  is small enough such that

$$f(s_0, y_0, \bar{a}) \leq f(t_0, x_0, \bar{a}) + 2L_f \epsilon \sqrt{\omega_w(\sqrt{\epsilon})} < \rho_0 + \sqrt{\epsilon}$$

and so, by definition of  $H$  we have that

$$\frac{\partial \phi_2}{\partial s}(s_0, y_0, \rho_0 + \sqrt{\epsilon}) + \frac{\partial \phi_2}{\partial y}(s_0, y_0, \rho_0 + \sqrt{\epsilon}) g(s_0, y_0, \bar{a}) \geq 0. \quad (42)$$

Besides,

$$\left\{ \begin{array}{l} \frac{\partial \phi_2}{\partial s}(s, y, \rho) = -\frac{\partial \Phi}{\partial s}(t_0, x_0, s, y) = -2 \frac{(t_0 - s)}{\epsilon^2} \\ \frac{\partial \phi_2}{\partial y}(s, y, \rho) = -\frac{\partial \Phi}{\partial y}(t_0, x_0, s, y) = -2 \frac{(x_0 - y)}{\epsilon^2} \\ \frac{\partial \phi_1}{\partial t}(t, x, \rho) = \frac{\partial \Phi}{\partial t}(t, x, s_0, y_0) = -2 \frac{(t - s_0)}{\epsilon^2} + \sigma \frac{T}{t^2} + \frac{r}{8T} \\ \frac{\partial \phi_1}{\partial x}(t, x, \rho) = \frac{\partial \Phi}{\partial x}(t, x, s_0, y_0) = -2 \frac{(x - y_0)}{\epsilon^2} \\ \qquad \qquad \qquad + (\delta + \omega_w(\sqrt{\epsilon})) \xi'(\|x - x_\delta\|) \frac{(x - x_\delta)}{\|x - x_\delta\|} \end{array} \right.$$

Using these expressions in (40) and (42) we have

$$\begin{aligned} 0 \leq \frac{r}{8T} &\leq -\sigma \frac{T}{t_0^2} + 2 \frac{\langle x_0 - y_0, g(t_0, x_0, \bar{a}) - g(s_0, y_0, \bar{a}) \rangle}{\epsilon^2} \\ &\quad - (\delta + \omega_w(\sqrt{\epsilon})) \xi'(\|x_0 - x_\delta\|) \frac{\langle x_0 - x_\delta, g(t_0, x_0, \bar{a}) \rangle}{\|x_0 - x_\delta\|} \\ &\leq \frac{2L_g}{\epsilon^2} \|x_0 - y_0\| (\|x_0 - y_0\| + |t_0 - s_0|) + (\delta + \omega_w(\sqrt{\epsilon})) \|g(t_0, x_0, \bar{a})\| \\ &\leq 4L_g \omega_w(\epsilon) + (\delta + \omega_w(\sqrt{\epsilon})) M_g. \end{aligned}$$

Letting successively  $\epsilon$  and  $\delta$  go to zero, we have that  $r \leq 0$  which contradicts the inequality (29). Then  $z \geq w$ . Finally, taking into account that any solution is at the same time a subsolution and a supersolution, we obtain the uniqueness of solution.  $\square$



**Note 4.1** If  $(t_0, x_0, \rho_0, s_0, y_0)$  realizes the minimum of

$$\phi^{\sqrt{\epsilon}}(t, x, \rho, s, y) = z(t, x, \rho) - w(s, y, \rho + \sqrt{\epsilon}) - \Phi(t, x, s, y), \quad (43)$$

then  $\exists (t_0, x_0, \bar{\rho}_0, s_0, y_0)$  that also realizes the minimum of  $\phi^{\sqrt{\epsilon}}$  and the point  $(t_0, x_0, \bar{\rho}_0)$  has the property:  $H_A(t_0, x_0, \bar{\rho}_0, z) = +\infty$ .

**Proof.** Let us suppose that

$$\min(H_A(t_0, x_0, \rho_0, z), H_*(t_0, x_0, \rho_0, \nabla \phi_1)) = H_A(t_0, x_0, \rho_0, z),$$

then

$$H_A(t_0, x_0, \rho_0, z) \leq 0. \quad (44)$$

In consequence, there exists  $a' \in A$ , such that

$$\left\{ \begin{array}{l} f(t_0, x_0, a') > \rho_0 \\ \text{and} \\ z(t_0, x_0, f(t_0, x_0, a')) \leq z(t_0, x_0, \rho_0). \end{array} \right. \quad (45)$$

From the continuity of  $f$  and the lower-semicontinuity of  $z$ , it is obvious that there exists at least one  $\bar{a} \in A$  such that

$$f(t_0, x_0, \bar{a}) = \max\{f(t_0, x_0, a') : a' \in A \text{ and (45)}\} \quad (46)$$

We define:

$$\bar{\rho}_0 = f(t_0, x_0, \bar{a}). \quad (47)$$

Since  $w$  is a subsolution, it is a non-decreasing function of the variable " $\rho$ " and as  $\bar{\rho}_0 + \sqrt{\epsilon} > \rho_0 + \sqrt{\epsilon}$ , we have

$$w(s_0, y_0, \bar{\rho}_0 + \sqrt{\epsilon}) \geq w(s_0, y_0, \rho_0 + \sqrt{\epsilon}). \quad (48)$$

From the inequalities (45) and (48) it follows that

$$w(s_0, y_0, \bar{\rho}_0 + \sqrt{\epsilon}) - z(t_0, x_0, \bar{\rho}_0) \geq w(s_0, y_0, \rho_0 + \sqrt{\epsilon}) - z(t_0, x_0, \rho_0).$$

So, we have that

$$\begin{aligned} w(s_0, y_0, \bar{\rho}_0 + \sqrt{\epsilon}) - z(t_0, x_0, \bar{\rho}_0) + \Phi(t_0, x_0, s_0, y_0) \\ \geq w(s_0, y_0, \rho_0 + \sqrt{\epsilon}) - z(t_0, x_0, \rho_0) + \Phi(t_0, x_0, s_0, y_0). \end{aligned}$$

This last inequality proves that  $(t_0, x_0, \bar{\rho}_0, s_0, y_0)$  is also a point where  $\phi^{\sqrt{\epsilon}}$  has a minimum.

Let us now show that

$$H_A(t_0, x_0, \bar{\rho}_0, z) = +\infty \quad (49)$$

If we suppose that  $H_A(t_0, x_0, \bar{\rho}_0, z) < \infty$ , then

$$H_A(t_0, x_0, \bar{\rho}_0, z) \leq 0. \quad (50)$$

In consequence, it would exist  $a'' \in A$ , such that

$$\left. \begin{array}{l} f(t_0, x_0, a'') > \bar{\rho}_0 \\ \text{and} \\ z(t_0, x_0, f(t_0, x_0, a'')) \leq z(t_0, x_0, \bar{\rho}_0). \end{array} \right\} \quad (51)$$

From (45)-(47) and (51) we have

$$z(t_0, x_0, f(t_0, x_0, a'')) \leq z(t_0, x_0, \bar{\rho}_0) = z(t_0, x_0, f(t_0, x_0, \bar{a})) \leq z(t_0, x_0, \rho_0)$$

and

$$f(t_0, x_0, a'') > \bar{\rho}_0 = f(t_0, x_0, \bar{a}) > \rho_0. \quad (52)$$

These inequalities contradict the maximality of  $f(t_0, x_0, \bar{a})$  given by (46). So, by *reductio ad absurdum* we get (49). □

## Conclusions

In the minimax optimal control problem analyzed in this paper, by extending the state of the system, we have embedded the original problem into another one where it is possible to establish a dynamical programming principle and a HJB equation with unique solution in the viscosity sense. The optimal cost of the auxiliary problem is that unique solution and this property enables us to characterize the optimal cost and to deal with the numerical analysis of the problem.

## Acknowledgments

The authors would like to thank:

- CONICET for support given through the grant PID BID-CONICET N°4796.
- The Universidad Nacional de Rosario, Argentina, for support given through the grant PID N°19/1028.
- The authorities of INRIA for the support given through the Cooperation Project INRIA-Instituto de Matemática Beppo Levi.

## References

- [1] Barles G., *Discontinuous viscosity solutions of first order Hamilton-Jacobi equations: A guided visit*, Nonlinear Analysis, Theory, Methods and Applications, **20**, **9**, 1123-1134, (1993).
- [2] Barles G., *Solutions de viscosité des équations de Hamilton Jacobi*, Springer-Verlag, New York (1994).
- [3] Barles G., Ch. Daher and M. Romano, *Optimal control on the  $L^\infty$  norm of a diffusion process*, SIAM J. on Control and Optimization, **32**, **3**, 612-634 (1994).
- [4] Barles G., Ch. Daher and M. Romano, *Convergence of numerical schemes for parabolic equations arising in finance theory*, Math. Models and Methods in Applied Sc., **5**, **1**, 125-143 (1995).
- [5] Barron E.N., *Differential games with maximum cost*, Nonlinear Analysis, Theory, Methods and Applications, **14**, **11**, 971-989 (1990).

- [6] Barron E.N., *Averaging in Lagrange and minimax problems of optimal control*, SIAM J. on Control and Optimization, **31**, 6, 1630-1652 (1993).
- [7] Barron E.N. and H. Ishii, *The Bellman equation for minimizing the maximum cost*, Nonlinear Analysis, Theory, Methods and Applications, **13**, 9, 1067-1090 (1989).
- [8] Barron E.N. and R. Jensen, *Relaxed minimax control*, SIAM J. on Control and Optimization, **33**, 4, 1028-1039 (1995).
- [9] Barron E.N., R. Jensen and J.L. Menaldi, *Optimal control and differential games with measures*, Nonlinear Analysis, Theory, Methods and Applications, **21**, 4, 241-268 (1993).
- [10] Barron E.N. and W. Liu, *Semicontinuous solutions for Hamilton-Jacobi equations and the  $L^\infty$  control problem*, Applied Math. Optim. **34**, 325-360 (1996).
- [11] Başar T. and P. Bernhard,  *$H^\infty$  optimal control and related minimax design problems. A dynamic game approach*, Birkhäuser, Berlin (1991).
- [12] Di Marco S.C., *Sobre la optimización minimax y tiempos de detención óptimos*, Thesis, University of Rosario, Argentine (1996).
- [13] Di Marco S.C. and R.L.V. González, *Une procédure numérique pour la minimisation du coût maximum*, Comptes Rendus Acad. Sc. Paris, Série I, Tome **321**, 869-874 (1995).
- [14] Di Marco S.C. and R.L.V. González, *A numerical procedure for minimizing the maximum cost*, in System modelling and optimization, J. Dolezal and J. Fidler (Eds.), Chap. **33**, 285-291, Chapman & Hall Eds., London (1996); (extended version: Rapport de Recherche N°2454, INRIA, Rocquencourt (1995)).
- [15] Di Marco S.C. and R.L.V. González, *Deterministic minimax optimal control problem with infinite horizon*, in Proceedings Fourth WG-7.6 Working Conference on Optimization-Based Computer Aided Modelling and Design, 9.1-9.3. ESIEE, Noisy le Grand (1996).
- [16] Di Marco S.C. and R.L.V. González, *A minimax optimal control problem with infinite horizon*, Rapport de Recherche N°2945, INRIA, Rocquencourt (1996).
- [17] Di Marco S.C. and R.L.V. González, *A stochastic minimax optimal control problem on Markov chains with infinite horizon*, Rapport de Recherche N°2946, INRIA, Rocquencourt (1996).
- [18] Di Marco S.C. and R.L.V. González, *A finite state stochastic minimax optimal control problem with infinite horizon*, in Numerical Analysis and Its Applications, L. Vulkov, J. Wasniewski, P. Yalamov (Eds.), Lectures Notes of Computer Science N°1196, 134-141, Springer-Verlag, Berlin (1997).
- [19] González R.L.V. and L.S. Aragone, *A Bellman's equation for minimizing the maximum cost*, Rapport de Recherche N°3290, INRIA, Rocquencourt (1997).
- [20] Polak E. and L. He, *Rate-preserving discretization strategies for semiinfinite programming and optimal control*, SIAM J. on Control and Optimization, **30**, 4, 548-572 (1992).



---

Unité de recherche INRIA Rocquencourt

Domaine de Voluceau - Rocquencourt - B.P. 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Lorraine : Technopôle de Nancy-Brabois - Campus scientifique

615, rue du Jardin Botanique - B.P. 101 - 54602 Villers lès Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38330 Montbonnot St Martin (France)

Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - B.P. 93 - 06902 Sophia Antipolis Cedex (France)

---

Éditeur

INRIA - Domaine de Voluceau - Rocquencourt, B.P. 105 - 78153 Le Chesnay Cedex (France)

<http://www.inria.fr>

ISSN 0249-6399



★ R R . 3 3 9 4 ★