



# Global convergence properties of conjugate gradient methods for optimization

Jean Charles Gilbert, Jorge Nocedal

## ► To cite this version:

Jean Charles Gilbert, Jorge Nocedal. Global convergence properties of conjugate gradient methods for optimization. SIAM Journal on Optimization, 1992, 2 (1), pp.21-42. 10.1137/0802003 . inria-00075291

**HAL Id: inria-00075291**

**<https://inria.hal.science/inria-00075291>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License



UNITÉ DE RECHERCHE  
INRIA-ROCQUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
B.P.105  
78153 Le Chesnay Cedex  
France  
Tél.: (1) 39 63 55 11

# Rapports de Recherche

N° 1268

*Programme 5*  
*Automatique, Productique,*  
*Traitement du Signal et des Données*

## GLOBAL CONVERGENCE PROPERTIES OF CONJUGATE GRADIENT METHODS FOR OPTIMIZATION

Jean Charles GILBERT  
Jorge NOCEDAL

Juillet 1990



\* R R - 1 2 6 8 \*

INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

**GLOBAL CONVERGENCE PROPERTIES OF  
CONJUGATE GRADIENT METHODS FOR OPTIMIZATION**

**PROPRIÉTÉS DE CONVERGENCE GLOBALE DES  
MÉTHODES DE GRADIENT CONJUGUÉ EN OPTIMISATION**

by

*Jean Charles Gilbert,<sup>1</sup> and Jorge Nocedal<sup>2</sup>*

Rapport de Recherche n° 1268

June 1990

---

<sup>1</sup> INRIA, Domaine de Voluceau, Rocquencourt, B.P. 105, 78153 Le Chesnay Cedex (France).

<sup>2</sup> Department of Electrical Engineering and Computer Science, Northwestern University, Evanston IL 60208. This author was supported by the U.S. Department of Energy, under Grant DE-FG02-87ER25047-A001, and by National Science Foundation Grant No. CCR-8902096.

GLOBAL CONVERGENCE PROPERTIES OF  
CONJUGATE GRADIENT METHODS FOR OPTIMIZATION

PROPRIÉTÉS DE CONVERGENCE GLOBALE DES  
MÉTHODES DE GRADIENT CONJUGUÉ EN OPTIMISATION

by

*Jean Charles Gilbert and Jorge Nocedal*

ABSTRACT

We study the convergence of nonlinear conjugate gradient methods without restarts, and with practical line searches. The analysis covers two classes of methods that are globally convergent on smooth, non-convex functions. Some properties of the Fletcher-Reeves method play an important role in the first family, whereas the second family shares an important property with the Polak-Ribière method. Numerical experiments are presented.

RÉSUMÉ

Nous étudions la convergence des méthodes de gradient conjugué pour la minimisation des fonctions sans contrainte. Les méthodes sont supposées être implémentées sans redémarrage et avec une recherche linéaire réaliste (non exacte). L'analyse couvre deux classes de méthodes qui sont globalement convergentes pour les fonctions régulières non nécessairement convexes. Dans la première famille, ce sont certaines propriétés de la méthode de Fletcher-Reeves qui jouent un rôle crucial, tandis que la seconde famille partage avec la méthode de Polak-Ribière une propriété importante. Des résultats numériques sont présentés.

*Key words:* conjugate gradient method, global convergence, unconstrained optimization, large-scale optimization.

*Abbreviated title:* Conjugate gradient methods.

## 1. Introduction.

The object of this paper is to study the convergence properties of several conjugate gradient methods for nonlinear optimization. We consider only the case where the methods are implemented without regular restarts, and ask under what conditions are they globally convergent for general smooth nonlinear functions. The analysis will allow us to highlight differences among various conjugate gradient methods, and will suggest new implementations.

Our problem is to minimize a function of  $n$  variables,

$$\min f(x), \quad (1.1)$$

where  $f$  is smooth, and its gradient  $g$  is available. We consider iterations of the form

$$d_k = \begin{cases} -g_k & \text{for } k = 1 \\ -g_k + \beta_k d_{k-1} & \text{for } k \geq 2, \end{cases} \quad (1.2)$$

$$x_{k+1} = x_k + \alpha_k d_k, \quad (1.3)$$

where  $\beta_k$  is a scalar, and  $\alpha_k$  is a steplength obtained by means of a one-dimensional search. We call this iteration a *conjugate gradient method* if  $\beta_k$  is such that (1.2)-(1.3) reduces to the linear conjugate gradient method in the case when  $f$  is a strictly convex quadratic and  $\alpha_k$  is the exact one-dimensional minimizer. Some of the results of this paper, however, also apply to methods of the form (1.2)-(1.3) that do not reduce to the linear conjugate gradient method.

The best known formulas for  $\beta_k$  are called the Fletcher-Reeves (FR), Polak-Ribière (PR) and Hestenes-Stiefel (HS) formulas, and are given by

$$\beta_k^{\text{FR}} = \|g_k\|^2 / \|g_{k-1}\|^2, \quad (1.4)$$

$$\beta_k^{\text{PR}} = \langle g_k, g_k - g_{k-1} \rangle / \|g_{k-1}\|^2, \quad (1.5)$$

$$\beta_k^{\text{HS}} = \langle g_k, g_k - g_{k-1} \rangle / \langle d_{k-1}, g_k - g_{k-1} \rangle. \quad (1.6)$$

Here,  $\langle \cdot, \cdot \rangle$  is the scalar product used to compute the gradient and  $\|\cdot\|$  denotes its associated norm. The numerical performance of the Fletcher-Reeves (1964) method is somewhat erratic: it is sometimes as efficient as the Polak-Ribière and Hestenes-Stiefel methods, but it is often much slower. Powell (1977) gives an argument showing that, under some circumstances, the Fletcher-Reeves method with exact lines searches will produce very small displacements, and will normally not recover unless a restart along the gradient direction is performed. In spite of these drawbacks, Zoutendijk (1970) has shown that the method cannot fail. He proved that the Fletcher-Reeves method with exact line searches is globally convergent on general functions. Al-Baali (1985) extended this result to inexact line searches.

The Hestenes-Stiefel and Polak-Ribière methods appear to perform very similarly in practice, and are to be preferred over the Fletcher-Reeves method. Nevertheless, in a remarkably laborious paper, Powell (1984) was able to show that the Polak-Ribière method

with exact line searches can cycle infinitely without approaching a solution point. The same result applies to the Hestenes-Stiefel method since the two methods are identical when  $\langle g_k, d_{k-1} \rangle = 0$ , which holds when line searches are exact. Since the steplength of Powell's example would probably be accepted by any practical line search, it appears unlikely that a satisfactory global convergence result can be found for the Polak-Ribière and Hestenes-Stiefel methods. In contrast, Al-Baali's convergence result for the less efficient Fletcher-Reeves method, is very satisfactory. This disconcerting state of affairs motivated the present study.

In this paper we will consider various choices of  $\beta_k$  and various line search strategies that result in globally convergent methods. In §2 we describe the approach used in our analysis, and summarize some of the previous work in the area. §3 establishes global convergence for the class of methods with  $|\beta_k| \leq \beta_k^{\text{FR}}$ , and describes a modification of the Polak-Ribière formula. In §4 we consider methods that use only non-negative values for  $\beta_k$ , and which are, in some sense, related to the Polak-Ribière method. In particular, we show that a suggestion of Powell (1985), to set  $\beta_k = \max\{\beta_k^{\text{FR}}, 0\}$ , results in global convergence, even for inexact line searches. Further remarks on the convergence results are made in §5, and the results of some numerical experiments are presented in §6.

We note that this paper does not study the rate of convergence of conjugate gradient methods. For some results on this subject see Crowder and Wolfe (1972), Cohen (1972), Powell (1976), Baptist and Stoer (1977) and Stoer (1977).

## 2. Preliminaries.

Some important global convergence results for conjugate gradient methods have been given by Polak and Ribière (1969), Zoutendijk (1970), Powell (1984) and Al-Baali (1985). In this section we will see that the underlying approach used for these analyses is essentially the same, and we will describe it in detail, since it is also the basis for the results presented in this paper. Before doing so, we describe our notation, state the assumptions we make about the objective function, and consider the line search strategy.

**Notation and definitions.** We denote the starting point by  $x_1$ , and define  $s_k := x_{k+1} - x_k$ , and  $y_k := g_{k+1} - g_k$ . We say that  $d_k$  is a *descent direction* if  $\langle g_k, d_k \rangle < 0$ . We will also make use of the angle  $\theta_k$  between  $-g_k$  and  $d_k$ :

$$\cos \theta_k := -\langle g_k, d_k \rangle / \|g_k\| \|d_k\|. \quad (2.1)$$

The Fletcher-Reeves, Polak-Ribière and Hestenes-Stiefel methods will be abbreviated as FR, PR and HS, respectively. For a derivation of these methods and a discussion of some of their properties see Gill, Murray and Wright (1981), and Fletcher (1987).

**Assumptions 2.1** (i) *The level set  $\mathcal{L} := \{x : f(x) \leq f(x_1)\}$  is bounded.* (ii) *In some neighborhood  $\mathcal{N}$  of  $\mathcal{L}$ , the objective function  $f$  is continuously differentiable, and its gradient is Lipschitz continuous, i.e., there exists a constant  $L > 0$  such that*

$$\|g(x) - g(\tilde{x})\| \leq L\|x - \tilde{x}\|, \quad (2.2)$$

for all  $x, \tilde{x} \in \mathcal{N}$ .

Note that these assumptions imply that there is a constant  $\bar{\gamma}$  such that

$$\|g(x)\| \leq \bar{\gamma}, \text{ for all } x \in \mathcal{L}. \quad (2.3)$$

Let us now turn our attention to the line search. An efficient strategy, studied by Wolfe (1969), consists in accepting a positive steplength  $\alpha_k$  if it satisfies the two conditions:

$$f(x_k + \alpha_k d_k) \leq f(x_k) + \sigma_1 \alpha_k \langle g_k, d_k \rangle \quad (2.4)$$

$$\langle g(x_k + \alpha_k d_k), d_k \rangle \geq \sigma_2 \langle g_k, d_k \rangle, \quad (2.5)$$

where  $0 < \sigma_1 < \sigma_2 < 1$ . We will sometimes refer also to more ideal line search conditions. To this end let us define the following strategy: a positive steplength  $\alpha_k$  is accepted if

$$f(x_k + \alpha_k d_k) \leq f(x_k + \hat{\alpha}_k d_k), \quad (2.6)$$

where  $\hat{\alpha}_k$  is the *smallest* positive stationary point of the function  $\xi_k(\alpha) := f(x_k + \alpha d_k)$ . Assumptions 2.1 ensure that  $\hat{\alpha}_k$  exists. Note that both the first local minimizer as well as the global minimizer of  $f$  along the search direction satisfy (2.6).

Any of these line search strategies is sufficient to establish the following very useful result.

**Theorem 2.1** *Suppose that Assumptions 2.1 hold, and consider any iteration of the form (1.3), where  $d_k$  is a descent direction and  $\alpha_k$  satisfies one of the following line search conditions: (i) the Wolfe conditions (2.4)-(2.5), or (ii) the ideal line search condition (2.6). Then*

$$\sum_{k \geq 1} \cos^2 \theta_k \|g_k\|^2 < \infty. \quad (2.7)$$

This result was essentially proved by Zoutendijk (1970) and Wolfe (1969 and 1971). We shall call equation (2.7), the *Zoutendijk condition*.

We can now describe the basic ideas used for the convergence analysis. The first results, by Polak and Ribière (1969) and Zoutendijk (1970), assume exact line searches. The term *exact line search* can be ambiguous. Sometimes, it implies that a one-dimensional minimizer is found, but often it simply means that the orthogonality condition

$$\langle g_k, d_{k-1} \rangle = 0, \quad (2.8)$$

is satisfied. Throughout the paper we will indicate in detail the conditions required of the line search. Let us suppose that  $d_{k-1}$  is a descent direction and that the line search satisfies Zoutendijk's condition and condition (2.8). From (1.2) and (2.8) we have that

$$\cos \theta_k = \|g_k\| / \|d_k\|, \quad (2.9)$$

which shows that  $d_k$  is a descent direction. Substituting this relation in Zoutendijk's condition (2.7) we obtain

$$\sum_{k \geq 1} \frac{\|g_k\|^4}{\|d_k\|^2} < \infty. \quad (2.10)$$

If one can show that  $\{\|d_k\|/\|g_k\|\}$  is bounded, which means that  $\{\cos \theta_k\}$  is bounded away from zero, then (2.10) immediately gives

$$\lim_{k \rightarrow \infty} g_k = 0. \quad (2.11)$$

This is done by Polak and Ribière (1969) for their method, assuming that  $f$  is *strongly convex*, i.e.,  $\langle g(x) - g(\tilde{x}), x - \tilde{x} \rangle \geq c \|x - \tilde{x}\|^2$ , for some positive constant  $c$  and for all  $x$  and  $\tilde{x}$  in  $\mathcal{L}$ .

For general functions, however, it is usually impossible to bound  $\{\|d_k\|/\|g_k\|\}$  *a priori*, and only a weaker result than (2.11) can be obtained, namely

$$\liminf_{k \rightarrow \infty} \|g_k\| = 0. \quad (2.12)$$

To obtain this result one proceeds by contradiction. Suppose that (2.12) does not hold, which means that the gradients remain bounded away from zero: there exists  $\gamma > 0$  such that

$$\|g_k\| \geq \gamma, \quad (2.13)$$

for all  $k \geq 1$ . Then (2.10) implies that

$$\sum_{k \geq 1} \frac{1}{\|d_k\|^2} < \infty. \quad (2.14)$$

We conclude that the iteration can fail only if  $\|d_k\| \rightarrow \infty$  sufficiently rapidly. The method of proof used by Zoutendijk consists in showing that, if (2.13) holds, then  $\|d_k\|^2$  can grow at most linearly, i.e.,

$$\|d_k\|^2 \leq c k,$$

for some constant  $c$ . This contradicts (2.14), proving (2.12).

The analysis for *inexact* line searches that satisfy Zoutendijk's condition can proceed along the same lines if one can show that the iteration satisfies

$$\cos \theta_k \geq c \|g_k\|/\|d_k\|, \quad (2.15)$$

for some positive constant  $c$ . Then, this relation can be used instead of (2.9) to give (2.10), and the rest of the analysis is as in the case of exact line searches.

Al-Baali (1985) shows that the FR method gives (2.15) if the steplength satisfies the *strong* Wolfe conditions:

$$f(x_k + \alpha_k d_k) \leq f(x_k) + \sigma_1 \alpha_k \langle g_k, d_k \rangle \quad (2.16)$$

$$|\langle g(x_k + \alpha_k d_k), d_k \rangle| \leq -\sigma_2 \langle g_k, d_k \rangle, \quad (2.17)$$



where  $0 < \sigma_1 < \sigma_2 < 1$ . In fact, it is necessary to require that  $\sigma_2 < \frac{1}{2}$  for the result to hold. He thus shows that (2.12) holds for the FR method.

Al-Baali's result is remarkable in another respect too. By establishing (2.15), which by (2.1) is equivalent to

$$\langle g_k, d_k \rangle \leq -c \|g_k\|^2, \quad (2.18)$$

he proved that the FR method using the strong Wolfe conditions (with  $\sigma_2 < \frac{1}{2}$ ) always generates descent directions. Prior to this result it was believed that it was necessary to enforce the descent condition while doing the line search.

In this paper we use the approach described above to establish the global convergence of various algorithms with inexact line searches. As we do so, we will repeatedly encounter (2.18), which appears to be a natural way of guaranteeing descent for conjugate gradient methods. We call (2.18) the *sufficient descent condition*. The first class of methods we consider, in §3, are related to the FR method. We show that any method of the form (1.2)-(1.3) is globally convergent if  $\beta_k$  satisfies  $|\beta_k| \leq \beta_k^{\text{FR}}$ . The result readily suggests a new implementation of the PR method that preserves its efficiency and assures its convergence.

In §4, we study methods with  $\beta_k \geq 0$  that are, in some sense, related to the PR method. A particular case is the following adaptation of the PR method, which consists in restricting  $\beta_k$  to positive values: we let

$$\beta_k = \max\{\beta_k^{\text{PR}}, 0\}. \quad (2.19)$$

The motivation for this strategy arises from Powell's analysis of the PR method. Powell (1984) assumes that the line search always finds the first stationary point, and shows that there is a twice continuously differentiable function and a starting point such that the sequence of gradients generated by the PR method stays bounded away from zero. Since Powell's example requires that some consecutive search directions become almost contrary, and since this can only be achieved (in the case of exact line searches) when  $\beta_k < 0$ , Powell (1985) suggests modifying the PR method as in (2.19). In §4 we show that this choice of  $\beta_k$  does indeed result in global convergence, both for exact and inexact line searches. Moreover we show that the analysis also applies to a family of methods with  $\beta_k \geq 0$  that share a common property with the PR method – we call this *Property (\*)*.

### 3. Iterations constrained by the FR method.

In this section we will see that it is possible to obtain global convergence if the parameter  $\beta_k$  is appropriately bounded in magnitude. We consider a method of the form (1.2)-(1.3), where  $\beta_k$  is any scalar such that

$$|\beta_k| \leq \beta_k^{\text{FR}}, \quad (3.1)$$

for all  $k \geq 2$ , and where the steplength satisfies the strong Wolfe conditions (2.16)-(2.17) with  $\sigma_2 < \frac{1}{2}$ . Note that Zoutendijk's result, Theorem 2.1, holds in this case, since the strong Wolfe conditions imply the Wolfe conditions (2.4)-(2.5). The next two results are very based upon the work of Al-Baali (1985) for the FR method, and are slightly stronger than those given by Touati-Ahmed and Storey (1990).

**Lemma 3.1** Suppose that Assumptions 2.1 hold. Consider any method of the form (1.2)-(1.3), where  $\beta_k$  satisfies (3.1), and where the steplength satisfies the Wolfe condition (2.17) with  $\sigma_2 < \frac{1}{2}$ . Then, the method generates descent directions  $d_k$  satisfying

$$-\sum_{j=0}^{k-1} \sigma_2^j \leq \frac{\langle g_k, d_k \rangle}{\|g_k\|^2} \leq -2 + \sum_{j=0}^{k-1} \sigma_2^j, \quad k = 1, \dots \quad (3.2)$$

**Proof.** The proof is by induction. The result clearly holds for  $k = 1$  since all three terms equal  $-1$ . Assume that (3.2) holds for some  $k \geq 1$ . This implies that  $\langle g_k, d_k \rangle < 0$ , since

$$-2 + \sum_{j=0}^{k-1} \sigma_2^j < -2 + \frac{1}{1 - \sigma_2} = \frac{2\sigma_2 - 1}{1 - \sigma_2} < 0, \quad (3.3)$$

by the condition  $\sigma_2 < \frac{1}{2}$ . From (1.2) and (1.4) we have

$$\frac{\langle g_{k+1}, d_{k+1} \rangle}{\|g_{k+1}\|^2} = -1 + \beta_{k+1} \frac{\langle g_{k+1}, d_k \rangle}{\|g_{k+1}\|^2} = -1 + \frac{\beta_{k+1}}{\beta_{k+1}^{\text{FR}}} \frac{\langle g_k, d_k \rangle}{\|g_k\|^2}. \quad (3.4)$$

Using the line search condition (2.17) we have

$$|\beta_{k+1} \langle g_{k+1}, d_k \rangle| \leq -\sigma_2 |\beta_{k+1}| \langle g_k, d_k \rangle,$$

which together with (3.4) gives

$$-1 + \sigma_2 \frac{|\beta_{k+1}|}{\beta_{k+1}^{\text{FR}}} \frac{\langle g_k, d_k \rangle}{\|g_k\|^2} \leq \frac{\langle g_{k+1}, d_{k+1} \rangle}{\|g_{k+1}\|^2} \leq -1 - \sigma_2 \frac{|\beta_{k+1}|}{\beta_{k+1}^{\text{FR}}} \frac{\langle g_k, d_k \rangle}{\|g_k\|^2}.$$

From the left hand side of the induction hypothesis (3.2) we obtain

$$-1 - \sigma_2 \frac{|\beta_{k+1}|}{\beta_{k+1}^{\text{FR}}} \sum_{j=0}^{k-1} \sigma_2^j \leq \frac{\langle g_{k+1}, d_{k+1} \rangle}{\|g_{k+1}\|^2} \leq -1 + \sigma_2 \frac{|\beta_{k+1}|}{\beta_{k+1}^{\text{FR}}} \sum_{j=0}^{k-1} \sigma_2^j.$$

Using the bound (3.1), and since  $\sum_{j=0}^{k-1} \sigma_2^{j+1} = \sum_{j=0}^k \sigma_2^j - 1$ , we conclude that (3.2) holds for  $k + 1$ .  $\square$

Lemma 3.1 achieves three objectives: (i) it shows that all search directions are descent directions, and the upper bound in (3.2) shows that the sufficient descent condition (2.18) holds; (ii) the bounds on  $\langle g_k, d_k \rangle$  impose a limit on how fast  $\|d_k\|$  can grow when the gradients are not small, as we will see this in the next theorem; (iii) from (2.1) and (3.2) we see that there are positive constants  $c_1$  and  $c_2$  such that

$$c_1 \frac{\|g_k\|}{\|d_k\|} \leq \cos \theta_k \leq c_2 \frac{\|g_k\|}{\|d_k\|}. \quad (3.5)$$

Therefore, for the FR method or any method with  $|\beta_k| \leq \beta_k^{\text{FR}}$ , we have that  $\cos \theta_k$  is proportional to  $\|g_k\|/\|d_k\|$ . We will make good use of this fact later on.

**Theorem 3.2** Suppose that Assumptions 2.1 hold. Consider any method of the form (1.2)-(1.3), where  $\beta_k$  satisfies (3.1), and where the steplength satisfies the strong Wolfe conditions (2.16)-(2.17), with  $0 < \sigma_1 < \sigma_2 < \frac{1}{2}$ . Then

$$\liminf_{k \rightarrow \infty} \|g_k\| = 0.$$

**Proof.** From (2.17) and Lemma 3.1 we have

$$\begin{aligned} |\langle g_k, d_{k-1} \rangle| &\leq -\sigma_2 \langle g_{k-1}, d_{k-1} \rangle \\ &\leq \sigma_2 \sum_{j=0}^{k-2} \sigma_2^j \|g_{k-1}\|^2 \\ &\leq \frac{\sigma_2}{1 - \sigma_2} \|g_{k-1}\|^2. \end{aligned}$$

Thus, from (1.2) and (3.1)

$$\begin{aligned} \|d_k\|^2 &\leq \|g_k\|^2 + 2|\beta_k| |\langle g_k, d_{k-1} \rangle| + \beta_k^2 \|d_{k-1}\|^2 \\ &\leq \|g_k\|^2 + \frac{2\sigma_2}{1 - \sigma_2} |\beta_k| \|g_{k-1}\|^2 + \beta_k^2 \|d_{k-1}\|^2 \\ &\leq \left( \frac{1 + \sigma_2}{1 - \sigma_2} \right) \|g_k\|^2 + \beta_k^2 \|d_{k-1}\|^2. \end{aligned}$$

Applying this relation repeatedly, defining  $\hat{\sigma} := (1 + \sigma_2)/(1 - \sigma_2) \geq 1$ , and using the condition  $|\beta_k| \leq \beta_k^{\text{FR}}$ , we have

$$\begin{aligned} \|d_k\|^2 &\leq \hat{\sigma} \|g_k\|^2 + \beta_k^2 (\hat{\sigma} \|g_{k-1}\|^2 + \beta_{k-1}^2 \|d_{k-2}\|^2) \\ &\leq \hat{\sigma} \|g_k\|^4 \sum_{j=1}^k \|g_j\|^{-2}. \end{aligned}$$

Let us now assume that  $\|g_k\| \geq \gamma > 0$  for all  $k$ . This implies, by (2.3), that

$$\|d_k\|^2 \leq \frac{\hat{\sigma} \gamma^4}{\gamma^2} k. \quad (3.6)$$

We now follow the reasoning described in §2. From the left inequality in (3.5) and Zoutendijk's result (2.7) we obtain (2.10). If the gradients are bounded away from zero, (2.10) implies (2.14). We conclude the proof by noting that (3.6) and (2.14) are incompatible.  $\square$

This theorem suggests the following globally convergent modification of the PR method. It differs from that considered by Tonati-Ahmed and Storey (1990) in that it allows for negative values of  $\beta_k$ . For all  $k \geq 2$  let

$$\beta_k = \begin{cases} -\beta_k^{\text{FR}} & \text{if } \beta_k^{\text{PR}} < -\beta_k^{\text{FR}} \\ \beta_k^{\text{PR}} & \text{if } |\beta_k^{\text{PR}}| \leq \beta_k^{\text{FR}} \\ \beta_k^{\text{FR}} & \text{if } \beta_k^{\text{PR}} > \beta_k^{\text{FR}}. \end{cases} \quad (3.7)$$

This strategy avoids one of the main disadvantages of the FR method, as we will now discuss.

We have observed in numerical tests that the FR method with inexact line searches sometimes slows down away from the solution: the steps become very small and this behavior can continue for a very large number of iterations, unless the method is restarted. This behavior was observed earlier by Powell (1977), who provides an explanation, under the assumption of exact line searches. It turns out that his argument can be extended to the case of inexact line searches, due to (3.5). The argument is as follows. Suppose that at the iteration  $k$  an unfortunate search direction is generated, such that  $\cos \theta_k \approx 0$ , and that  $x_{k+1} \approx x_k$ . Thus  $\|g_{k+1}\| \approx \|g_k\|$ , and

$$\beta_{k+1}^{\text{FR}} \approx 1. \quad (3.8)$$

Moreover, by (3.5),

$$\|g_{k+1}\| \approx \|g_k\| \ll \|d_k\|.$$

From this relation, (3.8) and (1.2) we see that  $\|d_{k+1}\| \approx \|d_k\| \gg \|g_{k+1}\|$ , which by (3.5) implies that  $\cos \theta_{k+1} \approx 0$ . The argument can therefore start all over again. In §6 we give a numerical example demonstrating this behavior.

The PR method would behave quite differently from the FR method in this situation. If  $g_{k+1} \approx g_k$ , then  $\beta_{k+1}^{\text{PR}} \approx 0$ , so that by (1.2) and (3.5)  $\cos \theta_{k+1} \gg \cos \theta_k$ . Thus the PR method would recover from that situation. Let us now consider the behavior of method (3.7) in these circumstances. We have seen that  $\beta_{k+1}^{\text{FR}} \approx 1$ , and  $\beta_{k+1}^{\text{PR}} \approx 0$ , in this case. The method (3.7) will thus set  $\beta_{k+1} = \beta_{k+1}^{\text{PR}}$ , as desired. It is reassuring that the modification (3.7), which falls back on the FR method to ensure global convergence, avoids the inefficiencies of this method.

The previous discussion highlights a property of the PR method that is not shared by the FR method: when the step is small,  $\beta_k^{\text{PR}}$  will be small. This property is essential for the analysis given in the next section, where a method that possesses it, will be said to have *Property (\*)*.

It is natural to ask if the bound  $|\beta_k| \leq \beta_k^{\text{FR}}$  can be replaced by

$$|\beta_k| \leq c \beta_k^{\text{FR}}, \quad (3.9)$$

where  $c > 1$  is some suitable constant. We have not been able to establish global convergence in this case (although, by modifying Lemma 3.1, one can show that the descent property of the search directions can still be obtained provided  $\sigma_2 < 1/(2c)$ ). In fact, one can prove the following negative result.

**Proposition 3.3** *Consider the method (1.2)-(1.3), with a line search that always chooses the first positive stationary point of  $\xi_k(\alpha) := f(x_k + \alpha d_k)$ . There exists a twice continuously differentiable objective function of three variables, a starting point and a choice of  $\beta_k$  satisfying (3.9) for some constant  $c > 1$ , such that the sequence of gradients  $\{\|g_k\|\}$  is bounded away from zero.*

**Proof.** The objective function is taken from the fourth example of Powell (1984). It is twice continuously differentiable. For this function, there is a starting point from which the PR method with a line search providing the first stationary point fails to converge, in the sense that  $\|g_k\| \geq \gamma > 0$  for all  $k$ . Therefore using (2.3) we have for all  $k \geq 2$

$$|\beta_k^{\text{PR}}| \leq \frac{2\bar{\gamma}^2}{\gamma^2}.$$

Now, suppose that we computed (but not used)  $\beta_k^{\text{FR}}$ . We would see that for all  $k \geq 2$

$$\beta_k^{\text{FR}} \geq \frac{\gamma^2}{\bar{\gamma}^2}.$$

Combining the two inequalities we obtain

$$|\beta_k^{\text{PR}}| \leq \frac{2\bar{\gamma}^4}{\gamma^4} \beta_k^{\text{FR}}.$$

Therefore, if the constant  $c$  in (3.9) is chosen larger than  $2\bar{\gamma}^4/\gamma^4$ , the PR parameter  $\beta_k^{\text{PR}}$  in Powell's example would always satisfy (3.9).  $\square$

We end this section by making an observation about the restart criterion of Powell (1977). Even though this criterion was designed to ensure the convergence of Beale's method, we will apply it to the PR method, and see that it has some of the flavor of the modifications described in this section. Powell (1977) suggests to restart if the following inequality is violated,

$$|\langle g_k, g_{k-1} \rangle| \leq \nu \|g_{k-1}\|^2,$$

where  $\nu$  is a small positive constant. (Powell actually uses  $g_k$  instead of  $g_{k-1}$  in the right hand side, but one can argue for either choice.) From (1.4) and (1.5),

$$\beta_k^{\text{PR}} = \beta_k^{\text{FR}} - \frac{\langle g_k, g_{k-1} \rangle}{\|g_{k-1}\|^2}.$$

Applying the restart criterion to the PR method we see that a restart is *not* necessary as long as

$$\beta_k^{\text{FR}} - \nu \leq \beta_k^{\text{PR}} \leq \beta_k^{\text{FR}} + \nu.$$

Once more,  $\beta_k^{\text{FR}}$  appears as a measure of the adequacy of  $\beta_k^{\text{PR}}$ , but this measure is quite different from (3.1). In the next section we will view Powell's restart criterion from a somewhat different angle.

#### 4. Methods related to the PR method with non-negative $\beta_k$ .

We now turn our attention to methods with  $\beta_k \geq 0$  for all  $k$ . In §2 we mentioned that a motivation for placing this restriction comes from the example of Powell, in which the PR method cycles without obtaining the solution. Another reason for keeping  $\beta_k \geq 0$  is that it allows us to easily enforce the descent property of the algorithm, as we will now discuss.

Let us consider the iteration (1.2)-(1.3) with any  $\beta_k \geq 0$ . We will require the sufficient descent condition

$$\langle g_k, d_k \rangle \leq -\sigma_3 \|g_k\|^2, \quad (4.1)$$

for some  $0 < \sigma_3 \leq 1$  and for all  $k \geq 1$ . In contrast to the FR method, the strong Wolfe conditions (2.16)-(2.17) no longer ensure (4.1). Note, from (1.2), that

$$\langle g_k, d_k \rangle = -\|g_k\|^2 + \beta_k \langle g_k, d_{k-1} \rangle. \quad (4.2)$$

Therefore, to obtain descent for an inexact line search algorithm, one needs to ensure that the last term is not too large. Suppose that we perform a line search along  $d_{k-1}$ , enforcing the Wolfe (or strong Wolfe) conditions, to obtain  $x_k$ . If  $\langle g_k, d_{k-1} \rangle \leq 0$ , the non-negativity of  $\beta_k$  implies that the sufficient descent condition (4.1) holds. Moreover if (4.1) is not satisfied then  $\langle g_k, d_{k-1} \rangle > 0$ , which means that a one-dimensional minimizer has been bracketed. In this case it is easy to apply a line search algorithm, such as that given by Lemaréchal (1981), Fletcher (1987) or Moré and Thuente (1990), to reduce  $|\langle g_k, d_{k-1} \rangle|$  sufficiently and obtain (4.1). This will be discussed in detail in §6.

We now prove a global convergence result for methods which are related to the PR method, but that allow only non-negative values of  $\beta_k$ . The idea of our analysis is simple, but is somewhat concealed in the proofs. We establish the results by contradiction, assuming that the gradients are bounded away from zero:

$$\text{for some } \gamma > 0, \|g_k\| \geq \gamma, \text{ for all } k \geq 1. \quad (4.3)$$

Lemma 4.1 shows that in this case the direction of search changes slowly, asymptotically, and Lemma 4.2 proves that a certain fraction of the steps are not too small. In Theorem 4.3 we show that these two results contradict the assumption that the iterates stay in the bounded level set  $\mathcal{L}$ . We conclude that a subsequence of the iterates converges to a stationary point.

For the results that follow, we do not specify a particular line search strategy. We only assume that the line search satisfies the following three properties: (i) all iterates remain in the level set  $\mathcal{L}$  defined in Assumptions 2.1:

$$\{x_k\} \subset \mathcal{L}; \quad (4.4)$$

(ii) the Zoutendijk condition (2.7) holds; and (iii) the sufficient descent condition (4.1) holds. We mentioned in §2 that the Wolfe line search, as well as the ideal line search (2.6), ensure Zoutendijk's condition and reduce  $f$  at each step, which implies (4.4). An exact line search satisfies the sufficient descent condition (4.1), because in this case  $\langle g_k, d_k \rangle = -\|g_k\|^2$ , and in §6 we describe an inexact line search procedure that satisfies the Wolfe conditions and (4.1) when  $\beta_k \geq 0$ . Therefore the results of this section apply to both ideal and practical line searches.

For the rest of the section, we assume that convergence does not occur in a finite number of steps, i.e.,  $g_k \neq 0$  for all  $k$ .

**Lemma 4.1** *Suppose that Assumptions 2.1 hold. Consider the method (1.2)-(1.3), with  $\beta_k \geq 0$ , and with any line search satisfying both the Zoutendijk condition (2.7) and the sufficient descent condition (4.1). If (4.3) holds, then  $d_k \neq 0$  and*

$$\sum_{k \geq 2} \|u_k - u_{k-1}\|^2 < \infty, \quad (4.5)$$

where  $u_k := d_k / \|d_k\|$ .

**Proof.** First, note that  $d_k \neq 0$ , for otherwise (4.1) would imply  $g_k = 0$ . Therefore,  $u_k$  is well defined. Now, let us define  $r_k := -g_k / \|d_k\|$  and  $\delta_k := \beta_k \|d_{k-1}\| / \|d_k\|$ . From (1.2), we have for  $k \geq 2$ :

$$u_k = r_k + \delta_k u_{k-1}, \quad (4.6)$$

which gives both

$$\langle r_k, u_k \rangle = \|r_k\|^2 + \delta_k \langle r_k, u_{k-1} \rangle, \quad (4.7)$$

$$1 = \|r_k\|^2 + \delta_k^2 + 2\delta_k \langle r_k, u_{k-1} \rangle. \quad (4.8)$$

Using (4.6) and (4.8), we obtain

$$\begin{aligned} \|u_k - u_{k-1}\|^2 &= \|r_k\|^2 + (\delta_k - 1)^2 + 2(\delta_k - 1)\langle r_k, u_{k-1} \rangle \\ &= 2(1 - \delta_k - \langle r_k, u_{k-1} \rangle) \\ &= 2\left(1 - \delta_k^2 - (1 + \delta_k)\langle r_k, u_{k-1} \rangle\right) / (1 + \delta_k), \end{aligned}$$

where the last step was obtained by multiplying and dividing by  $(1 + \delta_k)$ , which is nonzero since  $\beta_k \geq 0$ . Next, by using (4.8), (4.7) and the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} \|u_k - u_{k-1}\|^2 &= 2\left(\|r_k\|^2 - (1 - \delta_k)\langle r_k, u_{k-1} \rangle\right) / (1 + \delta_k) \\ &= 2\langle r_k, u_k - u_{k-1} \rangle / (1 + \delta_k) \\ &\leq 2\|r_k\| \|u_k - u_{k-1}\| / (1 + \delta_k). \end{aligned}$$

Therefore, since  $1 + \delta_k \geq 1$ , we have for all  $k \geq 2$ ,

$$\|u_k - u_{k-1}\| \leq 2\|r_k\|. \quad (4.9)$$

Now, by (2.1) and (4.1), we have

$$\cos \theta_k \geq \sigma_3 \frac{\|g_k\|}{\|d_k\|}.$$

This relation and Zoutendijk's condition (2.7) imply

$$\sum_{k \geq 2} \frac{\|g_k\|^4}{\|d_k\|^2} = \sum_{k \geq 2} \|r_k\|^2 \|g_k\|^2 < \infty.$$

Using (4.3), we obtain

$$\sum_{k \geq 2} \|r_k\|^2 < \infty,$$

which together with (4.9) completes the proof.  $\square$

Of course, condition (4.5) does not imply the convergence of the sequence  $\{u_k\}$ , but shows that the search directions  $u_k$  change slowly, asymptotically.

Lemma 4.1 applies to any choice of  $\beta_k \geq 0$ . To proceed we need to require, in addition, that  $\beta_k$  be small when the step is small. We saw in §3 that the PR method possesses this property and that it prevents the inefficient behavior of the FR method from occurring. We now state this property formally.

**Property (\*).** Consider a method of the form (1.2)-(1.3), and suppose that

$$0 < \gamma \leq \|g_k\| \leq \bar{\gamma}, \quad (4.10)$$

for all  $k \geq 1$ . Under this assumption we say that the method has Property (\*) if there exist constants  $b > 1$  and  $\lambda > 0$  such that for all  $k$ :

$$|\beta_k| \leq b, \quad (4.11)$$

and

$$\|s_{k-1}\| \leq \lambda \implies |\beta_k| \leq \frac{1}{2b}. \quad (4.12)$$

It is easy to see that under Assumptions 2.1 the PR and HS methods have Property (\*). For the PR method, using the constants  $\gamma$  and  $\bar{\gamma}$  in (4.10), we can choose  $b := 2\bar{\gamma}^2/\gamma^2$  and  $\lambda := \gamma^2/(2L\bar{\gamma}b)$ . Then, we have from (1.5) and (4.10)

$$|\beta_k^{\text{PR}}| \leq \frac{(\|g_k\| + \|g_{k-1}\|)\|g_k\|}{\|g_{k-1}\|^2} \leq \frac{2\bar{\gamma}^2}{\gamma^2} = b,$$

and when  $\|s_{k-1}\| \leq \lambda$  we have from (2.2)

$$|\beta_k^{\text{PR}}| \leq \frac{\|y_{k-1}\|\|g_k\|}{\|g_{k-1}\|^2} \leq \frac{L\lambda\bar{\gamma}}{\gamma^2} = \frac{1}{2b}.$$

For the HS method, we assume that the descent condition (4.1) and the second Wolfe condition (2.5) are satisfied. Then

$$\begin{aligned} \langle d_{k-1}, y_{k-1} \rangle &= \langle d_{k-1}, g_k \rangle - \langle d_{k-1}, g_{k-1} \rangle \\ &\geq -(1 - \sigma_2) \langle g_{k-1}, d_{k-1} \rangle \\ &\geq (1 - \sigma_2) \sigma_3 \|g_{k-1}\|^2 \\ &\geq (1 - \sigma_2) \sigma_3 \gamma^2. \end{aligned}$$

Using this in (1.6) we obtain

$$|\beta_k^{\text{HS}}| \leq \frac{2\bar{\gamma}^2}{(1 - \sigma_2) \sigma_3 \gamma^2} =: b.$$



Now define  $\lambda := (1 - \sigma_2)\sigma_3\gamma^2/(2L\bar{\gamma}b)$ . Using (2.2) we see that if  $\|s_{k-1}\| \leq \lambda$ , then

$$|\beta_k^{\text{HS}}| \leq \frac{L\lambda\bar{\gamma}}{(1 - \sigma_2)\sigma_3\gamma^2} = \frac{1}{2b}.$$

It is clear that many other choices of  $\beta_k$  give rise to algorithms with Property (\*). For example, if  $\beta_k$  has Property (\*), so do  $|\beta_k|$  and  $\beta_k^+ := \max\{\beta_k, 0\}$ .

The next Lemma shows that if the gradients are bounded away from zero, and if the method has Property (\*), then a fraction of the steps cannot be too small. We let  $\mathbf{N}^*$  denote the set of positive integers, and for  $\lambda > 0$  we define  $\mathcal{K}^\lambda := \{i \in \mathbf{N}^* : i \geq 2, \|s_{i-1}\| > \lambda\}$ , i.e., the set of integers corresponding to steps that are larger than  $\lambda$ . We will need to consider groups of  $\Delta$  consecutive iterates, and for this purpose we define

$$\mathcal{K}_{k,\Delta}^\lambda := \{i \in \mathbf{N}^* : k \leq i \leq k + \Delta - 1, \|s_{i-1}\| > \lambda\}.$$

Let  $|\mathcal{K}_{k,\Delta}^\lambda|$  denote the number of elements of  $\mathcal{K}_{k,\Delta}^\lambda$  and let  $\lfloor \cdot \rfloor$  and  $\lceil \cdot \rceil$  denote respectively the floor and ceiling operators.

**Lemma 4.2** *Suppose that Assumptions 2.1 hold. Consider the method (1.2)-(1.3), with any line search satisfying (4.4), the Zoutendijk condition (2.7) and the sufficient descent condition (4.1), and assume that the method has Property (\*). Suppose also that (4.3) holds. Then there exists  $\lambda > 0$  such that, for any  $\Delta \in \mathbf{N}^*$  and any index  $k_0$ , there is a greater index  $k \geq k_0$  such that*

$$|\mathcal{K}_{k,\Delta}^\lambda| > \frac{\Delta}{2}.$$

**Proof.** We proceed by contradiction. Suppose that

$$\begin{cases} \text{for any } \lambda > 0, \text{ there exists } \Delta \in \mathbf{N}^* \text{ and } k_0 \text{ such that} \\ \text{for any } k \geq k_0, \text{ we have } |\mathcal{K}_{k,\Delta}^\lambda| \leq \frac{\Delta}{2}. \end{cases} \quad (4.13)$$

Assumptions 2.1, (4.4) and (4.3) imply that (4.10) holds. Since the method has Property (\*), there exists  $\lambda > 0$  and  $b > 1$  such that (4.11) and (4.12) hold for all  $k$ . For this  $\lambda$ , let  $\Delta$  and  $k_0$  be given by (4.13).

For any given index  $l \geq k_0 + 1$ , we have

$$\begin{aligned} \|d_l\|^2 &\leq (\|g_l\| + |\beta_l| \|d_{l-1}\|)^2 \\ &\leq 2\|g_l\|^2 + 2\beta_l^2 \|d_{l-1}\|^2 \\ &\leq 2\bar{\gamma}^2 + 2\beta_l^2 \|d_{l-1}\|^2, \end{aligned}$$

where the second inequality follows from the fact that, for any scalars  $a$  and  $b$ , we have  $2ab \leq a^2 + b^2$ , and hence  $(a + b)^2 \leq 2a^2 + 2b^2$ . By induction, we obtain

$$\|d_l\|^2 \leq c(1 + 2\beta_l^2 + 2\beta_l^2 2\beta_{l-1}^2 + \cdots + 2\beta_l^2 2\beta_{l-1}^2 \cdots 2\beta_{k_0}^2), \quad (4.14)$$

where  $c$  depends on  $\|d_{k_0-1}\|$ , but not on the index  $l$ . Let us consider a typical term in (4.14):

$$2\beta_l^2 2\beta_{l-1}^2 \cdots 2\beta_k^2, \quad (4.15)$$

where

$$k_0 \leq k \leq l. \quad (4.16)$$

We now divide the  $2(l - k + 1)$  factors of (4.15) into groups of  $2\Delta$  elements, i.e., if  $N := \lfloor (l - k + 1)/\Delta \rfloor$ , then (4.15) can be divided into  $N$  or  $N + 1$  groups, as follows:

$$(2\beta_{l_1}^2 \cdots 2\beta_{k_1}^2), \dots, (2\beta_{l_N}^2 \cdots 2\beta_{k_N}^2), \quad (4.17)$$

and possibly

$$(2\beta_{l_{N+1}}^2 \cdots 2\beta_{k_N}^2), \quad (4.18)$$

where  $l_i = l - (i - 1)\Delta$ , for  $i = 1, \dots, N + 1$ , and  $k_i = l_{i+1} + 1$ , for  $i = 1, \dots, N$ . Note from (4.16) that  $k_i \geq k_0$ , for  $i = 1, \dots, N$ , so that we can apply (4.13) for  $k = k_i$ . We thus have

$$p_i := |\mathcal{K}_{k_i, \Delta}^\lambda| \leq \frac{\Delta}{2}, \quad i = 1, \dots, N. \quad (4.19)$$

This means that in the range  $[k_i, k_i + \Delta - 1]$  there are exactly  $p_i$  indices  $j$  such that  $\|s_{j-1}\| > \lambda$ , and thus there are  $(\Delta - p_i)$  indices with  $\|s_{j-1}\| \leq \lambda$ . Using this fact, (4.11) and (4.12), we examine a typical factor in (4.17),

$$\begin{aligned} 2\beta_{l_i}^2 \cdots 2\beta_{k_i}^2 &\leq 2^{\Delta} b^{2p_i} \left( \frac{1}{2b} \right)^{2(\Delta - p_i)} \\ &= 2^{\Delta - 2\Delta + 2p_i} b^{2p_i - 2\Delta + 2p_i} \\ &= (2b^2)^{2p_i - \Delta} \\ &\leq 1, \end{aligned}$$

since by (4.19)  $2p_i - \Delta \leq 0$  and  $2b^2 \geq 1$ . Therefore each of the factors in (4.17) is less than or equal to 1, and so is their product. For the last group of factors, given in (4.18), we simply use (4.11)

$$2\beta_{l_{N+1}}^2 \cdots 2\beta_{k_N}^2 < (2b^2)^\Delta.$$

We conclude that each term on the right hand side of (4.14) is bounded by  $(2b^2)^\Delta$ , and as a result we have

$$\|d_l\|^2 \leq c(l - k_0 + 2), \quad (4.20)$$

for a certain positive constant  $c$  independent of  $l$ . In other words, we have shown that  $\|d_l\|^2$  grows at most linearly, and we now obtain a contradiction as described in §2. Recalling that (4.1) implies condition (2.15) and using the Zoutendijk condition (2.7), we obtain that

$$\gamma^4 \sum_{k \geq 1} \frac{1}{\|d_k\|^2} \leq \sum_{k \geq 1} \frac{\|g_k\|^4}{\|d_k\|^2} < \infty.$$

This contradicts (4.20), concluding the proof.  $\square$

**Theorem 4.3** *Suppose that Assumptions 2.1 hold. Consider the method (1.2)-(1.3) with the following three properties: (i)  $\beta_k \geq 0$  for all  $k$ ; (ii) the line search satisfies (4.4), the Zoutendijk condition (2.7) and the sufficient descent condition (4.1); (iii) Property (\*) holds. Then  $\liminf \|g_k\| = 0$ .*

**Proof.** We proceed by contradiction, assuming (4.3). Therefore, the conditions of Lemma 4.1 and Lemma 4.2 hold. Defining  $u_i := d_i/\|d_i\|$ , as before, we have for any two indices  $l, k$ , with  $l \geq k$ :

$$\begin{aligned} x_l - x_{k-1} &= \sum_{i=k}^l \|s_{i-1}\| u_{i-1} \\ &= \sum_{i=k}^l \|s_{i-1}\| u_{k-1} + \sum_{i=k}^l \|s_{i-1}\| (u_{i-1} - u_{k-1}). \end{aligned}$$

Taking norms,

$$\sum_{i=k}^l \|s_{i-1}\| \leq \|x_l - x_{k-1}\| + \sum_{i=k}^l \|s_{i-1}\| \|u_{i-1} - u_{k-1}\|.$$

By (4.4) and Assumptions 2.1 we have that the sequence  $\{x_k\}$  is bounded, and thus there exists a positive constant  $B$  such that  $\|x_k\| \leq B$ , for all  $k \geq 1$ . Thus

$$\sum_{i=k}^l \|s_{i-1}\| \leq 2B + \sum_{i=k}^l \|s_{i-1}\| \|u_{i-1} - u_{k-1}\|. \quad (4.21)$$

Let  $\lambda > 0$  be given by Lemma 4.2. Following the notation of this lemma, we define  $\Delta := \lceil 8B/\lambda \rceil$ . By Lemma 4.1, we can find an index  $k_0$  such that

$$\sum_{i \geq k_0} \|u_i - u_{i-1}\|^2 \leq \frac{1}{4\Delta}. \quad (4.22)$$

With this  $\Delta$  and  $k_0$ , Lemma 4.2 gives an index  $k \geq k_0$  such that

$$|\mathcal{K}_{k,\Delta}^\lambda| > \frac{\Delta}{2}. \quad (4.23)$$

Next, for any index  $i \in [k, k + \Delta - 1]$ , we have by the Cauchy-Schwarz inequality and (4.22)

$$\begin{aligned} \|u_{i-1} - u_{k-1}\| &\leq \sum_{j=k}^{i-1} \|u_j - u_{j-1}\| \\ &\leq (i-k)^{1/2} \left( \sum_{j=k}^{i-1} \|u_j - u_{j-1}\|^2 \right)^{1/2} \\ &\leq \Delta^{1/2} \left( \frac{1}{4\Delta} \right)^{1/2} = \frac{1}{2}. \end{aligned}$$

Using this relation and (4.23) in (4.21), with  $l = k + \Delta - 1$ , we have

$$2B \geq \frac{1}{2} \sum_{i=k}^{k+\Delta-1} \|s_{i-1}\| > \frac{\lambda}{2} |\mathcal{K}_{k,\Delta}^\lambda| > \frac{\lambda\Delta}{4},$$

Thus  $\Delta < 8B/\lambda$ , which contradicts the definition of  $\Delta$ .  $\square$

Since the PR and HS methods have Property (\*), the previous theorem applies to them provided we restrict  $\beta_k$  to be non-negative. This suggests, among other, the following formulae:

$$\beta_k = \max\{\beta_k^{\text{PR}}, 0\}, \quad (4.24)$$

$$\beta_k = |\beta_k^{\text{PR}}|, \quad (4.25)$$

and the corresponding formulae for the HS method. Of particular interest are inexact line searches, such as the Wolfe search. We formally state the convergence result for (4.24) a choice of  $\beta_k$  suggested by Powell (1985).

**Corollary 4.4** *Suppose that Assumptions 2.1 hold. Consider the method (1.2)-(1.3) with  $\beta_k = \max\{\beta_k^{\text{PR}}, 0\}$ , and with a line search satisfying the Wolfe conditions (2.4)-(2.5) and the sufficient descent condition (4.1). Then  $\liminf \|g_k\| = 0$ .*

We conclude this section by noting the relationship between (4.24), which can be viewed as an automatic restarting procedure, and Powell's restarting criterion. The latter states that a restart is not needed as long as

$$|\langle g_k, g_{k-1} \rangle| \leq \nu \|g_k\|^2, \quad (4.26)$$

where we now used  $g_k$  and not  $g_{k-1}$  in the right hand side, and where  $\nu$  is a small positive constant. By (1.5) the condition  $\beta_k^{\text{PR}} \geq 0$  is equivalent to

$$\langle g_k, g_{k-1} \rangle \leq \|g_k\|^2.$$

Thus (4.24) can be viewed as a less restrictive restarting test than (4.26). It follows that the global convergence result of Corollary 4.4 also applies to the PR method with Powell's restart (4.26), provided  $\nu \leq 1$ .

## 5. Discussion.

In §3 we saw that global convergence is obtained for any  $\beta_k$  in the interval  $\mathcal{I}_1 = [-\beta_k^{\text{PR}}, \beta_k^{\text{FR}}]$ , and in §4 we proved global convergence for any  $\beta_k$  with Property (\*) contained in the interval  $\mathcal{I}_2 = [0, \infty)$ . We now ask whether these results can be combined to obtain larger intervals of admissible  $\beta_k$ . In particular, since the PR method has Property (\*), we ask whether global convergence is obtained by restricting  $\beta_k^{\text{PR}}$  to the larger interval  $\mathcal{I}_1 \cup \mathcal{I}_2$ , i.e., by letting

$$\beta_k = \begin{cases} \beta_k^{\text{PR}} & \text{if } \beta_k^{\text{PR}} \geq -\beta_k^{\text{FR}} \\ -\beta_k^{\text{FR}} & \text{otherwise.} \end{cases}$$

Interestingly enough, global convergence cannot be guaranteed, and this is shown in the fourth example of Powell (1984). In this example, the sequence  $\{\beta_k^{\text{PR}}/\beta_k^{\text{FR}}\}$  has exactly three accumulation points:

$$-1/3, 1 \text{ and } 10.$$

Therefore, there exists an index  $k_0$  such that  $\beta_k = \beta_k^{\text{PR}} \geq -\beta_k^{\text{FR}}$ , for all  $k \geq k_0$ . Now, the function can be modified and the starting point can be changed so that the PR method generates, from the new initial point  $\tilde{x}_1$ , a sequence  $\{\tilde{x}_k\}$  with  $\tilde{x}_k = x_{k+k_0-2}$ , for  $k \geq 2$ . In this modified example, we have  $\tilde{\beta}_k^{\text{PR}} \geq -\tilde{\beta}_k^{\text{FR}}$ , for all  $k \geq 2$ , but the sequence of gradients is bounded away from zero.

There is another example in which intervals of admissible  $\beta_k$  cannot be combined. Any method of the form (1.2)-(1.3) with a line search giving  $\langle g_k, d_{k-1} \rangle = 0$  for all  $k$ , and with  $\beta_k \in \mathcal{I}_3 = [-1, 1]$ , is globally convergent. This is easy to see, since in this case

$$\|d_k\|^2 \leq \|g_k\|^2 + \|d_{k-1}\|^2 \leq \dots \leq \bar{\gamma}^2 k,$$

where  $\bar{\gamma}$  is an upper bound on  $\|g(x)\|$ . Therefore  $\|d_k\|^2$  grows at most linearly, and global convergence follows by the arguments given in §2. On the other hand, Corollary 4.4 shows that the PR method is convergent if restricted to  $\mathcal{I}_2 = [0, \infty)$ . However the PR method may not converge if  $\beta_k^{\text{PR}}$  is restricted to  $\mathcal{I}_3 \cup \mathcal{I}_2 = [-1, \infty)$ . The argument is again based on the counter-example of Powell and on the fact that  $\beta_k^{\text{PR}} \geq -1/4$  for all  $k$  (this is proved by means of the Cauchy-Schwarz inequality; see Powell (1984)). Therefore in this example  $\beta_k^{\text{PR}} \in [-1, \infty)$ , but convergence is not obtained.

Therefore we are not able to generalize the results of §§3 and 4, and instead look more closely at the conditions used in these sections. We ask under what conditions is  $\beta_k^{\text{PR}} \geq 0$ , or  $\beta_k^{\text{PR}} \geq -\beta_k^{\text{FR}}$ . For strictly convex quadratic functions and exact line-searches, the PR method coincides with the FR method. Since  $\beta_k^{\text{FR}}$  is always positive, so is  $\beta_k^{\text{PR}}$ . Let us now consider strongly convex functions. It turns out that in this case  $\beta_k^{\text{PR}}$  can be negative, and in fact be less than  $-\beta_k^{\text{FR}}$ .

**Proposition 5.1** *There exists a  $C^\infty$  strongly convex function of two variables and a starting point  $x_1$  for which the PR method with exact line-searches gives  $\beta_3^{\text{PR}} < -\beta_3^{\text{FR}} < 0$ .*

**Proof.** Let us introduce the following strictly convex quadratic function  $\tilde{f}$  of two variables  $x = (x_{(1)}, x_{(2)})$ :

$$\tilde{f}(x) := x_{(1)}^2 + \frac{1}{2}x_{(2)}^2,$$

with gradient and Hessian (the Euclidean scalar product is assumed)

$$\nabla \tilde{f}(x) = \begin{pmatrix} 2x_{(1)} \\ x_{(2)} \end{pmatrix}, \quad \nabla^2 \tilde{f}(x) = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}.$$

Starting from the point  $x_1 = (-3, 3)$ , the PR method with exact line searches gives

$$\nabla \tilde{f}(x_1) = \begin{pmatrix} -6 \\ 3 \end{pmatrix}, \quad \bar{\alpha}_1 = \frac{5}{9} \quad \text{and} \quad \tilde{x}_2 = \frac{1}{3} \begin{pmatrix} 1 \\ 4 \end{pmatrix}.$$

Next, it finds

$$\nabla \tilde{f}(\tilde{x}_2) = \frac{2}{3} \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \quad \tilde{\beta}_2^{\text{PR}} = \frac{4}{81}, \quad \tilde{d}_2 = -\frac{10}{27} \begin{pmatrix} 1 \\ 4 \end{pmatrix} \quad \text{and} \quad \bar{\alpha}_2 = \frac{9}{10}.$$

The third point is the solution point  $x_* = (0, 0)$ .

We now perturb the function  $\tilde{f}$  inside the ball  $B(0, 1) := \{x : x_{(1)}^2 + x_{(2)}^2 < 1\}$ , defining

$$f(x) := \tilde{f}(x) + \epsilon\psi(x),$$

where the function  $\psi$  will be such that

$$\psi(x) = 0, \quad \forall x \notin B(0, 1), \quad (5.1)$$

and  $\epsilon$  will be a small positive number. As the line joining  $x_1$  and  $\tilde{x}_2$  does not intersect the closure of  $B(0, 1)$ , we see that the PR method on this new function  $f$ , starting from the same point  $x_1$ , will give  $x_2 = \tilde{x}_2$  and  $d_2 = \tilde{d}_2$ . We now show how to choose the function  $\psi$  and the number  $\epsilon > 0$  so that  $f$  is strongly convex and  $\beta_3^{\text{PR}}$  is negative.

We take for  $\psi$  a function of the form

$$\psi(x) := \eta(x)\ell(x),$$

where  $\ell$  is the linear function

$$\ell(x) := 4x_{(1)} - x_{(2)},$$

and  $\eta$  is a  $C^\infty$  function satisfying

$$\eta(x) = \begin{cases} 1 & \text{if } x \in B(0, \frac{1}{2}) \\ 0 & \text{if } x \notin B(0, 1). \end{cases}$$

Clearly,  $\psi$  satisfies (5.1), and has bounded second order derivatives. Therefore, by choosing  $\epsilon$  sufficiently small, say  $0 < \epsilon < \epsilon_1$ , the Hessian of  $f$  will be uniformly positive definite and  $f$  will be a  $C^\infty$  strongly convex function.

Now, when the function  $f$  is determined in this manner, there is a unique minimum of  $f$  from  $x_2$  in the direction  $d_2$ . As

$$\nabla f(0) = \nabla \tilde{f}(0) + \epsilon \nabla \psi(0) = \epsilon \begin{pmatrix} 4 \\ -1 \end{pmatrix}$$

is orthogonal to  $d_2 = \tilde{d}_2$ , the one-dimensional minimum is still obtained at  $x_3 = (0, 0)$  (but this is no longer the solution point). Therefore,

$$\beta_3^{\text{PR}} + \beta_3^{\text{FR}} = \frac{2|\nabla f(0)|^2 - \langle \nabla f(0), \nabla \tilde{f}(x_2) \rangle}{|\nabla \tilde{f}(x_2)|^2} = \frac{34\epsilon^2 - 4\epsilon/3}{20/9}.$$

We see that  $\beta_3^{\text{PR}} < -\beta_3^{\text{FR}} < 0$ , if  $0 < \epsilon < \epsilon_2 := 2/51$ .

By taking  $\epsilon \in (0, \min(\epsilon_1, \epsilon_2))$ , we obtain the desired result.  $\square$

This proposition shows that the convergence result given by Polak and Ribière (1969), which was obtained for strongly convex functions and exact line-searches, is not a consequence of Theorem 4.3, since the latter requires  $\beta_k \geq 0$ . Nor is it a consequence of Theorem 3.2, because Proposition 5.1 shows that  $\beta_k^{\text{PR}}$  can lie outside the interval  $[-\beta_k^{\text{FR}}, \beta_k^{\text{FR}}]$ .

| Problem | Name                           | Reference                     | $n$         |
|---------|--------------------------------|-------------------------------|-------------|
| 2       | Calculus of variations 2       | Gill and Murray (1973)        | 100, 200    |
| 3       | Calculus of variations 3       | Gill and Murray (1973)        | 100, 200    |
| 6       | Generalized Rosenbrock         | More et al. (1981)            | 100, 500    |
| 8       | Penalty 1                      | Gill and Murray (1979)        | 100, 1000   |
| 9       | Penalty 2                      | Gill and Murray (1979)        | 100         |
| 10      | Penalty 3                      | Gill and Murray (1979)        | 100, 1000   |
| 28      | Extended Powell singular       | More et al. (1981)            | 100, 1000   |
| 31      | Brown almost linear            | More et al. (1981)            | 100, 200    |
| 38      | Tridiagonal 1                  | Buckley and LeNir (1983)      | 100, 1000   |
| 39      | Linear minimal surface         | Toint (1983)                  | 121, 961    |
| 40      | Boundary-value problem         | Toint (1983)                  | 100         |
| 41      | Broyden tridiagonal nonlinear  | Toint (1983)                  | 100         |
| 42      | Extended ENGVL1                | Toint (1983)                  | 1000, 10000 |
| 43      | Extended Freudenstein and Roth | Toint (1983)                  | 100, 1000   |
| 45      | Wrong extended Wood            | Toint (1983)                  | 100         |
| 46(1)   | Matrix square root (ns=1)      | Liu and Nocedal (1988)        | 100         |
| 46(2)   | Matrix square root (ns=2)      | Liu and Nocedal (1988)        | 100         |
| 47      | Sparse matrix square root      | Liu and Nocedal (1988)        | 100, 1000   |
| 48      | Extended Rosenbrock            | More et al. (1981)            | 1000, 10000 |
| 49      | Extended Powell                | More et al. (1981)            | 100, 1000   |
| 50      | Tridiagonal 2                  | Toint (1983)                  | 100, 1000   |
| 51      | Trigonometric                  | More et al. (1981)            | 100, 1000   |
| 52      | Penalty 1 (2nd version)        | More et al. (1981)            | 1000, 10000 |
| 53      | INRIA ults0.4 (u0=0.95)        | Gilbert and Lemaréchal (1989) | 403         |
| 54      | INRIA ulcr1.2                  | Gilbert and Lemaréchal (1989) | 455         |
| 55      | INRIA ulcr1.3                  | Gilbert and Lemaréchal (1989) | 1559        |

Table 1: List of test functions.

## 6. Numerical Experiments

We have tested several of the algorithms suggested by the convergence analysis of this paper, on the collection of large test problems given in Table 1.

The starting points used are those given in the references. For the problems of Moré *et al* we set the parameter factor=1; for test problems 8, 9 and 10, starting point 3 from the reference was used. We verified that, in each run, all the methods converged to the same solution point; otherwise the problem was not included in the test set. The problems are not numbered consecutively because they belong to a larger test set. Since conjugate gradient methods are mainly useful for large problems, our test problems have at least 100 variables.

The following are the methods tested; they differ only in the choice of  $\beta_k$  and possibly, in the line search.

1. FR: The Fletcher-Reeves method.
2. PR-FR: The Polak-Ribière method constrained by the FR method, as in (3.7).
3. PR: The Polak-Ribière method.
4. PR<sup>+</sup>: The Polak-Ribière method allowing only positive values of  $\beta_k^{\text{PR}}$ , as in (4.24).

For the line search we used the algorithm of Moré and Thuente (1990). This algorithm finds a point satisfying the strong Wolfe conditions (2.16)-(2.17). We used the values  $\sigma_1 = 10^{-4}$  and  $\sigma_2 = 0.1$  which, by Theorem 3.2, ensure that methods FR and PR-FR are globally convergent. The line search for the PR and PR<sup>+</sup> methods was performed as follows. We first found a point satisfying the strong Wolfe conditions, using the values of  $\sigma_1$  and  $\sigma_2$  mentioned above. If at this point the directional derivative of  $f$  is negative, we know that the sufficient descent condition (4.1) holds, and terminate the line search (this was discussed at the beginning of §4.). On the other hand, if the directional derivative is positive, the algorithm of Moré and Thuente has bracketed a one-dimensional minimizer, and if the line search iteration is continued it will give, in the limit, a point  $x_k$  with  $\langle g_k, d_{k-1} \rangle = 0$ . By continuity and (4.2) it is clear that the line search will find a point satisfying the sufficient descent condition (4.1) in a finite number of iterations. In the numerical tests we set  $\sigma_3 = 10^{-2}$  in (4.1).

Our numerical experience with conjugate gradient methods indicates that it is advantageous to perform a reasonably accurate line search. Therefore, in addition to setting  $\sigma_2$  to the small number 0.1, we ensured that the line search evaluated the function at least twice. The choice of the initial trial value for the line search is also important. For the first iteration we set it to  $1/\|g_1\|$ , and for subsequent iterations we used the formula recommended by Shanno and Phua (1980), which is based on quadratic interpolation.

The tests were performed on a SPARCstation 1, using FORTRAN in double precision. All runs were stopped when

$$\|g(x_k)\|_\infty < 10^{-5}(1 + |f(x_k)|),$$

except for the INRIA problems for which the runs were stopped when the value of the function reached the thresholds given in the reference. The results in Tables 2 and 3 are given in the form: (number of iterations)/(number of function evaluations). The number



given under the column “mod” for method PR-FR denotes the number of iterations for which  $|\beta_k^{\text{PR}}| > \beta_k^{\text{FR}}$ . For method  $\text{PR}^+$ , “mod” denotes the number of iterations for which  $\beta_k^{\text{PR}} < 0$ . If the limit of 9999 function evaluations was exceeded the run was stopped; this is indicated by “\*”. The sign “\*\*” means that the run stopped because the line search procedure described above failed to find a steplength. This generally occurs when the stopping criterion is very demanding.

| P     | N    | FR        | PR-FR     |      | PR        | PR <sup>+</sup> |     |
|-------|------|-----------|-----------|------|-----------|-----------------|-----|
|       |      | it/f-g    | it/f-g    | mod  | it/f-g    | it/f-g          | mod |
| 2     | 100  | 405/827   | 405/820   | 351  | 400/812   | 400/812         | 0   |
| 3     | 100  | 1313/2627 | 1313/2627 | 1313 | 1299/2599 | 1299/2599       | 0   |
| 6     | 100  | *         | 261/547   | 95   | 256/529   | 254/525         | 1   |
| 8     | 100  | 10/36     | 15/49     | 12   | 9/39      | 12/47           | 2   |
| 9     | 100  | 7/20      | 8/22      | 6    | 8/25      | 7/20            | 2   |
| 10    | 100  | 116/236   | 93/191    | 91   | 118/244   | 119/244         | 1   |
| 28    | 100  | 1426/2855 | 1291/2584 | 1289 | 120/280   | 168/382         | 3   |
| 31    | 100  | 2/3       | 2/3       | 1    | 1/4       | 1/4             | 0   |
| 38    | 100  | 70/142    | 70/142    | 47   | 71/144    | 71/144          | 0   |
| 39    | 121  | *         | 59/122    | 4    | 59/122    | 59/122          | 0   |
| 40    | 100  | 175/351   | 175/351   | 175  | 132/266   | 132/266         | 0   |
| 41    | 100  | 29/60     | 24/50     | 1    | 24/50     | 24/50           | 0   |
| 42    | 1000 | 10/27     | 9/25      | 8    | 10/34     | 9/30            | 2   |
| 43    | 100  | 16/41     | 14/39     | 13   | 16/44     | 13/37           | 1   |
| 45    | 100  | *         | 74/166    | 66   | 37/90     | 45/109          | 3   |
| 46(1) | 100  | 617/1238  | 253/510   | 248  | 257/518   | 257/518         | 0   |
| 46(2) | 100  | 886/1776  | 251/506   | 243  | 251/506   | 251/506         | 0   |
| 47    | 100  | 151/306   | 59/122    | 50   | 60/124    | 60/124          | 0   |
| 48    | 1000 | 79/185    | 71/172    | 66   | 26/73     | 23/70           | 3   |
| 49    | 100  | 1426/2855 | 1291/2584 | 1289 | 117/281   | 168/382         | 3   |
| 50    | 100  | 72/146    | 72/146    | 52   | 72/146    | 72/146          | 0   |
| 51    | 100  | 202/409   | 42/94     | 12   | 45/103    | 45/103          | 0   |
| 52    | 1000 | 3/10      | 3/10      | 2    | 4/12      | 4/12            | 2   |

Table 2: Smaller problems.

It is interesting to note that  $\beta_k^{\text{PR}}$  was constrained in most of the iterations of the method PR-FR, but was only rarely modified in the  $\text{PR}^+$  method. Many of the problems were run again for a larger number of variables. The results are given in Table 3.

The performance of methods PR-FR, PR and  $\text{PR}^+$  is comparable. Overall,  $\text{PR}^+$  appears to be better than PR. The FR method is clearly the least efficient, requiring an exceedingly large number of function evaluations in some problems.

In these runs the methods were implemented without restarting. We also performed tests in which the methods were restarted along the steepest descent direction every  $n$  iterations. (Since  $n$  is large, very few restarts were performed.) The FR method improved

| P  | N     | FR        | FR-PR     |      | PR        | PR <sup>+</sup> |     |
|----|-------|-----------|-----------|------|-----------|-----------------|-----|
|    |       | it/f-g    | it/f-g    | mod  | it/f-g    | it/f-g          | mod |
| 2  | 200   | 703/1424  | 701/1420  | 596  | 701/1420  | 701/1420        | 0   |
| 3  | 200   | 2808/5617 | 2808/5617 | 2808 | 2631/5263 | 2631/5263       | 0   |
| 6  | 500   | *         | 1107/2231 | 433  | 1068/2151 | 1067/2149       | 1   |
| 8  | 1000  | 12/39     | 9/34      | 7    | 6/28      | 10/42           | 2   |
| 10 | 1000  | 138/281   | 145/299   | 142  | 165/338   | 165/338         | 0   |
| 28 | 1000  | 533/1102  | 1369/2741 | 1366 | 212/473   | 97/229          | 3   |
| 31 | 200   | 2/4       | 2/4       | 1    | 1/5       | 1/5             | 0   |
| 38 | 1000  | 264/531   | 263/529   | 217  | 262/527   | 262/527         | 0   |
| 39 | 961   | *         | 143/290   | 5    | 142/287   | 142/287         | 0   |
| 42 | 10000 | 6/26      | 6/26      | 5    | 7/28      | 6/26            | 1   |
| 43 | 1000  | 10/27     | 15/38     | 15   | 10/33     | 9/29            | 2   |
| 47 | 1000  | 422/849   | 114/233   | 92   | 113/231   | 113/231         | 0   |
| 48 | 10000 | 61/143    | 130/283   | 123  | 24/73     | 19/62           | 4   |
| 49 | 1000  | 568/1175  | 1369/2741 | 1366 | 212/473   | 97/229          | 3   |
| 50 | 1000  | 274/551   | 273/549   | 245  | 274/551   | 274/551         | 0   |
| 51 | 1000  | 231/467   | 40/91     | 5    | 40/92     | 40/92           | 0   |
| 52 | 10000 | 4/15      | 4/15      | 4    | 3/13      | 3/13            | 1   |
| 53 | 403   | **        | 233/494   | 130  | 237/508   | 237/508         | 0   |
| 54 | 455   | **        | 44/91     | 7    | 44/87     | 44/87           | 0   |
| 55 | 1559  | 23/47     | 23/49     | 15   | 23/47     | 23/47           | 0   |

Table 3: Larger problems.

substantially, but this method was still the least efficient of the four. The other 3 methods performed similarly with and without restarts, and we will not present the results here.

We now give an example that illustrates the inefficient behavior of the FR method, as predicted in §3. For problem 45 with  $n = 100$ , we observed that for hundreds of iterations  $\cos \theta_k$  stays fairly constant, and is of order  $10^{-2}$ , while the steps  $\|x_k - x_{k-1}\|$  are of order  $10^{-2}$  to  $10^{-3}$ . This causes the algorithm to require a very large number of iterations to approach the solution. A restart along the steepest descent direction terminates this cycle of bad search directions and tiny steps. A similar behavior was observed in several other problems.

## REFERENCES

- Al-Baali, M. (1985). Descent property and global convergence of the Fletcher-Reeves method with inexact line search, *IMA Journal of Numerical Analysis*, Vol. 5, pp. 121-124.
- Baptist, P. and Stoer, J. (1977). On the relation between quadratic termination and convergence properties of minimization algorithms, Part II, Applications, *Numerische Mathematik*, Vol. 28, pp. 367-392.
- Buckley, A. and LeNir, A. (1983). QN-like variable storage conjugate gradients, *Mathematical Programming*, Vol. 27, pp.155-175.
- Cohen, A. (1972). Rate of convergence of several conjugate gradient algorithms, *SIAM Journal on Numerical Analysis*, Vol. 9, pp. 248-259.
- Crowder, H.P. and Wolfe, P. (1972). Linear convergence of the conjugate gradient method, *IBM Journal of Research and Development*, Vol. 16, pp. 431-433.
- Fletcher, R. and Reeves, C.M. (1964). Function minimization by conjugate gradients, *Computer Journal*, Vol. 7, pp. 149-154.
- Fletcher, R. (1987). *Practical Methods of Optimization*, Wiley.
- Gilbert, J.Ch. and Lemaréchal, C. (1989). Some numerical experiments with variable-storage quasi-Newton algorithms, *Mathematical Programming B*, Vol. 45, pp. 407-436.
- Gill, P.E. and Murray, W. (1973). The numerical solution of a problem in the calculus of variations, in D.J. Bell, ed., *Recent Mathematical Developments in Control*, Academic Press, New York, pp. 97-122.
- Gill, P.E. and Murray, W. (1979). Conjugate-gradient methods for large-scale nonlinear optimization, Technical report SOL 79-15, Department of Operations Research, Stanford University, Stanford, CA.
- Gill, P.E., Murray, W. and Wright, M.H. (1981). *Practical Optimization*, Academic Press.
- Lemaréchal, C. (1981). A view of line searches, in Auslander, Oettli and Stoer, eds., *Optimization and Optimal Control*, Lecture Notes in Control and Information Science 30, pp. 59-78, Springer Verlag.
- Liu, D.C. and Nocedal, J. (1988). Test results of two limited memory methods for large scale optimization, Report NAM 04, Department of Electrical Engineering and Computer Science, Northwestern University.

- Moré, J.J., Garbow, B.S. and Hillstom, K.E. (1981). Testing unconstrained optimization software, *ACM Transactions on Mathematical Software*, Vol. 7, pp. 17-41.
- Moré, J.J. and Thuente, D.J. (1990). On line search algorithms with guaranteed sufficient decrease, Mathematics and Computer Science Division Preprint MCS-P153-0590, Argonne National Laboratory, Argonne II 60439.
- Polak, E. and Ribière, G. (1969). Note sur la convergence de méthodes de directions conjuguées, *Revue Française d'Informatique et de Recherche Opérationnelle*, Vol. 16, pp. 35-43.
- Powell, M.J.D. (1976). Some convergence properties of the conjugate gradient method, *Mathematical Programming*, Vol. 11, pp. 42-49.
- Powell, M.J.D. (1977). Restart procedures of the conjugate gradient method, *Mathematical Programming*, Vol. 12, pp. 241-254.
- Powell, M.J.D. (1984). Nonconvex minimization calculations and the conjugate gradient method, in *Lecture Notes in Mathematics*, Vol. 1066, pp. 122-141, Springer-Verlag, Berlin.
- Powell, M.J.D. (1985). Convergence properties of algorithms for nonlinear optimization, Report DAMTP 1985/NA1, Department of Applied Mathematics and Theoretical Physics, University of Cambridge, England. Presented at the 1984 SIAM Summer Meeting in Seattle.
- Shanno, D.F. and Phua, K.H. (1980). Remark on algorithm 500: minimization of unconstrained multivariate functions, *ACM Transactions on Mathematical Software*, Vol. 6, pp. 618-622.
- Stoer, J. (1977). On the relation between quadratic termination and convergence properties of minimization algorithms, Part I, Theory, *Numerische Mathematik*, Vol. 28, pp. 343-366.
- Toint, Ph.L. (1983). Test problems for partially separable optimization and results for the routine PSPMIN, Report 83/4, Department of Mathematics, Facultés Universitaires de Namur, Namur, Belgium.
- Touati-Ahmed, D. and Storey, C. (1990). Efficient hybrid conjugate gradient techniques, *Journal of Optimization Theory and Applications*, Vol. 64, pp. 379-397.
- Wolfe, P. (1969). Convergence conditions for ascent methods, *SIAM Review*, Vol. 11, pp. 226-235.
- Wolfe, P. (1971). Convergence conditions for ascent methods II: some corrections, *SIAM Review*, Vol. 13, pp. 185-188.
- Zoutendijk, G. (1970). Nonlinear Programming, computational methods, in *Integer and Nonlinear Programming*, pp. 37-86, ed. J. Abadie, North-Holland, Amsterdam.

**ISSN 0249 - 6399**