



# Optimality of the parameterization of quadrics and their intersections

Daniel Lazard

## ► To cite this version:

| Daniel Lazard. Optimality of the parameterization of quadrics and their intersections. Journées de clôture Visi3D et CoSTIC, 2001, Paris, France. inria-00099402

HAL Id: inria-00099402

<https://inria.hal.science/inria-00099402>

Submitted on 26 Sep 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

December 11, 2001

and Laurent DuPont

Sylvain Lazard, Sylvain Petitjean

In collaboration with

Daniel Lazard

Optimality of the Parameterization of  
Quadratics and their Intersections

the number of radicals involved  
means here optimality in term of  
Optimality

## Quadratics of rank 1 or 2

Rank 1 :

$$x^2$$

Gauss-Jordan reduction leads to equations :

Rational parameterization = no radicals

Clearly optimal

Rank 2 :

$$x^2 - ay^2$$

Optimal : There are regular rational points iff  $\sqrt{a}$  is rational.

Rational parameterization in  $\mathbb{Q}(\sqrt{a})$

$x^2 + y^2 - 3z^2$  has no rational point (Proof over  $\mathbb{Z}/4\mathbb{Z}$ )

Example where a square root is needed :

point

Gauss-Jordan gives equation  $x^2 + y^2 - z^2$  when starting from this  $\Leftrightarrow$

The conic  $ax^2 + by^2 - cz^2$  has a rational point over  $\mathbb{Q}$   $\Leftrightarrow$

There exists a parametrization rational over  $\mathbb{Q}$

$\Leftarrow$  At most one square root

$$x = 2uv, y = \frac{u^2 - aqv^2}{u^2 + av^2}, z = \frac{\sqrt{bc}}{u^2 + av^2}, t = w$$

parametrization :

$$ax^2 + by^2 - cz^2$$

Gauss-Jordan reduction leads to equation :

Quadratics of rank 3

Another square root is needed iff the quadratic has no rational point.

$\sqrt{d}$  always needed

We claim :

Invariant up to a square by change of coordinates

$Q := abcd =$  discriminant of the quadratic.

2 square roots

$$x = \frac{pq\sqrt{v}}{uv+qv}, y = \frac{q\sqrt{av}}{uv-qv}, z = \frac{a\sqrt{v}}{uv-qv}, t = \frac{q}{uv-qv}$$

Parameterization :

$$ax^2 + qy^2 - cz^2 - dt^2$$

Gauss-Jordan reduction leads to equation :

Quadratics of signature (2,2)

rational frame.

- $\Leftrightarrow$  It has a rational point over  $\mathbb{Q}$  and equation  $x^2 + y^2 - z^2 - \mathfrak{d}t^2$  in some  $\mathbb{Q}\sqrt{\mathfrak{d}}$

$\Leftrightarrow$  The field of a rational parameterization contains  $\mathbb{Q}(\sqrt{\mathfrak{d}})$

- $\Leftrightarrow$   $\mathbb{Q}$  has equation  $x^2 + y^2 - z^2 - t^2$ , in some rational frame over  $\mathbb{K}$
- $\Leftrightarrow$   $\mathbb{Q}$  has a rational point over  $\mathbb{K}$  and  $\mathfrak{d}$  is a square in  $\mathbb{K}$
- $\Leftrightarrow$   $\mathbb{Q}$  contains a line which is rational over  $\mathbb{K}$
- $\Leftrightarrow$   $\mathbb{Q}$  admits a rational parameterization over  $\mathbb{K}$ , linear in one parameter

## Quadratics of signature (2,2) (continued)

Proof over  $\mathbb{Z}/4\mathbb{Z}$  and  $\mathbb{Z}/8\mathbb{Z}$

and no rational parameterization over  $\mathbb{Q}(\sqrt{3})$   
has no rational point over  $\mathbb{Q}$

$$x^2 + y^2 - 3z^2 - 11t^2$$

One depending on the parameter,  $\sqrt{\Delta}$

A constant one,  $\sqrt{\delta}$ , the discriminant of the quadratic

**Parameterization through this quadratic involves 2  $\sqrt{\_}$ :**

the pencil contains a quadratic of signature  $(2, 2)$  with a rational point  
(i.e.  $\det(\lambda P + u Q)$  has no multiple (projective) root),

If the intersection is not empty nor singular

of the intersection (**linear equation to solve**)

There is exactly one quadratic of the pencil passing through a point outside

The intersection is contained in any quadratic of the pencil  $\lambda P + u Q$

$P$  and  $Q$  quadratics

Intersections of quadratics

If the intersection is non singular,  $\Delta$  is needed

Putting the parameterization of the quadric in  $\mathbb{P}$  gives an equation between the parameters.

Iso-morphism of the intersection

with a curve in  $\mathbb{P}_1 \times \mathbb{P}_1$ , the spaces of parameters

De-homogenizing and re-homogenizing :

bi-rational equivalence with a curve in  $\mathbb{P}^2$

of degree 4 with 2 ordinary singular points at infinity.

Non singular intersection has genus 1  $\iff$

No rational parameterization over  $\mathbb{C}$   $\iff$

$$\varphi = 3x^2 + y^2 - z^2 - t^2$$

$$\psi = 5y^2 + 6xy + 2z^2 - t^2 + 6zt;$$

This surface has no rational point in the case :

has a rational point.

$$z^2 = \det(\varphi(x,y,1,0) \psi - \psi(x,y,1,0) \varphi)$$

$\Leftrightarrow$  The surface

defined by such a pair has a square discriminant

the quadratic passing through a point on a rational line

the intersection contains pair of conjugate points of degree 2

If  $\sqrt{\delta}$  is not needed,

a surface of degree 8 has a rational point

if

$\sqrt{\delta}$  is not needed

there are **28 singular intersections** corresponding to **46 pencils**.

Up to a **real projective transformation**,

**singular point of  $\Phi$**

- $\Leftrightarrow$  The intersection has a **singular (complex) point** which is also a  **$\Phi$**  is a cone and all quadratics of the pencil are tangent at its vertex
- $\Leftrightarrow$   **$\Phi$**  has rank  $\leq 2$  (pair of planes) or
- $0$  is a multiple root of  $\det(\Phi + \lambda Q)$

## Singular intersections

Algorithm sketched by Sylvain Lazard  
 managed through the number of multiple roots of  $\det(\lambda P + u Q)$

If the intersection does not consist in 4 lines  
 at most 2  $\swarrow$  for parameterizing all components  
 at most 1  $\swarrow$  may be unnecessary  
 it may be removed by finding a rational point on a conic  
 If the intersection consists in 4 non coplanar lines (skew quadrilateral)  
 at most 2  $\swarrow$  for each line  
 possibility of 3  $\swarrow$  for all lines together  
 always optimal for each line and for all lines together.  
 In the case of 4 concurrent lines  
 general degree 4 equation



Totally rational parameterization

and a factor linear in that variable (the cubic)

a univariate factor (the line)

$\iff$  Equation in the parameters factors (GCD computation) in

Parameterize through a (2,2) quartic : no  $\checkmark$

$\det(\lambda P + \mu Q)$  has 2 double roots, real or not

## Cubic and Line

One  $\checkmark$ , zero if one finds a rational point on the cone

$\iff$  equation which is linear in one of the parameter.

Plug it in the equation of  $\mathcal{C}$

Parameterize the cone corresponding to this root : 1 or 0  $\checkmark$

$\det(\lambda P + \mu Q)$  has a single multiple root

## Singular quartic

field of the planes

Second  $\wedge$  may be avoided for each conic with a rational point on the

First  $\wedge$  is needed if the planes are not rational

$\iff$  Linear equations in the parameters

equations of the planes : 1 or 0  $\wedge$

Parameterize through a  $(2,2)$  quartic and plug the parameters in the

Compute these planes : 1 or 0  $\wedge$

The corresponding quadratic is a pair of planes

$\det(\lambda P + u Q)$  has 1 (rational) double root

## Two conics

It may be avoided for the conic iff it has a rational point

At most 1 ✓

and a bilinear one (the conic)

2 univariate factors (the lines)

⇒ Equation in the parameters factors (GCD computation) in

Parameterize through a (2,2) quadratic : one ✓ iff the lines are irrational

The corresponding quadratics are a cone and a pair of planes

$\det(\lambda p + \mu q)$  has 2 (rational) double roots

## Conic and 2 Lines

4 Lines

$\det(\lambda P + \mu Q)$  has 2 double roots

The corresponding quadratics are pairs of planes

If the roots are rational

Compute the planes and their intersection : 0, 1 or 2  $\sqrt{d}$ , unavoidable

Else

Parameterize through a (2,2) quadratic : one  $\sqrt{d}$

$\Leftrightarrow$  Equation in the parameters factors (GCD computation)  
in 2 univariate factors of degree 2.

Each linear factor parameterizes one line with at most 2  $\sqrt{d}$

(all together 3  $\sqrt{d}$ )

If the roots of  $\det(\lambda P + \mu Q)$  generates the same field as  $\sqrt{d}$

2  $\sqrt{d}$  all together

Always optimal

Plug the parametrization of the quadratic in each plan of a pair :