

Clinical application of iCMR for physiological measurement

Andrew Taylor

▶ To cite this version:

Andrew Taylor. Clinical application of iCMR for physiological measurement. CI2BM09 - MICCAI Workshop on Cardiovascular Interventional Imaging and Biophysical Modelling, Sep 2009, London, United Kingdom. inria-00418434

HAL Id: inria-00418434 https://inria.hal.science/inria-00418434

Submitted on 18 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Clinical application of iCMR for physiological measurement

Andrew Taylor

University College London Institute of Child Health & Great Ormond Street Hospital, London, UK

This presentation will focus on the potential physiological applications of hybrid MR/X-ray interventional technology (Figure 1). In the first half of the presentation, three clinical areas with be described – 1) Assessment of pulmonary hypertension; 2) Assessment of cardiac function, great vessel flow and pressures during treatment with devices; and 3) Assessment of physiology prior to treatment in paediatric congenital heart disease.

<u>Figure 1</u> – 360° 'fish-eye' view of the combined bi-plane catheter lab and MR scanner. The MR scanner (right) is viewed through the open sliding doors



The second half of the presentation will describe how the complex MR and pressure data, obtained in the MR/X-ray hybrid lab, can be used to validate models built from more simply MR (and or CT) data sets (Figure 2). Such models may ultimately allow complex parameters such as pressure/volume loops to be defined, without the need for invasive pressure measurements.

<u>Figure 2</u> – Combined MR and CT fusion image showing the heart (pale blue), pulmonary valve stent (orange), coronary arteries (red) and aortic valve prosthesis (pink).

