



**HAL**  
open science

## Fast, model-free, analytical diffusion PDF profile estimation from the DWI signals

Jian Cheng, Aurobrata Ghosh, Tianzi Jiang, Rachid Deriche

### ► To cite this version:

Jian Cheng, Aurobrata Ghosh, Tianzi Jiang, Rachid Deriche. Fast, model-free, analytical diffusion PDF profile estimation from the DWI signals. the Sixteenth Annual Meeting of the Organization for Human Brain Mapping (OHBM), 2010, Barcelona, Spain. inria-00497244

**HAL Id: inria-00497244**

**<https://inria.hal.science/inria-00497244>**

Submitted on 2 Jul 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Fast, model-free, analytical diffusion PDF profile estimation from the DWI signals

### Poster No:

837 WTh-AM

### Authors:

Jian Cheng<sup>1</sup>, Aurobrata Ghosh<sup>1</sup>, Tianzi Jiang<sup>2</sup>, Rachid Deriche<sup>1</sup>

### Institutions:

<sup>1</sup>INRIA Sophia Antipolis, Sophia Antipolis, France, <sup>2</sup>Institute Of Automation, Chinese Academy Of Sciences, Beijing, China

### Introduction:

How to estimate the diffusion Probability Density Function (PDF) from the q-space DWI signals is an open problem in diffusion MRI field. The diffusion PDF and the diffusion attenuation of dMRI signal is related by 3 dimensional Fourier transform. Diffusion Orientation Transform (DOT) [2] was proposed based on the assumption of the mono-exponential decay. Thanks to this assumption, PDF profile at a given radius  $R_0$  could be calculated analytically. However the estimated PDF is a low-pass filtered PDF. And this kind of low-pass filtering comes from the intrinsic modeling error [2], which is not a well defined filtering. Although DOT has been extended to multi-exponential model [2] and can reduce the modeling error dramatically, a nonlinear fitting needs to be performed for every voxel, which will limit its usefulness. Here we propose a model-free, linear, robust, analytical PDF profile reconstruction method which avoids the exponential or multi-exponential assumption.

### Methods:

The diffusion Probability Density Function and the diffusional attenuation of dMRI signal is related by 3D Fourier transform. See formula (1). where  $E(q) = S(q)/S_0$  is the normalized DWI signal measured on  $q$ , and  $S_0$  is the baseline image acquired without diffusion gradients. Spherical Polar Fourier expression is a kind of orthonormal basis representation and it has been shown in [1] that it can represent the diffusion weighted image signal sparsely. See formula (2).  $R_n(|q|)$  is the Gaussian-Laguerre polynomial in formula (3), where  $\zeta$  is the scale parameter and  $Y_l^m$  is the  $l$  order  $m$  degree spherical harmonics (SH). After we estimate the coefficients of the signal under the basis from a least square fit or nonlinear robust estimation [1], we proved that there is a linear, analytical solution to get the coefficients of the PDF profile at a given radius  $R_0$  under spherical harmonics representation. To get this result, firstly consider the plane wave equation in spherical coordinates in formula (4). Then put formula (2) and (4) into (1), we have the expression of the PDF profile in formula (5), where  ${}_1F_1$  is the confluent hypergeometric function. Please note that  $f_{n,l}$  is independent with data and the summation over  $n$  actually is just a linear transformation which could be implemented as an matrix multiplication. Since the coefficients  $a_{n,l,m}$  also could be estimated through a linear matrix multiplication [1]. Thus calculating the coefficients for PDF profile from the DWI signal here is very fast. The main computation complex is in the estimation of  $a_{n,l,m}$ . But it is still very fast if least square fitting is used. Our suggestion here is to store  $a_{n,l,m}$  once it is estimated. Then estimating the coefficients for PDF profile at several given  $R_0$  could be performed on the stored  $a_{n,l,m}$ . That means we just need to calculate  $a_{n,l,m}$  once and could re-calculate different PDF profile in different radii very fast.

### Results:

We apply our method to a real human data. This data has 110 DWIs, 50 diffusion encoding gradients with a b-value of  $1126 \text{ s/mm}^2$ , 30 gradients with b-value of  $820 \text{ s/mm}^2$ , 10 gradients respectively with b-value of 100, 307,  $512 \text{ s/mm}^2$ , twice-refocused spin-echo EPI sequence,  $TE = 100 \text{ ms}$ ,  $1.5 \times 1.5 \times 1.5 \text{ mm}$  voxel resolution, three repetitions, corrected for subject motion. It should be noted that this sampling scheme is not appropriate for a good estimation of the coefficients of the signal, since the b-values here are too small. For a good representation of the signal, we should keep the b-values range from small values to large values. A discussion about evaluating the reconstruction error in different sampling schemes could be found in [1]. However, even though this is not a good sampling, the result is still impressive. We visualize the PDF profile in  $R_0 = 17 \text{ ms}$  for one slice. Two areas were enlarged for good visualization. The glyphs were colored by GFA calculated from the estimated PDF profiles. Some crossings were found. Please note that we did not do any normalization here, e.g. min-max normalization [3]. That is because of two reasons. 1) the PDF profiles in white matter seem sharper enough and the profiles in CSF and gray matter are almost isotropic. 2) we will lost the radial information if we do some normalization. Please note that the radial information in PDF is also important compared with its peaks. It tell us how much the probability  $P(R_0)$  is. From the result, we can see that the CSF has the largest probability (glyph size) compared with white matter and gray matter, just like the visualization of tensors in DTI. Tensors can not tell us crossing fibers, while PDF profiles can.

### Conclusions:

We proposed a fast analytical PDF profile reconstruction method based on spherical polar Fourier expression of the signal in Q-space. It can avoid the modeling error from mono-exponential model or multi-exponential model. The coefficients of the PDF profile under spherical harmonics expression could be linearly and analytically calculated from the coefficients of the signal. It is a linear transformation that is independent with data. This transformation matrix is just needed to calculated only once for a whole data set, which makes the method very fast. The experiments validate our method which could provide both the radial information and the direction information and could detect the fiber crossing even for the data with small b-values.

$$P(\mathbf{R}) = \int E(\mathbf{q}) e^{-2\pi i \mathbf{q} \cdot \mathbf{R}} d\mathbf{q} \quad \left[ q = |\mathbf{q}|, R_0 = |\mathbf{R}|, \mathbf{q} = q\mathbf{u}, \mathbf{R} = R_0\mathbf{r} \right] \quad (1)$$


$$E(\mathbf{q}) = \sum_{n=0}^N \sum_{l=0}^L \sum_{m=-l}^l a_{n,l,m} R_n(\|\mathbf{q}\|) Y_l^m(\mathbf{u}) \quad (2)$$


$$R_n(\|\mathbf{q}\|) = \left[ \frac{2}{\zeta^{3/2}} \frac{n!}{\Gamma(n+3/2)} \right]^{1/2} \exp\left(-\frac{\|\mathbf{q}\|^2}{2\zeta}\right) L_n^{1/2}\left(\frac{\|\mathbf{q}\|^2}{\zeta}\right) \quad (3)$$


$$e^{\pm 2\pi i \mathbf{q} \cdot \mathbf{R}} = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^l (\pm i)^l j_l(2\pi q R_0) Y_l^m(\mathbf{u}) Y_l^m(\mathbf{r}) \quad (4)$$

$$\begin{aligned} P(\mathbf{R}) &= 4\pi \sum_{l=0}^L \sum_{m=-l}^l (-1)^{l/2} \sum_{n=0}^N \left\{ a_{n,l,m} \int_0^{\infty} j_l(2\pi q R_0) R_n(\|\mathbf{q}\|) q^2 dq \right\} Y_l^m(\mathbf{r}) \\ &= \sum_{l=0}^L \sum_{m=-l}^l \left\{ 4\pi (-1)^{l/2} \zeta^{l/2+3/2} R_0^l \sum_{n=0}^N a_{n,l,m} f_{n,l} \right\} Y_l^m(\mathbf{r}) \end{aligned} \quad (5)$$

$$f_{n,l} = \left[ \frac{2}{\zeta^{3/2}} \frac{n!}{\Gamma(n+3/2)} \right]^{1/2} \sum_{i=0}^n \frac{(-1)^i \binom{n+0.5}{n-i} 2^{l/2+i-1/2} \pi^{l+1/2} \Gamma(l/2+i+3/2)}{i! \Gamma(l+3/2)} {}_1F_1(i+l/2+3/2, l+3/2, -2\pi^2 R_0^2 \zeta)$$





**References:**

- Assemla, H. (2009), 'Efficient and robust computation of pdf features from diusion mr signal', *Medical Image Analysis*, vol. 13, pp. 715-729.
- Ozarslan, E. (2006), 'Resolution of complex tissue microarchitecture using the diusion orientation transform (dot)', *NeuroImage*, vol. 31, pp. 1086-1103.
- Tuch, D. (2004), 'Q-ball Imaging', *Magnetic Resonance in Medicine*, vol. 52, pp. 1358-1372.

**Categories**

- Diffusion MRI (Imaging Techniques and Contrast Mechanism)