



# Hydrodynamic limit for a type of exclusion processes with slow bonds in dimension greater than 2

Tertuliano Franco

## ► To cite this version:

Tertuliano Franco. Hydrodynamic limit for a type of exclusion processes with slow bonds in dimension greater than 2. Journées MAS et Journée en l'honneur de Jacques Neveu, Aug 2010, Talence, France. inria-00514283

HAL Id: inria-00514283

<https://inria.hal.science/inria-00514283>

Submitted on 2 Sep 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journées MAS 2010, Bordeaux

Session : Systèmes à une infinité de particules en interaction et applications

## **Hydrodynamic limit for a type of exclusion processes with slow bonds in dimension greater than 2**

par **Tertuliano Franco**

Let  $\Lambda$  be a connected closed region with smooth boundary contained in the  $d$ -dimensional continuous torus  $\mathbf{T}^d$ . In the discrete torus  $N^{-1}\mathbf{T}_N^d$ , we consider a nearest neighbor symmetric exclusion process where occupancies of neighboring sites are exchanged at rates depending on  $\Lambda$  in the following way : if both sites are in  $\Lambda$  or in its complement  $\Lambda^C$ , the exchange rate is one ; If one site is in  $\Lambda$  and the other one is in  $\Lambda^C$  and the direction of the bond connecting the sites is  $e_j$ , then the exchange rate is defined as  $N^{-1}$  times the absolute value of the inner product between  $e_j$  and the normal exterior vector to  $\partial\Lambda$ . We show that this exclusion type process has a non-trivial hydrodynamical behavior under diffusive scaling and, in the continuum limit, particles are not blocked or reflected by  $\partial\Lambda$ . Thus the model represents a system of particles under hard core interaction in the presence of a permeable membrane which slows down the passage of particles between two complementar regions.

Joint work with Adriana Neumann and Glauco Valle.

*Adresse :*

Tertuliano FRANCO  
IMPA  
Estrada Dona Castorina 110, Jardim Botanico.  
Rio de Janeiro, 22460-320, Brazil  
E-mail : [tertu@impa.br](mailto:tertu@impa.br)

Session : Systèmes à une infinité de particules en interaction et applications