



**HAL**  
open science

## Two results concerning chirps and 2-microlocal exponents prescription

Bertrand Guilheneuf, Stéphane Jaffard, Jacques Lévy Véhel

► **To cite this version:**

Bertrand Guilheneuf, Stéphane Jaffard, Jacques Lévy Véhel. Two results concerning chirps and 2-microlocal exponents prescription. *Applied and Computational Harmonic Analysis*, 1998, 5 (4), pp.487-492. 10.1006/acha.1997.0241 . inria-00593247

**HAL Id: inria-00593247**

**<https://inria.hal.science/inria-00593247v1>**

Submitted on 13 May 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Two results concerning Chirps and 2-microlocal exponents prescription

Bertrand Guiheneuf\*, Stéphane Jaffard†, Jacques Lévy Véhel\*

\* Projet FRACTALES, INRIA Rocquencourt,  
78153 Le Chesnay - FRANCE,  
Bertrand.Guieheneuf@inria.fr, Jacques.Levy\_Vehel@inria.fr

† Département de Mathématiques, Université Paris XII,  
61 Av. du Gal. de Gaulle, 94010 Créteil Cedex FRANCE, and C.M.L.A.,  
E.N.S. de Cachan, Jaffard@cmla.ens-cachan.fr

**Abstract:** The local behavior of a function near a Hölder singularity located at a point  $x_0$  can be finely described by means of its 2-microlocal domain. We solve the issue of prescribing this domain. A more compact description can be given using two parameters, the Hölder exponent and the chirp exponent. We give a necessary condition satisfied by this couple of exponents considered as functions of  $x_0$ .

# 1 Introduction

Let  $\alpha$  be a positive real number and  $x_0 \in \mathbf{R}$ ; a function  $f : \mathbf{R} \rightarrow \mathbf{R}$  is  $C^\alpha(x_0)$  if there exists a polynomial  $P$  of degree less than  $\alpha$  such that

$$|f(x) - P(x - x_0)| \leq C|x - x_0|^\alpha. \quad (1)$$

The *Hölder exponent*  $h_f(x_0)$  is the supremum of all the values of  $\alpha$  such that (1) holds.  $h(\cdot)$  thus defines a function, the pointwise values of which describe the local variations of the irregularity of the function  $f$ . The Hölder exponent, as a measure of pointwise regularity of functions, is a powerful tool in many applications such as image analysis or signal analysis [9] [4]. The issue of prescribing  $h_f(x)$  has been solved in [6] and [3]: The class of all admissible functions  $h_f(x)$  (when  $f \in C^0(\mathbf{R})$ ) coincides with the class of lower limits of continuous functions.

However, characterizing the regularity with the sole Hölder exponent has several drawbacks. It does not give any information about the oscillatory behavior of the function near the Hölder singularities; furthermore  $h_f(x)$  is not stable under the action of (pseudo) differential operators. This precludes, for instance, the use of the Hilbert transform, commonly used for estimation on sampled signals. The 2-microlocal spaces  $C_{x_0}^{s,s'}$  [5, 8] have been introduced by Bony [2] to cope with such issues in the frame of p.d.e. analysis. The rough idea is to incorporate knowledge about the regularity “in a neighborhood of the point”. These spaces can be defined by decay conditions on the wavelet coefficients as follows, see [5].

If the  $\psi_{j,k} = 2^{j/2}\psi(2^jx - k)$  form an orthonormal basis of  $L^2(\mathbf{R})$ , with  $\psi$  in the Schwartz class, we define the wavelet coefficients of  $f$  by

$$c_{j,k} = 2^j \int f(x)\psi(2^jx - k)dx$$

(note that we do not use a  $L^2$  normalization here).

**Definition 1.1** *A distribution  $F$  belongs to the 2-microlocal space  $C_{x_0}^{s,s'}$  if its wavelet coefficients satisfy, for  $|x_0 - k2^{-j}|$  close enough to 0,*

$$\exists C > 0 : |c_{j,k}| \leq C2^{-j(s+s')} (2^{-j} + |k2^{-j} - x_0|)^{-s'}.$$

It follows from this definition that the *2-microlocal domain of  $F$  at  $x_0$* , i.e.  $\{(s, s') : F \in C_{x_0}^{s,s'}\}$ , is convex and  $F \in C_{x_0}^{s,s'} \implies F \in C_{x_0}^{s-\epsilon, s'+\epsilon}, \forall \epsilon > 0$ .

This set is thus completely determined by the function

$$S(\sigma, x_0) = \sup \left\{ s : F \in C_{x_0}^{s, \sigma-s} \right\}.$$

In addition,  $\sigma \rightarrow S(\sigma, x_0)$  is decreasing and concave. In Section 3 we will show that, conversely, all such functions are functions  $S(\sigma, x_0)$  associated with two-microlocal frontiers *for a fixed point*  $x_0$ . Note that this result has been proved independently using a different method by Y.Meyer in [10]. This mathematical result should also lead to improved estimation methods and new tools for signal processing.

The 2-microlocal domains for all values of  $x_0$  clearly give a very rich information which is superfluous in some applications [1]. It is sometimes sufficient to deal with two numbers: the usual Hölder exponent, and another exponent  $\beta(x_0)$  which measures the oscillatory behavior of  $f$  near  $x_0$ , and is defined as follows.

**Definition 1.2** *Let  $F^{(-n)}$  a  $n$ -th order primitive of  $F$ .  $F$  is called a  $(h, \beta)$ -type chirp at  $x_0$  if*

$$\forall n \in \mathbf{N} : F^{(-n)} \in C_{x_0}^{h+n(1+\beta)}.$$

The simplest example of a  $(h, \beta)$  chirp at  $x_0$  is supplied by the function  $|x - x_0|^h \sin(\frac{1}{|x-x_0|^\beta})$ . The interior of the set of points  $(h, \beta)$  such that a function  $f$  is an  $(h, \beta)$  type chirp at  $x_0$  is always a domain of the form  $h < h_0$ ,  $\beta < \beta_0$ ; this property, proved in [7], allows to define the *chirp exponent at  $x_0$*   $\beta(x_0)$  as this real number  $\beta_0$ .

In sharp contrast with the problem of the prescription of the sole Hölder exponent, we show in Section 2 that the couple  $(h(x_0), \beta(x_0))$  must satisfy a very strong a priori requirement: The function  $\beta(x)$  vanishes where it is continuous.

The following inclusions relate the 2-microlocal and Hölder spaces, [5]

$$C_{x_0}^{s, s'} \subset C_{x_0}^s \subset C_{x_0}^{s, -s}, \quad \forall (s + s') > 0. \quad (2)$$

It follows that the chirp exponents of  $f$  at  $x_0$  are  $(h, \beta)$  if and only if the boundary of the 2-microlocal domain of  $f$  at  $x_0$  is above the half-line starting at  $(s, s') = (h, -h)$  of slope  $-(\beta + 1)/\beta$ , and if this property no more holds for larger values of  $h$  or  $\beta$  (see figure 1).




Figure 1: Relation between 2-microlocal domain and chirps exponents

## 2 A condition on chirps exponents

In this section, we show that it is not possible, in general, to prescribe arbitrarily the chirp exponents of a function independently at each point.

**Proposition 2.1** *Let  $F$  be a function whose Hölder exponent  $h(x)$  satisfies*

$$\text{for all } x \quad 0 < h \leq h(x) \leq H < +\infty;$$

*then the chirp exponent  $\beta(x)$  vanishes on a dense set of points.*

**Proof:** Let  $I$  be a non empty interior interval. Assume  $\beta(x) > 0$  for a set  $E \subset I$ ,  $\overline{E} = I$ . Let  $\epsilon > 0$ ; for any  $x \in E$ , there exists a sequence  $(j_n, k_n) \rightarrow (+\infty, x)$  such that

$$|c_{j_n, k_n}| \geq |x - k_n 2^{-j_n}|^{h(x)+\epsilon} \geq |x - k_n 2^{-j_n}|^{H+\epsilon} \quad (3)$$

Moreover

$$\begin{aligned} |x - k_n 2^{-j_n}| &\geq (2^{-j_n})^{\frac{1}{1+\beta(x)}+\epsilon} \\ &\geq (2^{-j_n})^{1+\epsilon}. \end{aligned} \quad (4)$$

Estimate (3) follows from (2); and (4) holds because, if  $\beta \neq 0$ ,  $(c_{j,k})$  is rapidly decreasing as a function of  $2^{-j}$  in any domain  $2^{-j} \geq |x - k2^{-j}|^{1+\beta-\epsilon}$ . Thus,

$$|c_{j_n, k_n}| \geq (2^{-j_n})^{(1+\epsilon)(1+H)} \quad (5)$$

Let  $E_{j_n} = \{k_n 2^{-j_n} \text{ such that (5) holds}\}$  and  $F_j = \bigcup_{j_n > j} E_{j_n}$ . For all  $x \in E$ , the sequence  $(k_n 2^{-j_n})$  constructed above converges to  $x$ . We construct an  $x_0 \in I$  for which the coefficients  $c_{j,k}$  verify (5) in the cone  $|x_0 - k2^{-j}| \leq 2 \cdot 2^{-j}$  in the following way: Let  $c_{j,k}$  verify (5). We choose  $(j', k')$  such that  $j' > j$  and  $k' 2^{-j'} \in [k2^{-j} - 2^{-j}, k2^{-j} + 2^{-j}]$ . This is possible because  $\overline{F_j} = I, \forall j$ . Carrying on, we construct a sequence of points converging to a point  $x_0$  for which  $\beta(x_0) = 0$ .

### 3 2-microlocal prescription at one point

In this section, we deal with the prescription of the function  $\sigma \rightarrow S(\sigma, x_0)$  for a fixed  $x_0$ . The case where  $S(\sigma, x_0)$  is a line can be excluded since for the chirps  $|x - x_0|^h \sin(|x - x_0|^{-\beta})$ ,  $S(\sigma, x_0) = h(x_0) - \beta(x_0)\sigma$ .

**Proposition 3.1** *Let  $S(\sigma)$  be a concave and decreasing function defined on  $\mathbf{R}$ . Assume that  $S(\sigma)$  is not a line. Then, the wavelet coefficients*

$$c_{j,k} = \inf_{\sigma} 2^{-j\sigma} (2^{-j} + |2^{-j}k - x_0|)^{S(\sigma) - \sigma} \quad (6)$$

define a distribution  $F$ , the 2-microlocal frontier at  $x_0$  of which is  $S(\sigma)$ .

**Proof:** The points  $(S(\sigma), \sigma - S(\sigma))$  are clearly contained in the 2-microlocal domain of  $F$ . For the optimality, we will assume that, without loss of generality,  $x_0 = 0$ . For sake of simplicity, let  $k \geq 0$ :

$$\begin{aligned} u_{j,k} &= \ln(2^{-j}(k+1)) \\ v_{j,k} &= \frac{\ln(2^{-j})}{u_{j,k}} - 1 = -\frac{\ln(k+1)}{\ln(2^{-j}(k+1))} \end{aligned}$$

The 2-microlocal condition, for  $k > 0$ , at 0

$$|c_{j,k}| \leq C 2^{-j\sigma} (2^{-j} + k2^{-j})^{S(\sigma) - \sigma}$$

can be written

$$\ln(|c_{j,k}|) \leq \ln(C) + u_{j,k}(\sigma v_{j,k} + S(\sigma)), \quad (7)$$

and (6) becomes

$$\ln(c_{j,k}) = u_{j,k} \cdot \sup_{\sigma} (\sigma v_{j,k} + S(\sigma)) = u_{j,k} \cdot [-S(\cdot)]^*(v_{j,k}) \quad (8)$$

where  $f^*(\sigma)$  denotes the Legendre-Fenchel transform of  $f$ .

Let  $l(\vartheta) = [-S(\cdot)]^*(\vartheta)$ . As  $(-S(\cdot))$  is a continuous convex function,  $l^*(\sigma) = -S(\sigma)$ , and thus

$$\forall \sigma, \forall \eta > 0, \exists \vartheta_{\sigma, \eta} \quad l(\vartheta_{\sigma, \eta}) \leq \sigma \cdot \vartheta_{\sigma, \eta} + S(\sigma) + \eta. \quad (9)$$

Moreover,  $\vartheta_{\sigma, \eta} > 0$ , as  $S(\cdot)$  is decreasing.

Suppose now that there exist  $\sigma$  and  $\epsilon$  such that  $F \in C_0^{S(\sigma)+\epsilon, \sigma-S(\sigma)-\epsilon}$ . Relations (7) and (8) imply that

$$\exists C > 0, \forall (j, k) : \quad u_{j,k} l(v_{j,k}) \leq \ln(C) + u_{j,k} (\sigma \cdot v_{j,k} + S(\sigma) + \epsilon),$$

which can be rewritten

$$l(v_{j,k}) \geq \frac{\ln(C)}{u_{j,k}} + \sigma \cdot v_{j,k} + S(\sigma) + \epsilon. \quad (10)$$

Let  $\vartheta_{\sigma, \epsilon/2}$  be defined as above. Since  $\vartheta_{\sigma, \epsilon/2} > 0$ , there exist  $(j(n), k(n))$  such that

$$\lim_{n \rightarrow \infty} (u_{j(n), k(n)}, v_{j(n), k(n)}) = (-\infty, \vartheta_{\sigma, \epsilon/2}). \quad (11)$$

As  $S(\cdot)$  is not a line,  $[-S(\cdot)]^*$  is continuous on the right or on the left at each point of its definition domain. Thus, applying (10) for this sequence and taking the limit given by (11),

$$l(\vartheta_{\sigma, \epsilon/2}) \geq \sigma \cdot \vartheta_{\sigma, \epsilon/2} + S(\sigma) + \epsilon$$

which contradicts (9).

**N.B.** While the above local construction can obviously be made around a finite number of points, it follows from Section 2 that the 2-microlocal domain cannot be prescribed everywhere.



## References

- [1] A. Arneodo, E. Bacry, S. Jaffard, and J. F. Muzy. Oscillating singularities on Cantor sets: a grand-canonical multifractal formalism. *J. Statist. Phys.*, 87(1-2):179–209, 1997.
- [2] J-M Bony. Second microlocalization and propagation of singularities for semilinear hyperbolic equations. In “Hyperbolic equations and related topics” (Katata/Kyoto, 1984), pages 11–49. Academic Press, Boston, MA, 1986.
- [3] K. Daoudi, J. Lévy Véhel, and Y. Meyer. Construction of continuous functions with prescribed local regularity. *To appear in Journal of Constructive Approximation*.
- [4] K. Daoudi and J. Lévy Véhel. Speech Signal Modeling Based on Local Regularity Analysis. IASTED/IEEE International Conference on Signal and Image Processing (SIP’95). Las Vegas, November 20-23, 95. .
- [5] S. Jaffard. Pointwise smoothness, two-microlocalization and wavelet coefficients. *Publ. Mat.*, 35(1):155–168, 1991. Conference on Mathematical Analysis (El Escorial, 1989).
- [6] S. Jaffard. Functions with prescribed Hölder exponent. *Appl. Comput. Harmon. Anal.*, 2(4):400–401, 1995.
- [7] S. Jaffard. Lacunary wavelet series. *Preprint*, 1997.
- [8] S. Jaffard and Y. Meyer. Wavelet methods for pointwise regularity and local oscillations of functions. *Mem. Amer. Math. Soc.*, 123 (587), 1996.
- [9] J. Lévy-Véhel. Fractal approaches in signal processing. In “Fractal Geometry and Analysis, The Mandelbrot Festschrift”, C.J.G. Evertsz, H.-O. Peitgen & R.F. Voss Editors, World Scientific, 1996.
- [10] Y. Meyer. Wavelets, vibrations and scalings. Presses de l’université de Montreal. AMS CRM Series, 1997. *To appear*.