



HAL
open science

Specimens: "most of" generic NPs in a contextually flexible type theory

Christian Retoré

► **To cite this version:**

Christian Retoré. Specimens: "most of" generic NPs in a contextually flexible type theory. Genius III, Dec 2011, Paris, France. inria-00637326v1

HAL Id: inria-00637326

<https://inria.hal.science/inria-00637326v1>

Submitted on 4 Nov 2011 (v1), last revised 6 Nov 2011 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SPECIMENS: “MOST OF” GENERIC NPS
IN A CONTEXTUALLY FLEXIBLE TYPE THEORY

Christian Retoré (LaBRI & INRIA, Université de Bordeaux)

Overview This paper proposes to compute the meanings associated to sentences with generic NPs corresponding to the *most of* generalized quantifier. We call these generics *specimens* and they resemble stereotypes or prototypes in lexical semantics. The *meanings* are viewed as logical formulae that can be thereafter interpreted in your favorite models.

We rather depart from the dominant Fregean single untyped universe and go for type theory with hints from Hilbert ε calculus [8, 3] and from medieval philosophy see e.g. [6]. Our type theoretic analysis bears some resemblance with on going work in lexical semantics. [2, 4]

Our model also applies to classical examples involving a class (or a generic element of this class) which is provided by the context. An outcome of this study is that, in the minimalism-contextualism debate, see e.g. [5], if one adopts a type theoretical view, terms encode the purely semantic meaning component while their typing is pragmatically determined.

Terms for universal and specimen generics Here are two examples from the web. The first one involves a universal generic element and the second one a *specimen*:

- (1) The AKC notes that any dog may bite [...]
- (2) The Brits love France.

As Hilbert calculus shows, quantifiers, classes and generic elements are closely related. He introduced $\tau x. A$, an element such that $\forall x. A(x)$ is equivalent to $A(\tau x. A(x))$: A holds for every object if and only if it holds for this element $\tau x. A(x)$, i.e. it is the universal generic associated with A .¹ This view is rather confidential. Some exceptions are the work on definite NPs with ι choice function (in particular by von Heusinger see e.g. [9]) and [1] on generalized quantifiers.

Here we suggest to associate to any property A its specimen written $\angle x. A$. Intuitively, it enjoys all the properties that are true of *most of* A . Although this paper remains on the “syntactic side of semantics”, inhabited with logical formulae, let us give a few hints on how to interpret specimens and their possible reference in models.

Properties of $\angle x. A$ are the ones that are true of *most of* the A ² there can be no contradiction, since when P holds of the specimen, $\neg P$ does not. – in Hilbertian terms it is more like a τ than like an ε . For scalar values we prefer to have relations rather than functions: indeed the specimen has not, for instance, a single height but the relation *height*(*spec*, x) is true whenever x is in some interval (think of baby height charts).

As far as proofs are concerned, we know some situations which enables to assert that P holds of $\angle x. A$: when

¹There is the dual existential generic $\varepsilon x.A$ which satisfies $A(\varepsilon x.A(x)) \equiv A(\tau x.\neg A(x))$.

²Observe that we do not fix a precise ratio much larger than a half. Indeed, it is a vague quantifier. However *most of* as opposed to what is commonly said, is not a matter of cardinality but of measure: for instance in maths books it is said that most of number are not primes meaning $\lim_{n \rightarrow +\infty} \pi(n) \rightarrow 0$.

the universal quantifier holds, when all the *most of* properties are true of it,... But, as expected, we do not know any complete set of rules. We also know it can be refuted when there are only a small minority of A enjoys P , or when there is another property Q disjoint from P and true of $\angle x. A$.

The specimen can be foreseen in ancient and medieval logic: the predication on object as member of some class, formal generic elements with a given ontological class, essential and accidental properties ... In particular, predicates that apply to several unrelated classes were distinguished from “homogenous” predicates that apply to (the generic element of) a class, – as in Abu’l Barakat or Dun Scott. [6]

A flexible typed calculus with a specimen operator

As in Montague semantics we assume that a lexicon associates typed λ -terms with each word, and we start from a syntactic analysis (saying what applies to what). The logical formula depicting the meaning is obtained by applying lexical λ -terms one to another, according to the syntax. In addition to this montagovian term depicting argumental structure each word is also provided with a finite number of λ -terms which are optionally used to convert the type when needed. For instance the lexicon provide for some human entries a term which convert them into vehicles when needed, e.g if a VP like “*is parked up for the night*” is applied to “*Nic*”.

Instead of simply typed λ -calculus we use second order λ -calculus, namely Girard system F (1971), see e.g. [7]. Base types are constant types (the usual ones of TY_n , \mathbf{t} , \mathbf{e}_i , lots of entity types), or variable types, α , β , ... When T_1 and T_2 are types, so is $T_1 \rightarrow T_2$ and when T is a type and α is a type variable, $\Pi\alpha. T$ is a type as well – α usually appears in T but not necessarily.

As opposed to other type theories e.g. (I)TT, the system is conceptually and formally extremely simple, quite powerful,... and paradox free. Term building operations include the ones of simply typed λ -calculus:

\boxed{vc} Constants (resp. variables) of a given type T are terms: $c : T$ (resp. $x : T$).

$\boxed{e\lambda}$ If u is a term of type $T_1 \rightarrow T_2$ and v is a term of type T_1 , then $u(v)$ is a term of type T_2 .

$\boxed{i\lambda}$ If u is a term of type T_2 and x a variable of type T_1 , then $\lambda x. u$ is a term of type $T_1 \rightarrow T_2$.

These usual operations are completed by quite similar operations handling quantification over all types:

$\boxed{e\Lambda}$ If u is a term of type $\Pi\alpha. U$ and T is a type, then $u\{T\}$ is a (specialized) term of type $U[\alpha := T]$

$\boxed{i\Lambda}$ If u is a term of type T and if there is no occurrence of the type variable α in the type of any free variable (u works uniformly for every type α), then $\Lambda\alpha.u$ is a term of type $\Pi\alpha. T$ (that’s the universal view of u).

Remember usual beta-reduction is $(\lambda x^T. u)t^T \rightsquigarrow u[x := t]$ Here, beta-reduction for types and Λ works just the same: $(\Lambda\alpha. u)\{T\} \rightsquigarrow u[\alpha := T]$.

In F, instead of having a constant \forall_α of type $(\alpha \rightarrow \mathbf{t}) \rightarrow \mathbf{t}$ for every type α over which we would like to quantify we shall have one constant \forall of type $\Pi\alpha. (\alpha \rightarrow \mathbf{t}) \rightarrow \mathbf{t}$ that

will be applied to T to obtain the quantifier over the type T : $\forall\{human\}(\lambda x^{human.mortal^{human \rightarrow t}}(x))$.

We introduce a constant \triangleleft of type $\Pi\alpha$. α mapping each property to its specimen. When applied to a type T , this constant \triangleleft yields the element $\triangleleft\{T\}$ of type T which is assumed to be the specimen of T ($\triangleleft\{T\}$ is the F term for $\triangleleft x.T$ when types and properties are identified): it is to be interpreted consequently when interpreting the resulting formula. We could also use the type raised version, mapping each property A to the average element of type A as some did for the choice function.

Computing the readings: semantic terms and contextual typing It is easily seen that our model will provide the right formula for the example (2): $love(\triangleleft\{brits\}, France)$ It resembles the ι choice function, apart that it selects an element about which we can assert properties but which does not exist *stricto sensu*, as medieval universals, Hilbert's $\tau x.A$, etc.

We actually started our reflexion on such generics from classical examples in the minimalism-contextualism debate. Such statements can be both true and false depending on the class in which the object is considered, which is provided by the context.

(3) Carlotta is tall.

If Carlotta is a two year old girl it can be both true ("My daughter is tall and thin for a 2 year old.") and false ("My two-year-old can't get his own cup [...] because he can't reach, [...]") depending on her class – her type in our type theoretic framework.

We noticed that the specimen notion together with the flexibility of F typing succeed to capture this phenomenon. Many of optional λ -terms encode the ontological relations and in the case of a two-year old girl like Carlotta, she can be viewed as a child, and also as a female human being, as a human being etc.

Here are the constants and the useful lexicon entries:

$height : \Pi\alpha. (\alpha \rightarrow float \rightarrow t)$

$< : float \rightarrow float \rightarrow t$

Carlotta $Carlotta : 2yoGirl$ (constant)

$h : 2yoGirl \rightarrow human$ (optional λ -term)

tall $\Lambda\alpha. \lambda x^{\alpha \vee \{float\}} \lambda h_s^{float \vee \{float\}} \lambda h^{float}$

$height\{\alpha\}(\triangleleft\{\alpha\}, h_s) \wedge height\{\alpha\}(x, h) \Rightarrow h_s < h$

type of tall: $\Pi\alpha. \alpha \rightarrow t$

The constant $height$ is a relation between members of a type and numbers ($float$) which are compared with $<$. The entry for $tall$ applies to any type T (second order is quite important here as well) and to a term u of type T . It says that the object u is taller than any possible height of the specimen of this class T .

If we do not use any optional λ -term, we apply $tall$ to the type $2yoGirl$, and to the constant $Carlotta^{2yoGirl}$ we get the reading where Carlotta is taller than the maximal height of the $2yoGirl$ specimen (think again of baby height charts). This is likely to be interpreted as true.

But if we apply $tall$ to the $human$ type, we cannot apply the result to the constant $Carlotta^{2yoGirl}$. But we can apply the $h : 2yoGirl \rightarrow human$ (optional λ -term) to the constant $Carlotta^{2yoGirl}$ and proceed: using the type $human$ since $h(Carlotta)$ is of type $human$. We thus obtain the formula

meaning that Carlotta is tall as a human being, which is unlikely to be interpreted as true.

The semantic machinery produces every possible reading and the context intervenes as a preference for some optional transformation(s). It should be discussed whether there is one or several natural types for an object. Our model can handle any solution: a single natural type, several privileged types,... — quite often, such ontological or metaphysical questions spontaneously pop up when dealing with the organization of the concepts in the lexicon.

Conclusion We applied the F typed λ -calculus to derive semantic readings in the presence of "most of" generics, that we call *specimens*. Our treatment also helps to determine the border between semantics and pragmatics: the term calculus models the semantics, while the typing flexibility of F represents the possible context adaptation.

Thanks This work owes a lot to Sarah-Jane Conrad (*meaning. ch, Sprachphilosophie, Universität Bern*). Indeed, her talk and our discussions initiated in Cerisy on the debate between contextualism and semantic minimalism, lead us to a new connection between logical semantics and type theory, here applied to generic elements.

References

- [1] V. M. Abrusci and C. Retoré. Quantification in ordinary language: from a critic of set-theoretic approaches to a proof-theoretic proposal. In P. Schröder-Heister, editor, *14th Congress of Logic, Methodology and Philosophy of Sciences*, 2011.
- [2] N. Asher. *Lexical Meaning in context – a web of words*. Cambridge University Press, 2011.
- [3] J. Avigad and R. Zach. The epsilon calculus. In E. N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Center for the Study of Language and Information, 2008.
- [4] C. Bassac, B. Mery, and C. Retoré. Towards a Type-Theoretical Account of Lexical Semantics. *Journal of Logic Language and Information*, 19(2):229–245, April 2010.
- [5] S.-J. Conrad. How much context can a language bear? In F. Recanati, editor, *Cerisy Context Conference*, 2011.
- [6] A. de Libera. *La philosophie médiévale*. Presses universitaires de France, 1993.
- [7] J.-Y. Girard, Y. Lafont, and P. Taylor. *Proofs and Types*. Number 7 in Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 1988.
- [8] D. Hilbert. Die logischen grundlagen der mathematik. *Mathematische Annalen*, 88:151–165, 1922.

- [9] K. von Stechow. Alternative semantics for definite nps. In K. Schwabe and S. Winkler, editors, *On Information Structure, Meaning and Form – Generalizations across languages*, volume 100 of *Linguistic Today*, pages 485–508. John Benjamins publishing company, 2007.